Stability and Secretion of Acetylcholinesterase Forms in Skeletal Muscle Cells

Claire Legay,¹ Fawzi A. Mankal,² Jean Massoulié,¹ and Bernard J. Jasmin²

¹Laboratoire de Neurobiologie Moléculaire et Cellulaire, Centre National de la Recherche Scientifique UMR 8544, Ecole Normale Supérieure, 75005 Paris, France, and ²Department of Cellular and Molecular Medicine, Faculty of Medicine, University of Ottawa, Ottawa, Ontario, Canada K1H 8M5

Muscle cells express a distinct splice variant of acetylcholinesterase (AChE_T), but the specific mechanisms governing this restricted expression remain unclear. In these cells, a fraction of AChE subunits is associated with a triple helical collagen, ColQ, each strand of which can recruit a tetramer of AChE_T. In the present study, we examined the expression of the various splice variants of AChE by transfection in the mouse C2C12 myogenic cells *in vitro*, as well as *in vivo* by injecting plasmid DNA directly into tibialis anterior muscles of mice and rats. Surprisingly, we found that transfection with an AChE_H cDNA, generating a glycoposphatidylinositol-anchored enzyme species, produced much more activity than transfection with AChE_T cDNA in both C2C12 cells and *in vivo*. This indicates that the exclusive expression of AChE_T in mature muscle is governed by specific splicing. Interaction of AChE_T subunits

with the complete collagen tail ColQ increased enzyme activity in cultured cells, as well as in muscle fibers *in vivo*. Truncated ColQ subunits, presenting more or less extensive C-terminal deletions, also increased AChE activity and secretion in C2C12 cells, although the triple helix could not form in the case of the larger deletion. This suggests that heteromeric associations are stabilized compared with isolated AChE_T subunits. Coinjections of AChE_T and ColQ resulted in the production and secretion of asymmetric forms, indicating that assembly, processing, and externalization of these molecules can occur outside the junctional region of muscle fibers and hence does not require the specialized junctional Golgi apparatus.

Key words: acetylcholinesterase; collagen tail; C2C12 muscle cells; skeletal muscle; alternative splicing; synaptic proteins; neuromuscular junctions

Acetylcholinesterase (AChE) (EC 3.1.1.7.) is concentrated at neuromuscular junctions. Although both muscle fibers and motoneurons synthesize this enzyme *in vivo* (Brimijoin, 1979; Couraud and Di Giambardino, 1980; Anglister, 1991), there is currently more information available on the pattern of AChE expression in muscle cells.

In mammals, alternative splicing of the AChE gene produces several types of subunits possessing identical catalytic activity (Massoulié et al., 1998). The AChE catalytic domain, which is common to all subunit types, is encoded by exons 2, 3, and 4, whereas alternative C-terminal peptides are generated by splicing of exons 5 [hydrophobic (H)] and 6 [tailed (T)] to yield the AChE_H and AChE_T catalytic subunits, respectively. A third class of mRNA in which the intron after the invariant exon 4 is retained has been identified in *Torpedo*, mouse and rat, but not in man (Sikorav et al., 1988; Legay et al., 1993; Li et al., 1993). This latter mRNA species has been designated as the Readthrough (R) transcript (Taylor and Radic, 1994).

AChE exists as a family of molecular forms that may be

classified as amphiphilic and nonamphiphilic according to their hydrophobic interactions and as homomeric and heteromeric according to their quaternary structure. The homomeric forms include amphiphilic and nonamphiphilic monomers (G₁^a, G₁^{na}), dimers (G₂^a, G₂^{na}), and tetramers (G₄^a, G₄^{na}). Heteromeric membrane-bound and collagen-tailed molecules result from the association of tetramers of AChE_T subunits with an hydrophobic anchoring subunit (P) or with a triple helical collagen tail (ColQ) (Massoulié et al., 1998). In collagen-tailed or asymmetric forms (A₄, A₈, A₁₂), the catalytic subunits are attached to the N terminus of each collagen strand through a proline-rich attachment domain (PRAD) (Bon et al., 1997). In the adult, AChE_H subunits are exclusively expressed in tissues of hematopoietic origin in which they anchored at the cell surface by a glycoposphatidylinositol (GPI). The AChE_T subunit is expressed in both muscle and neuronal cells.

The muscle-specific expression of AChE_T catalytic subunits may result from alternative splicing, mRNA stabilization, and differential post-translational processing. For example, the commitment of myogenic cells to the T splicing pattern occurs early during muscle differentiation (Legay et al., 1995). After fusion of these mononucleated cells into myotubes, T and R mRNAs coexist, but *in vivo* only T transcripts remain expressed after innervation of the muscle. Expression of AChE_T subunits is enhanced during myogenic differentiation in mammals as a result of mRNA stabilization rather than an increase in transcription (Fuentes and Taylor, 1993). Other regulatory events have been implicated because transfection of cDNAs, under the control of nonspecific viral promoters, led to a tissue-specific expression of different types of AChE subunits (Seidman et al., 1995). Post-

Received May 19, 1999; revised July 14, 1999; accepted July 19, 1999.

This work was supported by the Medical Research Council of Canada (MRC), the Centre National de la Recherche Scientifique, the Direction des Forces et de la Prospective, the European Community, and the Association Française contre les Myopathies (AFM). F.A.M. was supported by a Ministry of Ontario Graduate Studentship. B.J.J. is an MRC Scientist. We thank Dr. Roxanne Chan for fruitful discussions and critical reading of this manuscript. Also, we thank Monique Lambergeon and John Lunde for expert technical assistance.

Correspondence should be addressed to Dr. Claire Legay, Laboratoire de Neurobiologie Moléculaire et Cellulaire, Centre National de la Recherche Scientifique, Unité Mixte de Recherche 8544, Ecole Normale Supérieure, 46 rue d'Ulm, 75005 Paris, France.

Copyright © 1999 Society for Neuroscience 0270-6474/99/198252-08\$05.00/0

translational mechanisms may therefore contribute to the preferential expression of AChE_T subunits in muscle cells. In the present study, we examined how several AChE mRNA constructs are expressed after transfection in C2C12 cells in culture.

MATERIALS AND METHODS

Cultures. C2C12 cells were seeded on Matrigel-coated (Collaborative Research, Bedford, MA) 35 mm culture plates and maintained at 37°C in a water-saturated atmosphere containing 5% CO₂. Myoblasts were grown in a proliferation medium consisting of DMEM supplemented with 20% horse serum, 10% fetal calf serum, 292 ng/ml L-glutamine, and 100 U/ml of penicillin–streptomycin. Once the cells reached ~90% confluence, the amount of serum in the media was reduced (5% horse serum) to stimulate differentiation into myotubes.

cDNA constructs. cDNA clones encoding the AChE catalytic subunits were from rat (Legay et al., 1993). They contained no 3' untranslated region and only 12 bp from the mouse 5' untranslated sequence (GTC-CTGGCAGTC) to facilitate translation initiation. The RHT minigene contained the three common exons, followed by the 3' alternative splicing domain (Fig. 1A). The RH construct also contained the three common exons, followed by intron R and the coding region of exon H. Both RH and RHT sequences were obtained by reverse transcription-PCR using total RNA isolated from rat spleen. The cDNA clone encoding the collagen tail (tQ1) was from *Torpedo* (Krejci et al., 1991). Mutagenesis of tQ1 was performed by PCR, using a forward common oligonucleotide and various mutagenized 3' oligonucleotides. The structure of the constructs is shown in Figure 5A. All constructs were inserted at the *Xba*I site of the pEF-BOS vector, under the control of the promoter of elongation factor EF-1 α (Mizushima and Nagata, 1990). Plasmid DNA was prepared using the Qiagen (Chatsworth, CA) mega-prep procedure.

Transfection of cultured cells. At ~50% confluence, C2C12 myoblasts were transfected using the mammalian transfection system–calcium phosphate kit (Promega, Madison, WI) as described previously (Gramolini et al., 1998). Briefly, cells were incubated for 6 hr with 5 μ g of DNA encoding AChE and 5 μ g of DNA encoding the various collagen subunits, together with 3 μ g of a cytomegalovirus-LacZ plasmid (Clontech, Palo Alto, CA), which was included to quantify transfection efficiency. They were then shocked with DMEM containing 15% glycerol for 2 min. After transfection, myoblasts were returned to the proliferation medium (see above) until they reached ~90% confluence. The medium was then replaced by the differentiation medium (see above), and the cells were allowed to differentiate into myotubes for 3 d before analysis.

In vivo gene transfer. These experiments were performed using the tibialis anterior muscles of mouse, as described previously (Chan et al., 1999). Briefly, 25 μ l of a DNA solution containing 2.5 μ g/ml of the appropriate plasmids were injected directly into TA muscles of 4-week-old rats or mice. Seven to 14 d later, injected muscles were excised and rapidly frozen in isopentane precooled with liquid nitrogen. Cryostat tissue sections (10 μ m) were processed histochemically for the simultaneous detection of β -galactosidase and AChE activity (Gramolini et al., 1998).

AChE extraction and biochemical analysis. Three-day-old myotubes were washed twice in PBS and scraped into 400 μ l (per 35 mm plate) of a high-salt detergent buffer containing 10 mM Tris-HCl, pH 7.0, 10 mM EDTA, 1 M NaCl, 1% Triton X-100 or Brij-96, and 1.0 mg/ml of bacitracin and 0.25 mg/ml of aprotinin, as protease inhibitors. Cells were homogenized in a Polytron set at maximum speed, twice for 15 sec. After low-speed centrifugation, supernatants were collected, transferred to fresh tubes, and stored at –80°C for further analysis. For some experiments, cells were first harvested in PBS and immediately frozen at –80°C.

Total AChE activity was determined using the spectrophotometric method of Ellman et al. (1961) as described elsewhere (Jasmin and Gisiger, 1990; Duval et al., 1992). The various AChE molecular forms were separated by velocity sedimentation in 5–20% sucrose gradients in the presence of 1% Brij-96 or Triton X-100. Amphiphilic forms were characterized by the fact that their sedimentation coefficient was lower in the presence of Brij-96 than in the presence of Triton X-100, whereas nonamphiphilic forms were not affected by the detergents (Bon et al., 1991). Treatment of the extract with phosphatidylinositol-phospholipase C (PI-PLC) was performed as described by Legay et al. (1993). The amount of protein present in each sample was determined using the bicinchoninic acid protein assay reagent (Pierce, Rockford, IL).

(A) AChE Catalytic Subunit cDNA Constructs

(B)

Figure 1. *A* is a schematic representation of the various cDNAs encoding AChE splice variants. See Materials and Methods for details. *B* shows AChE activity associated with the myotubes and in the media after transfection of the different constructs. For each construct, a minimum of six distinct cultures (in triplicate) were analyzed. AChE activity is expressed in relation to a cotransfected LacZ construct used to control for transfection efficiency. The values are given as mean $\pm$ SEM.

β -Galactosidase activity was assessed using the β -Galactosidase enzyme assay system (Promega).

To assess secreted AChE activity produced by nontransfected and transfected myotubes, the differentiation media were prepared with horse serum that was pretreated with diisopropyl fluorophosphate (Sigma, St. Louis, MO) (Boudreau-Larivière and Jasmin, 1999).

AChE histochemical staining. Myotubes and cryostat muscle sections were fixed briefly with 4% paraformaldehyde and processed for AChE histochemistry, using the procedure of Karnovsky and Roots (1964).

Immunofluorescence. Myotubes were rinsed in PBS, fixed with 4% paraformaldehyde in PBS for 15 min at room temperature, and washed thoroughly with buffer A (0.5% glycine in PBS) with or without 0.1% Triton X-100. Blockade of nonspecific binding was achieved by incubating the cells in buffer A that contained 5% normal goat serum for 15 min. Cells were then incubated at room temperature with a rabbit anti-rat AChE antibody (A63, diluted 1/400) (Marsh et al., 1984) diluted in buffer A. One hour later, the myotubes were washed with buffer A and incu-

bated for 1 hr at room temperature with Cy3- or fluorescein isothiocyanate-conjugated donkey anti-rabbit IgG antibodies (BioCan, Mississauga, Ontario, Canada). After extensive washing with PBS, the myotubes were mounted in Citifluor (Canterbury, UK) and observed with a Zeiss (Oberkochen, Germany) Axiophot photomicroscope equipped with epifluorescence.

RESULTS

Expression of AChE splice variants in C2C12 myotubes

C2C12 myoblasts were transiently transfected with various constructs encoding the distinct splice variants of AChE catalytic subunits (Fig. 1A). Total AChE activity was assayed in 3-d-old myotubes, and the pattern of AChE molecular forms was analyzed by velocity sedimentation in sucrose gradients. Nontransfected cells contained an extremely low level of AChE activity (Figs. 1B, 2). This activity could not be detected histochemically under the conditions used for visualization of AChE in transfected cells (Fig. 3). In agreement with recent findings (Luo et al., 1998; Boudreau-Larivière and Jasmin, 1999), no asymmetric forms of AChE could be detected in these myotubes. In contrast, total AChE activity produced in transfected cells was significantly higher ($p < 0.05$; Student's t test) and varied according to the particular construct used for transfection. For example, total enzyme activity per dish (cell-associated plus secreted) in cells transfected with constructs containing the H and T cDNAs was more than 30-fold and sixfold higher, respectively, than in nontransfected cells (Figs. 1, 3). In our experiments, transfected C2C12 cells showed no visible morphological difference from controls for any of the constructs that we used (Fig. 3).

After 3 d of differentiation, C2C12 cells transfected with AChE_T cDNA contained mostly monomers (G_1^a), together with low levels of dimers (G_2^a) and tetramers (G_4^{na}), respectively, appearing as a shoulder and a minor peak in the sedimentation profile (Fig. 2). The corresponding media contained G_1^a and G_4^{na} in similar amounts, along with a lower proportion of G_2^a , as revealed by comparison of the molecular form profiles obtained in Triton X-100 versus Brij-96-containing sucrose gradients (data not shown). The minor 13 S component, which was occasionally observed in AChE_T-transfected COS cells (Bon and Massoulié, 1997), was never detected in transfected C2C12 cells. Production of this component seems to be correlated with high levels of AChE_T expression (Simon et al., 1998).

Analysis of the pattern of AChE molecular forms revealed also that C2C12 cells transfected with the H construct produced cell-associated G_2^a and secreted G_2^{na} (Fig. 2). As expected, the sedimentation of G_2^a was shifted after PI-PLC treatment, thereby demonstrating that these cell-associated dimers were GPI-anchored. In addition, these amphiphilic dimers were targeted to membrane of the muscle cells, as revealed by immunofluorescence staining of nonpermeabilized H-transfected C2C12 cells (Fig. 4A). Similarly, histochemical experiments, performed on cryostat sections, showed that the H construct could also be highly expressed along the sarcolemma of muscle fibers after *in vivo* gene transfer (Fig. 4B,C).

We designed two constructs to examine whether alternative splicing is specifically directed toward the T exon in myogenic cells and whether the H exon may be used in the absence of the downstream region. The RH construct contained the common exons 2, 3, and 4 encoding the catalytic domain, followed by intron R and the coding sequence of exon 5; in addition, the RHT construct contained intron 5' and exon 6, i.e., the entire alternative splicing domain (see Fig. 1A). As shown in Figure 1B, total AChE activity was significantly higher ($p < 0.05$; Student's t test)

Figure 2. Representative examples of AChE molecular form profiles seen in myotubes, as well as in the media, after transfection with the different AChE cDNA constructs. For some experiments, the cell extracts were treated with PI-PLC before velocity sedimentation to determine whether the molecular forms contained a GPI-anchor. The data are expressed in arbitrary units.

in cells transfected with the T cDNA compared with that seen in RH-transfected cells, although a larger proportion of AChE activity was secreted with the latter construct (~47% in the case of RH-transfected cells vs ~37% for T-transfected cells). Transfection of the RHT construct into C2C12 cells yielded ~20% more activity than for AChE_T-transfected cells (Fig. 1B).

C2C12 cells transfected with RH constructs produced and secreted G_1^{na} (4.8 S in Triton X-100) together with a lower amount of cell-associated GPI-linked G_2^a . In the presence of Brij-96, the intact G_2^a form sedimented at the same position as the nonamphiphilic G_1^{na} monomers (4.5 S), but PI-PLC treatment produced nonamphiphilic dimers (G_2^{na}) that sedimented at 6.5 S and therefore appeared as a distinct shoulder. Amphiphilic and nonamphiphilic molecules were also characterized by com-

Figure 3. Histochemical analysis of AChE expression in transfected myotubes. *A* shows that transfection of the AChE_T cDNA resulted in modest levels of enzyme activity in a few myotubes, as revealed by histochemical staining using the method of Karnovsky and Roots (1964) (see Materials and Methods). *B* demonstrates that cotransfection of the AChE_T construct with ColQ increased enzyme activity. Consistent with our biochemical analysis, transfection of AChE_H cDNAs (*C*) and cotransfection of AChE_T with Q_N/H_C (*D*) resulted in the most dramatic increase in staining intensity. In parallel experiments, we observed no detectable staining of untransfected cells (data not shown). Note that the morphology of the myotubes is not affected by the expression of these various constructs. Scale bar, 600 μ m.

parison of their sedimentation coefficients in the presence of Triton X-100 and Brij-96 (data not shown).

RHT-transfected cells produced a similar pattern of AChE forms as AChE_T-transfected cells but with a slightly higher proportion of the G₁^a form. In these cells, the sedimentation profiles were clearly not affected by incubating the cell extract with PI-PLC, indicating that the 4.5 S peak does not correspond to GPI-anchored dimers. The sedimentation profiles obtained for RHT-transfected cells did not reveal any detectable nonamphiphilic monomers.

Influence of ColQ subunit on AChE expression

We transfected C2C12 cells with cDNAs encoding AChE_T alone or together with tQ1 (Krejci et al., 1991). Total AChE activity (cell-associated and secreted) was significantly higher ($p < 0.05$; Student's *t* test) in cotransfected cells, indicating that a fraction of the enzyme was stabilized by its association with the collagen subunit (Fig. 5*B*). As expected, these cotransfections resulted in the production of asymmetric forms of AChE. The sedimentation profiles show a major peak at 16 S, corresponding to the A₁₂ form, as well as minor peaks at 14 and ~9–10 S, probably representing A₈ and a mixture of A₄ and G₄ forms (Fig. 6). The G₄ form may be generated by proteolysis of collagen-tailed molecules. Although the sedimentation profiles do not exclude the presence of globular 13 S molecules, their formation is extremely unlikely because they were not observed when AChE_T was expressed alone.

The N-terminal domain of ColQ (tQ_N) (Fig. 5*B*) was sufficient to increase the total activity by threefold to fourfold. Cotransfection with the chimeric protein Q_N/H_C in which the N-terminal of ColQ is linked to a GPI-addition signal peptide also increased by

Figure 4. AChE_H cDNA can be expressed in muscle cells. *A*, Transfected muscle cells in culture stained by immunofluorescence with the A63 antibody directed against AChE. *B*, *C*, Cryostat sections from *in vivo* injected whole muscle stained with the histochemical staining procedure of Karnovsky and Roots (1964). Note that, as expected, the labeling appears associated with the sarcolemma because the H construct contained a GPI addition signal. Scale bars: *A*, *B*, 50 μ m; *C*, 100 μ m.

more than fivefold total AChE activity. Similar to what was observed in COS cells, these cotransfected subunits resulted in the production of GPI-anchored G₄a together with more modest levels of G₁^a and G₂^a (Fig. 6).

To examine whether coexpression with ColQ modified the subcellular distribution of AChE, as well as the total activity, we performed immunofluorescence experiments on cotransfected C2C12 cells using a rabbit polyclonal antibody raised against rat AChE (Marsh et al., 1984). As shown in Figure 7, a strong labeling was seen in permeabilized cells cotransfected with the AChE_T and tQ1 constructs but not in intact nonpermeabilized cells. As shown in Figures 6 and 7, the cells secreted a significant amount of asymmetric forms that did not attach to the cell surface. In agreement with these findings, treatment with collagenase did not release a significant amount of AChE activity from these cells (data not shown). *In vivo*, coinjections of AChE_T and ColQ cDNAs resulted in AChE staining around the cells in extrasynaptic regions (Fig. 8).

The complete ColQ subunits are thus able to recruit the catalytic AChE_T subunits and target them to the cell surface. We

(A) Collagen Subunit cDNA Constructs **(B)**

Figure 5. *A* is a schematic representation of the various cDNA constructs encoding the collagenic subunit, as well as the chimeric Q_N/H_C cDNA. *B* shows AChE activity associated with the myotubes and in the media after transfection and cotransfection of the different constructs. Note that cotransfection with any of the structural subunits resulted in significant increases in AChE enzyme activity ($p < 0.05$). For each construct, a minimum of six distinct cultures (in triplicate) were analyzed. AChE activity is expressed in relation to a cotransfected LacZ construct used to control for transfection efficiency.

tried to define the respective roles of C-terminal peptide motifs in the trimerization and secretion of ColQ. We thus generated three mutants of tQ1, termed tQC2, tQC1, and tQN, as shown in Figure 5*A*, by deletions of the C terminus. In cotransfection experiments, the highest level of total AChE activity (cell-associated and secreted) was observed when the AChE_T cDNA was transfected together with the wild-type collagen subunit, whereas the lowest

Figure 6. Representative examples of AChE molecular form profiles seen in myotubes, as well as in the media, after cotransfection with the different AChE and collagenic subunit cDNA constructs. The data are expressed in arbitrary units. A small peak, which was observed in the culture medium, sedimented slightly faster than the cellular A₁₂ form, most likely because the differentiation media contained trace amounts of collagenase.

amount of activity was detected with cotransfection of the tQN mutant that contains the PRAD domain (Fig. 5*B*).

The tQC2 mutant, which lacks the cysteine-rich domain, nonetheless allowed the formation of the asymmetric forms A₁₂, A₈, and A₄ when cotransfected with the AChE construct (Fig. 6). The AChE sedimentation profiles for both cell-associated and secreted enzymes were identical in tQC2- and tQ1-transfected cells (Fig. 6). With these two constructs, C2C12 cells secreted only asymmetric forms of AChE. In the tQC1 construct, the C-terminal region was nearly completely deleted, except for two cysteines that follow the collagen domain and can form disulfide bonds between the strands in the triple helix. In the tQN construct, the collagen and C-terminal regions were deleted, leaving only the N-terminal domain, including the PRAD. These two constructs did not produce A forms (Fig. 6). Only G₄ was produced and secreted by these cells along with low amounts of G₁

Figure 7. Immunofluorescence experiments using the A63 antibody were performed to localize exogenous expression of AChE in transfected myotubes. AChE expression after transfection with AChE_T alone was restricted to the intracellular compartment of selected myotubes (*A*, *B*). Cotransfection of AChE_T with tQ1 (*C*, *D*) resulted in a much higher expression of AChE activity. Scale bar, 200 μ m.

and G₂, indicating therefore that the truncated tQC1 and tQN subunits recruited most of the G₁ subunits into tetramers.

DISCUSSION

We have examined some of the events that govern expression of AChE_T catalytic subunits in muscle by expressing several AChE cDNA constructs corresponding to various domains of the alternatively spliced sequence in transfected C2C12 cells and in muscle *in vivo*. Transfection of the AChE_T construct (T) into C2C12 cells resulted in the production of molecular forms that were basically identical to those seen previously in COS cells (Bon and Massoulié, 1997), although they were present in slightly different proportions. A small component of G₄^a was occasionally observed in these cell extracts that could represent the association of catalytically active tetramers with endogenous hydrophobic proteins, such as the 20 kDa subunit, in agreement with the existence of membrane-bound tetramers (G₄^a) in muscle (Fernandez et al., 1996). Although the H transcript is only weakly expressed in C2C12 myoblasts and is absent from mature myotubes (Legay et al., 1995), transfected H transcripts were translated and processed efficiently in muscle cells, and the resulting GPI-linked dimers became anchored to the sarcolemma, both in culture and *in vivo*. Therefore, these results indicate that specific expression of AChE_T subunits in muscle originates from the regulation of the splice choice from pre-mRNAs and not from a post-transcriptional elimination of H mRNAs or AChE_H subunits. In the present study, we observed that myogenic C2C12 cells express approximately fivefold higher levels of active AChE_H subunits compared with AChE_T subunits when transfected with the corresponding H and T cDNAs. In contrast, COS cells produced twice as much activity with AChE_T as with AChE_H when transfected with the same constructs (Bon et al., 1997). This indicates

Figure 8. *In vivo* expression of AChE_T constructs coinjected directly into skeletal muscle, along with tQ1 and LacZ. Serial cryostat sections were histochemically processed for β -galactosidase (*A*) and AChE enzyme activity (*B*). Note the coexpression of the two enzymes in some fibers, indicating successful transfection. These pictures were taken in extrajunctional regions of skeletal muscle fibers. No labeling was observed when only AChE_T cDNAs were injected. Also, note that not all LacZ-positive fibers expressed AChE activity, because it is expected that fibers do not necessarily uptake all injected constructs (Jones et al., 1997; Decrouy et al., 1998). Scale bar, 100 μ m.

that the processing and stabilization of the two types of subunits are cell-specific processes.

The restricted secretion of unassembled subunits and the preferential externalization of cysteine-bound dimers from overexpressing human embryonic kidney cells have led to the proposal that the T peptide acts as a retention signal (Velan et al., 1991; Kerem et al., 1993). It is not clear whether the free cysteine of the T peptide plays a crucial role in the ER retention. In this respect, it is noteworthy that, in the present study, we observed an active secretion of AChE_T monomers by transfected C2C12 cells, as reported previously in the case of COS cells (Bon and Massoulié, 1997). Thus, oligomerization is not an absolute prerequisite for externalization of AChE. In C2C12 cells, coexpression of the AChE_T subunit along with various constructs containing the PRAD increased the level of AChE activity, suggesting that the formation of PRAD-linked tetramers stabilized catalytic subunits that would otherwise be degraded. The association of the PRAD with tetramers could therefore mask a degradation signal.

We also examined the splicing choice made by C2C12 cells when transfected with either RH or RHT constructs. When the noncoding region after exon H was absent from the construct (RH), C2C12 cells produced mostly G₁^{na}, corresponding to the translation product of R transcripts, and a small but significant percentage of GPI-linked dimers that represented splicing toward the H exon. When the 3' region downstream of exon H was present (RHT), C2C12 cells produced mostly AChE_T subunits. Based on the present results, it appears therefore that the H acceptor splice site is weaker than the T site in myogenic cells. The absence of the T splicing site and of the upstream intronic sequence did not force the splicing choice toward exon H. Using a human construct similar to our rat RH clone, Seidman et al. (1995) failed to detect the expression of the R and H subunits in

transgenic *Xenopus* muscle. This apparent discrepancy with our results may originate from the species specificity of splicing factors and from intrinsic species-specific nucleotide sequences. For example, transfected mouse blood cells produced R transcripts with the mouse AChE gene but not with the human gene (Li et al., 1993). It is noteworthy that the noncoding region located between the H and T coding sequences contains a 100 nucleotide pyrimidine stretch between the putative branch points and the 3' splice junction, similar in length to those found upstream of exons that are specifically spliced in muscle, such as in the β -tropomyosin gene (Goux-Pelletan et al., 1990). Therefore, the structure of the sequence upstream of the T exon may reinforce the preferential splicing choice in C2C12 cells toward this exon.

Recently, Taylor and colleagues compared the splice choices operated by C2C12 cells that were transfected with a mouse genomic AChE construct or with a construct in which the constitutive introns were deleted (Luo et al., 1998). The latter construct, termed Δ i2–3,3–4, is in fact similar to our RHT construct except that it contains the endogenous AChE promoter and a 5' untranslated region. When transfected into C2C12 cells, this construct produced R, T, and H transcripts in decreasing order, whereas the genomic construct only produced the T transcript. In our experiments, we could not detect any H or R subunits in RHT-transfected cells. In their work, Luo et al. (1998) suggested that the third constitutive intron (intron 3') contains *cis*-elements that influence the downstream splicing. The difference between our results and those of Luo et al. may be related to the fact that they used the mouse gene with its own promoter, whereas we used the rat gene with the EF-1 α promoter or that their transfections were performed 1 d after induction of differentiation, whereas we performed transfections on myoblasts and induced differentiation the next day.

Cysteine- and proline-rich subdomains, which are conserved among vertebrates, have been characterized in the C-terminal sequence of the collagen subunit (Krejci et al., 1991, 1997). Deletion of the more distal cysteine-rich subdomain did not prevent the production of asymmetric forms of AChE, indicating that this region is not essential for the formation of the triple helix. On the other hand, deletions of both cysteine- and proline-rich subdomains did not allow association of the three strands of collagen. Recently, in a genetic analysis of patients presenting a disabling congenital myasthenic syndrome, Ohno and collaborators identified a mutation at a position equivalent to nucleotide 1133 in the *Torpedo* collagen subunit that resulted in the absence of asymmetric forms (Ohno et al., 1998). Altogether, these data indicate that integrity of the entire proline-rich domain is essential for initiating the triple helix assembly, which is known to proceed from the C to the N termini (Prockop and Kivirikko, 1995).

When the AChE_T construct was cotransfected with tQC1, only tetramers were secreted in the medium, suggesting that these tetramers are assembled by a single strand of truncated collagen. It has been shown that single strands of collagen are normally degraded within the cell (Nakai et al., 1992). The PRAD domain must be present in the secreted tetramers (Bon et al., 1997), but we do not know whether the entire collagen domain is conserved in these forms. One hypothesis may be that the association of AChE_T subunit with the N-terminal region of the collagen subunit prevents the intracellular degradation of at least the N-terminal domain of ColQ, thereby allowing secretion of hetero-oligomers. However, the absence of the C-terminal domains in tQC2, tQC1, and tQN mutants decreased the total AChE activity

in cotransfected cells compared with wild-type tQ1, suggesting that the truncated molecules are less stable but more efficiently secreted. Some natural variants lacking part of the ColQ C terminus have been found in rat (Krejci et al., 1997). Our results imply that these ColQ variants would still associate with tetramers and would be secreted at the neuromuscular junction *in vivo*. However, they would not become anchored in the basal lamina because the C-terminal region of ColQ appears necessary for a stable interaction with the extracellular matrix (Donger et al., 1998).

Coinjection experiments in tibialis anterior muscles showed that the presence of ColQ in muscle fibers may direct the secretion of AChE_T subunits, even in extrajunctional domains of the fibers. This suggests that the endplate versus nonendplate specialization of the Golgi apparatus (Jasmin et al., 1989, 1995) and other elements of the secretory pathway (Jasmin et al., 1990) observed previously in skeletal muscle fibers do not affect the capacity of these cells to secrete hetero-oligomers along their entire length. Accordingly, this indicates that the presence of hetero-oligomers in muscle fibers relies exclusively on the expression of AChE_T subunits (Michel et al., 1994; Chan et al., 1999) and ColQ subunits (E. Krejci, C. Legay, S. Thomine, J. Sketelji, and J. Massoulié, unpublished observations).

Finally, it should be noted that only asymmetric but not globular forms of AChE were secreted from C2C12 cells cotransfected with AChE_T and tQ1 or tQC2 constructs. The latter result is, in fact, in agreement with recent findings showing that no AChE activity is detected in the synaptic cleft of ColQ-deficient mice, suggesting that the secreted AChE essentially corresponds to asymmetric forms *in vivo* (Feng et al., 1999).

REFERENCES

- Anglister L (1991) Acetylcholinesterase from the motor nerve terminal accumulates on the synaptic basal lamina of the myofiber. *J Cell Biol* 115:755–764.
- Bon S, Massoulié J (1997) Quaternary associations of acetylcholinesterase. I. Oligomeric associations of T subunits with and without the amino-terminal domain of the collagen tail. *J Biol Chem* 272:3007–3015.
- Bon S, Rosenberry TL, Massoulié J (1991) Amphiphilic, glycoposphatidylinositol-specific phospholipase C (PI-PLC)-insensitive monomers and dimers of acetylcholinesterase. *Cell Mol Neurobiol* 11:157–172.
- Bon S, Coussen F, Massoulié J (1997) Quaternary associations of acetylcholinesterase. II. the polyproline attachment domain of the collagen tail. *J Biol Chem* 272:3016–3021.
- Boudreau-Larivière C, Jasmin BJ (1999) Calcitonin gene-related peptide decreases expression of acetylcholinesterase in mammalian myotubes. *FEBS Lett* 444:22–26.
- Brimijoin S (1979) Axonal transport and subcellular distribution of molecular forms of acetylcholinesterase in rabbit sciatic nerve. *Mol Pharmacol* 15:641–648.
- Chan RY, Boudreau-Larivière C, Angus LM, Mankal FA, Jasmin BJ (1999) An intronic enhancer containing an N-box motif is required for synapse- and tissue-specific expression of the acetylcholinesterase gene in skeletal muscle fibers. *Proc Natl Acad Sci USA* 96:4627–4632.
- Couraud JY, Di Giamberardino L (1980) Axonal transport of the molecular forms of acetylcholinesterase in chick sciatic nerve. *J Neurochem* 35:1053–1066.
- Decrouy A, Renauds J-M, Lunde JA, Dickson G, Jasmin BJ (1998) Mini- and full-length dystrophin induces the recovery of nitric oxide synthase at the sarcolemma of mdx4cv skeletal fibers. *Gene Ther* 5:59–64.
- Donger C, Krejci E, Pou Serradell A, Eymard B, Bon S, Nicole S, Chateau D, Gary F, Fardeau M, Massoulié J, Guicheney P (1998) Mutation in the human acetylcholinesterase-associated collagen gene, ColQ, is responsible for congenital myasthenic syndrome with endplate acetylcholinesterase deficiency (type Ic). *Am J Hum Genet* 63:967–975.

- Duval N, Massoulié J, Bon S (1992) H and T subunits of acetylcholinesterase from *Torpedo*, expressed in COS cells, generate all types of globular forms. *J Cell Biol* 118:641–653.
- Ellman GL, Courtney KD, Andres V, Featherstone RM (1961) A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem Pharmacol* 7:88–95.
- Feng G, Krejci E, Molgo J, Cunningham JM, Massoulié J, Sanes JR (1999) Genetic analysis of collagen Q: role in acetylcholinesterase and butyrylcholinesterase assembly and in synaptic structure and function. *J Cell Biol* 144:1349–1360.
- Fernandez HL, Moreno RD, Inestrosa NC (1996) Tetrameric (G4) acetylcholinesterase: structure, localization, and physiological regulation. *J Neurochem* 66:1335–1346.
- Fuentes ME, Taylor P (1993) Control of acetylcholinesterase gene expression during myogenesis. *Neuron* 10:679–687.
- Goux-Pelletan M, Libri D, d'Aubenton-Carafa Y, Fiszman M, Brody E, Marie J (1990) *In vitro* splicing of mutually exclusive exons from the chicken β -tropomyosin gene: role of the branch point location and very long pyrimidine stretch. *EMBO J* 9:241–249.
- Gramolini AO, Burton EA, Tinsley JM, Ferns MJ, Cartaud A, Cartaud J, Davies KE, Lunde JA, Jasmin BJ (1998) Muscle and neural isoforms of agrin increase utrophin expression in cultured myotubes via a transcriptional regulatory mechanism. *J Biol Chem* 273:736–743.
- Jasmin BJ, Gisiger V (1990) Regulation by exercise of the pool of G4 acetylcholinesterase characterizing fast muscles: opposite effects of running training in antagonist muscles. *J Neurosci* 10:1444–1454.
- Jasmin BJ, Cartaud J, Bornens M, Changeux JP (1989) Golgi apparatus in chick skeletal muscle: changes in its distribution during end plate development and after denervation. *Proc Natl Acad Sci USA* 86:7216–7222.
- Jasmin BJ, Changeux JP, Cartaud J (1990) Compartmentalization of cold-stable and acetylated microtubules in the subsynaptic domain of chick skeletal muscle fibre. *Nature* 344:673–675.
- Jasmin BJ, Antony C, Changeux JP, Cartaud J (1995) Nerve-dependent plasticity of the Golgi complex in skeletal muscle fibres: compartmentalization within the subneural sarcoplasm. *Eur J Neurosci* 7:470–479.
- Jones G, Meier T, Lichtsteiner M, Witzemann W, Sakmann B, Brenner HR (1997) Induction by agrin of ectopic and functional postsynaptic-like membrane in innervated muscle. *Proc Natl Acad Sci USA* 94:2654–2659.
- Karnovsky MJ, Roots L (1964) A direct-coloring thiocholine method for cholinesterases. *J Histochem Cytochem* 12:219–221.
- Kerem A, Kronman C, Bar-Nun S, Shafferman A, Velan B (1993) Interrelations between assembly and secretion of recombinant human acetylcholinesterase. *J Biol Chem* 268:180–184.
- Krejci E, Coussen F, Duval N, Chatel JM, Legay C, Puype M, Vandekerckhove J, Cartaud J, Bon S, Massoulié J (1991) Primary structure of a collagenic tail peptide of *Torpedo* acetylcholinesterase: co-expression with catalytic subunit induces the production of collagen-tailed forms in transfected cells. *EMBO J* 10:1285–1293.
- Krejci E, Thomine S, Boschetti N, Legay C, Sketelj J, Massoulié J (1997) The mammalian gene of acetylcholinesterase-associated collagen. *J Biol Chem* 272:22840–22847.
- Legay C, Bon S, Massoulié J (1993) Expression of a cDNA encoding the glycolipid-anchored form of rat acetylcholinesterase. *FEBS Lett* 315:163–166.
- Legay C, Huchet M, Massoulié J, Changeux JP (1995) Developmental regulation of acetylcholinesterase transcripts in the mouse diaphragm: alternative splicing and focalization. *Eur J Neurosci* 7:1803–1809.
- Li Y, Camp S, Taylor P (1993) Tissue-specific expression and alternative mRNA processing of the mammalian acetylcholinesterase gene. *J Biol Chem* 268:5790–5797.
- Luo ZD, Camp S, Mutero A, Taylor P (1998) Splicing of 5' introns dictates alternative splice selection of acetylcholinesterase pre-mRNA and specific expression during myogenesis. *J Biol Chem* 273:28486–28495.
- Marsh D, Grassi J, Vigny M, Massoulié J (1984) An immunological study of rat acetylcholinesterase: comparison with acetylcholinesterases from other vertebrates. *J Neurochem* 43:204–213.
- Massoulié J, Anselmet A, Bon S, Krejci E, Legay C, Morel N, Simon S (1998) Acetylcholinesterase: C-terminal domains, molecular forms and functional localization. *J Physiol (Paris)* 92:183–190.
- Michel RN, Vu CQ, Tetzlaff W, Jasmin BJ (1994) Neural regulation of acetylcholinesterase mRNAs at mammalian neuromuscular synapses. *J Cell Biol* 127:1061–1069.
- Mizushima S, Nagata S (1990) pEF-BOS, a powerful mammalian expression vector. *Nucleic Acids Res* 18:5322.
- Nakai A, Satoh M, Hirayoshi K, Nagata K (1992) Involvement of the stress protein HSP47 in procollagen processing in the endoplasmic reticulum. *J Cell Biol* 117:903–914.
- Ohno K, Brengman J, Tsujino A, Engel AG (1998) Human endplate acetylcholinesterase deficiency caused by mutations in the collagen-like (ColQ) of the asymmetric enzyme. *Proc Natl Acad Sci USA* 95:9654–9659.
- Prockop DJ, Kivirikko KI (1995) Collagens: molecular biology, diseases, and potentials for therapy. *Annu Rev Biochem* 64:403–434.
- Seidman S, Sternfeld M, Ben Aziz-Aloya R, Timberg R, Kaufer-Nachum D, Soreq H (1995) Synaptic and epidermal accumulations of human acetylcholinesterase are encoded by alternative 3'-terminal exons. *Mol Cell Biol* 15:2993–3002.
- Sikorav JL, Duval N, Anselmet A, Bon S, Krejci E, Legay C, Osterlund M, Reimund B, Massoulié J (1988) Complex alternative splicing of acetylcholinesterase transcripts in *Torpedo* electric organ; primary structure of the precursor of the glycolipid-anchored dimeric form. *EMBO J* 7:2983–2993.
- Simon S, Krejci E, Massoulié J (1998) A four-to-one association between peptide motifs: four C-terminal domains from cholinesterase assemble with one proline-rich attachment domain (PRAD) in the secretory pathway. *EMBO J* 17:6178–6187.
- Taylor P, Radic Z (1994) The cholinesterases: from genes to proteins. *Annu Rev Pharmacol Toxicol* 34:281–320.
- Velan B, Grosfeld H, Kronman C, Leitner M, Gozes Y, Lazar A, Flashner Y, Marcus D, Cohen S, Shafferman A (1991) The effect of elimination of intersubunit disulfide bonds on the activity, assembly, and secretion of recombinant human acetylcholinesterase. Expression of acetylcholinesterase Cys-580→Ala mutant. *J Biol Chem* 266:23977–23984.