Presynaptic Resurgent Na^+ Currents Sculpt the Action Potential Waveform and Increase Firing Reliability at a CNS Nerve Terminal

Jun Hee Kim,¹ Christopher Kushmerick,² and Henrique von Gersdorff¹

¹The Vollum Institute, Oregon Health and Science University, Portland, Oregon 97239, and ²Departamento de Fisiologia e Biofísica, Universidade Federal de Minas Gerais, Belo Horizonte, 31270–901, Minas Gerais, Brazil

Axonal and nerve terminal action potentials often display a depolarizing after potential (DAP). However, the underlying mechanism that generates the DAP, and its impact on firing patterns, are poorly understood at axon terminals. Here, we find that at calyx of Held nerve terminals in the rat auditory brainstem the DAP is blocked by low doses of externally applied TTX or by the internal dialysis of low doses of lidocaine analog QX-314. The DAP is thus generated by a voltage-dependent Na^+ conductance present after the action potential spike. Voltage-clamp recordings from the calyx terminal revealed the expression of a resurgent Na^+ current (I_{NaR}), the amplitude of which increased during early postnatal development. The calyx of Held also expresses a persistent Na^+ current (I_{NaP}), but measurements of calyx I_{NaP} together with computer modeling indicate that the fast deactivation time constant of I_{NaP} minimizes its contribution to the DAP. I_{NaP} is thus neither sufficient nor necessary to generate the calyx DAP, whereas I_{NaR} by itself can generate a prominent DAP. Dialysis of a small peptide fragment of the auxiliary $\beta 4 \text{Na}^+$ channel subunit into immature calyces (postnatal day 5–6) induced an increase in I_{NaR} and a larger DAP amplitude, and enhanced the spike-firing precision and reliability of the calyx terminal. Our results thus suggest that an increase of I_{NaR} during postnatal synaptic maturation is a critical feature that promotes precise and resilient high-frequency firing.

Introduction

To localize sounds, mammals use sub-millisecond differences in the timing of binaural signals (Carr et al., 2001). Preservation of the exact timing of action potential spikes along the ascending auditory pathway is thus thought to be crucial for the rapid and precise pinpointing of sound sources. The calyx of Held synapse is a pivotal element in a brainstem circuit that computes sound localization (Kandler and Friauf, 1993; Grothe, 2003). Fast Na^+ channel inactivation kinetics (Leão et al., 2005), together with high levels of expression of high-threshold (HT) Kv3 potassium channels (Wang and Kaczmarek, 1998; Nakamura and Takahashi, 2007), promote short action potential (AP) waveforms at the calyx terminal, making it well suited to preserve the timing of sensory signals (Dodson and Forsythe, 2004; Kaczmarek et al., 2005). Moreover, a fast and narrow AP waveform allows the mature calyx terminal to fire short spike trains at frequencies up to 1 kHz without failures (Wu and Kelly, 1993; Taschenberger and von Gersdorff, 2000). Sodium currents in the auditory system

may thus have been under evolutionary pressure for fast inactivation and rapid recovery from inactivation (Koishi et al., 2004).

After the action potential spike, recordings from the calyx of Held exhibit a depolarizing after potential (DAP) characterized by a fast upstroke and a slower decay phase (Borst et al., 1995). Previous studies indicate that passive discharge currents regulate the decay phase of the DAP (Borst et al., 1995), whereas active Na^+ and/or K^+ currents influence the amplitude of the DAP (Dodson et al., 2003; Paradiso and Wu, 2009). However, the specific mechanism that generates the calyx DAP has not been established. Moreover, a prominent DAP has also been observed in the action potential waveforms of other CNS nerve terminals (Geiger and Jonas, 2000; Debanne, 2004) and peripheral axons (Barrett and Barrett, 1982; David et al., 1995; Lin, 2008), where the DAP represents a period of subthreshold superexcitability for axons (Bowe et al., 1987). The DAP is therefore of great interest for neuropathies like multiple sclerosis that affect axonal excitability (Baker, 2005).

Pituitary nerve terminals (Ahern et al., 2000) and the calyx of Held (Huang and Trussell, 2008) express a persistent Na^+ current (I_{NaP}), and modeling studies have suggested that I_{NaP} may generate the DAP in some mammalian axons (Burke et al., 2009; McIntyre et al., 2002). In many neurons, non-inactivating I_{NaP} currents are active within the subthreshold region and facilitate bursts of neuronal discharges (Crill, 1996; Kay et al., 1998; Bean, 2007). By contrast, resurgent sodium current (I_{NaR}) is activated during membrane repolarization, following the action potential depolarization that inactivates the transient Na^+ current (I_{NaT}). I_{NaR} is generated by Na^+ channels transiently dwelling in an open state during return to closed states, and its

Received July 30, 2010; accepted Aug. 23, 2010.

J.H.K. was supported by the American Heart Association, C.K. was funded by Fundação de Amparo à Pesquisa do Estado de Minas Gerais and Conselho Nacional de Desenvolvimento Científico e Tecnológico (Brazil), and H.v.G. was supported by an R01 grant from the National Institute on Deafness and Other Communication Disorders. We thank Paul Brehm, Maxim Dobretsov, Ricardo M. Leão, Robert Renden, Geetha Srinivasan, Larry Trussell, and John Williams for discussions.

Correspondence should be addressed to Dr. Henrique von Gersdorff, The Vollum Institute, Oregon Health and Science University, 3181 SW Sam Jackson Park Road, Portland, Oregon 97239. E-mail: vongersd@ohsu.edu.

DOI:10.1523/JNEUROSCI.3982-10.2010

Copyright © 2010 the authors 0270-6474/10/3015479-12\$15.00/0

presence promotes rapid recovery from Na^+ channel inactivation (Raman and Bean, 2001). I_{NaR} thus facilitates neuronal repetitive firing (Raman and Bean, 1997; Afshari et al., 2004). Here, we propose that a developmentally upregulated I_{NaR} improves the calyx of Held's ability to fire without failures at high stimulation frequencies.

Materials and Methods

Brainstem slice preparation. All animal procedures were approved by the institutional animal use committee and followed National Institutes of Health guidelines. Transverse brainstem slices (200 μm thick) were prepared from Sprague Dawley rat pups on postnatal day 5 (P5)–P17. After rapid decapitation, the brainstem of rat pups was quickly removed from the skull and immersed in ice-cold, low-calcium artificial CSF (aCSF) containing the following (in mM): 125 NaCl, 2.5 KCl, 3 MgCl_2 , 0.1 CaCl_2 , 25 glucose, 25 NaHCO_3 , 1.25 NaH_2PO_4 , 0.4 ascorbic acid, 3 myo-inositol, and 2 Na-pyruvate, pH 7.3–7.4 when bubbled with carbogen (95% O_2 , 5% CO_2 ; osmolarity of 310–320 mOsm). After cutting with a vibratome slicer (VT1000, Leica), the slices were transferred to an incubation chamber containing normal aCSF bubbled with carbogen and maintained at 35°C for 30–45 min and thereafter at room temperature (RT). The normal aCSF was the same as the slicing aCSF, but with 1 mM MgCl_2 and 2 mM CaCl_2 .

Electrophysiology. Whole-cell patch-clamp recordings were performed in normal aCSF at RT (22–24°C). Slices were perfused at 2 ml/min and visualized using an infrared differential interference contrast microscope (Axioskop FS1, Zeiss) and a 40 $\times$ water-immersion objective coupled to a 2 $\times$ premagnification (Optovar, Zeiss) and a CCD camera (C73, Sony) with contrast enhancement controller (Hamamatsu). Presynaptic action potential recordings used a pipette solution containing the following (in mM): 130 K-gluconate, 20 KCl, 5 Na_2 -phosphocreatine, 10 HEPES, 4 Mg-ATP, 0.2 or 5.0 EGTA, and 0.3 GTP, pH adjusted to 7.3 with KOH. Presynaptic Na current recordings used a pipette solution containing the following (in mM): 130 Cs-gluconate, 20 CsCl, 10 TEA-Cl, 5 Na_2 -phosphocreatine, 10 HEPES, 4 Mg-ATP, 5 EGTA, and 0.3 GTP, pH adjusted to 7.3 with KOH. The external solution contained 10 mM TEA-Cl and 0.2 mM CdCl_2 to block presynaptic K^+ and Ca^{2+} channels. In some experiments, Alexa 555 (500 μM , Invitrogen) was included in the pipette solution for subsequent morphological analysis by confocal microscopy. Pipettes were pulled from borosilicate glass (World Precision Instruments) with a Sutter P-97 electrode puller (Sutter Instruments) to open tip resistances of 3–4 M Ω . Recordings were continued only if the initial uncompensated series resistance (R_s) was <20 M Ω . For voltage-clamp recordings, R_s was compensated 40–80%, and the average value of R_s after compensation was 7.8 ± 0.57 M Ω ($n = 19$).

For current-clamp recordings, we first established a cell-attached voltage-clamp configuration to set the pipette capacitance compensation (C-fast), and we then broke into the whole-cell configuration and set membrane potential at -80 mV. Once a stable whole-cell voltage-clamp configuration was achieved, we switched to the fast current-clamp configuration of the EPC-9 patch-clamp amplifier (HEKA Elektronik) (see also Taschenberger and von Gersdorff, 2000). Presynaptic resting membrane potentials were adjusted by small amounts of current injection (< -50 pA) to have a resting value near -80 mV (Borst et al., 1995). Recordings were not corrected for a calculated liquid junction potential of ~ 11 mV. The membrane time constant (τ_m) was measured from membrane voltage responses to -50 pA current injections (e.g., see Fig. 2b), and data points were fit by a single exponential function using Igor-Pro software (Wavemetrics). Presynaptic spikes were elicited with a bipolar platinum/iridium electrode (Frederick Haer Company) placed near the midline spanning the afferent fiber tract of the medial nucleus of the trapezoid body (MNTB). An Iso-Flex stimulator delivered 100 μs pulses (< 10 V constant voltage) and was driven by a Master 8 pulse generator (A.M.P.I.) triggered by a Macintosh computer (Apple). Data were filtered at 2.9 kHz and acquired at a 5–10 μs sampling rate, using an EPC-9 amplifier controlled by Pulse 8.4 HEKA software.

Confocal fluorescence microscopy. Slices used for patch-clamp recordings were subsequently fixed in 4% paraformaldehyde for 1 h and permeabilized with 0.5% Triton X-100 for 30 min. Calyces were filled with

Alexa 555 (500 μM , Invitrogen) via the patch pipette (see Fig. 7a; and supplemental Fig. 1, available at www.jneurosci.org as supplemental material). Slices were mounted onto Superfrost slides in photobleaching-protective medium. Stained slices were viewed with a 60 $\times$ oil-immersion objective using a confocal laser-scanning microscope (FluoView 300, Olympus; laser lines at 488 nm for green and 633 nm for red).

Analysis. Data were analyzed off-line using IgorPro software (Wave-metrics). Differences were considered statistically significant when p values were < 0.05 by Student's t test or by one-way ANOVA. Statistical analyses were performed using Prism 4.0 (GraphPad). Means $\pm$ SE are reported, unless otherwise noted. Significance is reported as * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$, using Student's t test or one-way ANOVA test.

Drugs and peptides. The $\beta 4_{\text{peptide}}$ and the scrambled $\beta 4_{\text{peptide}}$, both 95% pure, were synthesized by Gynmed. All other chemicals were obtained from Sigma-Aldrich except for TTX, which was from Tocris Bioscience.

Computational modeling. Model Na^+ currents and action potentials were performed by numerical integration of Hodgkin–Huxley-like equations and the cable equation using the NEURON programming environment (see supplemental materials, available at www.jneurosci.org as supplemental material). The model calyx consisted of two cylindrical compartments: the calyx nerve terminal (capacitance, 12.6 pF) and the axon (diameter, 3 μm ; length, 100 μm ; capacitance, 9.4 pF) (Leão et al., 2005). Conductances included were passive leak conductance, HT and low-threshold (LT) K^+ conductances, and transient, persistent, and resurgent Na^+ currents (Raman and Bean, 2001; Leão et al., 2005; Huang and Trussell, 2008). Auditory neuron HT and LT potassium current models were as previously published (Manis and Marx, 1991; Wang et al., 1998). For the parameters used and further details of the model, see supplemental materials (available at www.jneurosci.org as supplemental material).

Results

Presynaptic DAP and AP timing

Direct recordings from the rat calyx of Held revealed a prominent DAP following a single action potential spike (Fig. 1a) (see also Borst et al., 1995). This presynaptic DAP could be triggered by afferent fiber stimulation in the brainstem midline (Fig. 1a) or by a brief suprathreshold depolarizing current injection in a calyx terminal with a short axon stump (axon length in these cases was measured after filling the calyx with Alexa 555 and varied from 20 to 200 μm , $n = 5$ calyces) (Fig. 1b; supplemental Fig. 1, available at www.jneurosci.org as supplemental material). A long axon, extending back to the midline, was thus not necessary for generating the DAP, suggesting that the DAP was generated near the calyx nerve terminal.

The presynaptic DAP showed a clear rising phase that depolarized the calyx from the downward peak of the fast after-hyperpolarization (fAHP) to near -60 mV (Fig. 1a, inset). However, the membrane potential at the peak of the DAP rarely surpassed -60 mV and thus was always safely below spike threshold. The DAP amplitude measured from the resting potential to the peak of the DAP depended on the resting potential and was larger at more hyperpolarized potentials (supplemental Fig. 2, available at www.jneurosci.org as supplemental material). However, the DAP amplitude measured from the peak of fAHP to the peak of DAP had a fairly invariant amplitude of 7.7 ± 0.2 mV ($n = 22$ calyces) at adjusted resting membrane potentials ranging from -90 to -64 mV (supplemental Fig. 2, available at www.jneurosci.org as supplemental material). The 10 to 90% rise time of the DAP upstroke was dependent on the adjusted resting membrane potential (1.0 ± 0.1 ms at -90 mV, 1.7 ± 0.3 ms at -80 mV, 2.2 ± 0.2 ms at -70 mV, and 2.7 ± 0.3 ms at -65 mV; $n = 3, 3, 6$, and 5, respectively, for P13; $p < 0.01$, one-way ANOVA).

Figure 1. The presynaptic DAP at the calyx of Held. *a*, Afferent fiber stimulation evokes a single presynaptic AP spike followed by a depolarizing DAP in the calyx of Held (P13). Inset, Same trace on an expanded time and voltage scale showing the rise of the DAP and the fAHP. *b*, A brief suprathreshold depolarizing current injection (500 pA for 2 ms) evokes an AP followed by a DAP in a calyx terminal with a severed axon. *c*, A pair of APs in response to afferent fiber stimulation with 4 ms interval (black trace) and then combined with a short hyperpolarizing current injection (-100 pA) to the calyx to remove the DAP (red traces, arrowhead). Note the delay in the second AP (arrow) or failure to fire in the absence of the DAP.

The presynaptic DAP of a first AP influenced the exact timing of a second AP spike. For a pair of APs with a 4 ms interval (corresponding to 250 Hz activity, well within the physiological range of the calyx of Held), a hyperpolarizing current injection of ~ 100 pA removed the DAP and increased the latency of the second AP (latency, measured from the stimulus artifact to the peak of the AP, increased from 0.8 ± 0.06 to 0.9 ± 0.08 ms, $n = 5$, $p < 0.05$) (Fig. 1*c*). In addition, suppressing the DAP also reduced reliability as the failure rate for the second AP was increased. The increase in AP failures may be mediated by the back-propagation of the hyperpolarization down the axon (Paradiso and Wu, 2009). However, we also performed the same experiment on calyces with severed axons of different lengths using a brief depolarizing current injection. Similar to the afferent fiber stimulation case, the complete elimination of the DAP via a step-like hyperpolarizing current injection significantly increased the second AP latency ($n = 5$; data not shown). These results thus indicate the DAP is generated near the calyx terminal, likely at the axonal heminode that contains a large density of Na^+ channels (Leão et al., 2005), and also that the DAP influences the exact timing and reliability of subsequent APs.

Presynaptic DAP: developmental changes

To study the role of DAP during synaptic maturation, we characterized its kinetics at three postnatal ages: shortly after formation of the calyx terminal at P5–P6 (Kandler and Friauf, 1993; Rodríguez-Contreras et al., 2008); just before the onset of hearing at P9–P10; and at P15–P16, when the synapse assumes an adult-like morphology and rat pups respond to sound stimulation (Kandler and Friauf, 1993; Sonntag et al., 2009). Measured from the downward peak of the fAHP to the peak of the DAP (Fig. 2*a*, inset), the amplitude of DAP rise was 2.4 ± 0.4 mV at P5–P6, 6.7 ± 0.7 mV at P9–P10, and 13.8 ± 0.6 mV at P15–P16 ($n = 8$, 9 , and 4 , respectively; $p < 0.001$, one-way ANOVA). For the same ages listed above, the single exponential decay time constants (τ_{decay}) of DAP were 79 ± 3.6 , 30 ± 3.1 , and 15 ± 2.4 ms, respectively ($n = 8$, 9 , and 4 , respectively; $p < 0.001$). This decrease in τ_{decay} closely paralleled a decrease in τ_m s, which were 50.5 ± 3.7 , 22.4 ± 1.5 , and 11.4 ± 1.6 ms, respectively, for the same age groups ($n = 7$, 5 , and 5 , respectively; $p < 0.001$) (Fig. 2*b,c*). The faster membrane time constant in more mature calyces was due in part to reduced input resistance, which was 375 ± 25.1 , 322 ± 29.2 , and 191 ± 16.2 M Ω , respectively, at P5–P6, P9–P10, and

P15–P16 ($n = 7$, 5 , and 5 , respectively; $p < 0.001$). The 10 to 90% rise time of the DAP was also shorter in more mature rats (1.8 ± 0.13 ms in P15–P16 vs 3.0 ± 0.15 ms in P5–P6 calyces; resting potential, -80 mV; $n = 4$; $p < 0.01$). The DAP thus has a significantly faster rise and decay phase for more mature calyces.

What is the mechanism that governs the fAHP peak? We first note that the fAHP becomes progressively larger (more hyperpolarized) with increasing postnatal age (-69 ± 0.7 , -74 ± 0.4 , and -78 ± 0.5 mV; $n = 8$, 9 , and 8 , respectively, for P5–P6, P9–P10, and P15–P16; $p < 0.001$) (Fig. 2*a*). Previous studies indicate that TEA-sensitive Kv3 channels undergo a developmental increase that contributes to shorter AP waveforms and more reliable spiking at high frequencies (Wang and Kaczmarek, 1998; Nakamura and Takahashi, 2007). To test the involvement of Kv3 channels in the generation of the fAHP, we applied 1 mM TEA and found the peak fAHP for P14 calyces was significantly decreased by 7.1 ± 0.3 mV ($n = 3$) from -75 to -67.7 mV (Fig. 2*d*), indicating that Kv3 currents are a major determinant of the fAHP peak (Wang and Kaczmarek, 1998).

How does the DAP affect AP firing during a high-frequency train of spikes? We examined this by recording AP trains in P5–P6 and P15–P16 calyces (Fig. 2*e*). At P5–P6, a long AP train elicited by afferent fiber stimulation at 100 Hz had a large depolarizing plateau, and AP spikes became progressively smaller in amplitude during the train (Fig. 2*e*, arrow). Action potentials eventually failed in the latter part of the train. However, at P15–P16 calyx AP trains had almost no depolarizing plateau and little spike attenuation during the 100 Hz train (Fig. 2*e*). Superposition of the two spike trains at higher time resolution shows that the slow P5–P6 DAP summates to produce a large depolarizing plateau (Fig. 2*e*). This plateau will cause a progressive inactivation of Na^+ channels, which explains the attenuation of the spike amplitudes and the eventual failures in spiking (Klyachko et al., 2001; Leão and von Gersdorff, 2002). These developmental changes in the depolarizing plateau may be caused by an increased expression of Kv2-type channels in more mature calyces (Johnston et al., 2008). The fast decay of the DAP in more mature P15–P16 calyces is thus an important developmental change that helps to avoid DAP summation during a train and the subsequent failures in spiking seen in immature calyces.

Presynaptic Na^+ currents generate the DAP

We next examined the conductances that underlie the DAP in calyx terminals. We found that a low concentration of TTX (20 nM) significantly decreased the amplitude of DAP from 9.0 ± 1.1 to 1.0 ± 0.7 mV with only a slight reduction of AP spike amplitude by 10% ($n = 5$) (Fig. 3*a,f*). For a pair of APs, TTX (20 nM) also had a stronger effect on the second AP latency than the first AP (Fig. 3*a,c*), indicating that the TTX-sensitive DAP is critical to the timing of subsequent AP. To further test that a Na^+ current is the main conductance generating the DAP, we blocked Na^+ channels with 1 μM TTX, which completely inhibits Na^+ spikes, and then injected a large current (1 to 1.5 nA for 1.5 ms) to depolarize the calyx terminal up to $\sim +30$ mV, thus mimicking the peak potential reached during an AP. After the membrane potential reached $+30$ mV, we observed a fast repolarization of

Figure 2. The DAP kinetics during postnatal development. **a**, A presynaptic AP spike and DAP in the calyx of Held from three developmental stages: P6 (red), P10 (blue), and P16 (black). The AP spike was elicited by afferent fiber stimulation. Resting potential was -80 mV (dashed line). Inset, Same traces on an expanded time and voltage scale. Note the changes in rise and decay times of the DAP at different developmental ages. **b**, Membrane potential changes induced by a -50 pA hyperpolarizing current injection in calyces of different ages indicate that input resistance decreases with age. The P15 trace is also shown scaled to the P6 amplitude to facilitate comparison of decay times (gray trace). **c**, Summary of the single exponential τ_{decay} (left) of the DAP and the τ_m (right) for P5–P6, P9–P10, and P15–P16 calyces ($p < 0.001$, one-way ANOVA). **d**, A single AP at P10 calyx in the absence (control; black trace) or in the presence of TEA (1 mM; red trace); 1 mM TEA inhibited the fAHP in the calyx terminals. **e**, Action potential trains in response to 100 Hz afferent fiber stimulation in P5 (red) and P15 calyces (black). The P5 AP train adapted (black arrow). Right, Expanded figure of boxed area of AP trains superimposed. The P5 calyx train of APs showed a large depolarized plateau (large arrowhead) due to the summation of the slow DAPs, and its AP train had spike failures (small vertical arrowheads), whereas the P15 calyx AP train had no failures.

the membrane potential, but no DAP was generated ($n = 3$; P13 calyx) (Fig. 3b, black trace). This rapid decay is likely due to activation of high-threshold Kv3 currents (Dodson and Forsythe, 2004; Kaczmarek et al., 2005). The smaller current injections (0.5 nA), which depolarized the calyx to a peak value < 0 mV, resulted in a much slower decay of membrane potential back to the resting potential (Fig. 3b, red trace). In this case, decay was fitted by a double exponential function consisting of a fast component (1.2 ± 0.2 ms) and a slow component (15 ± 0.7 ms; 17% of amplitude; $n = 3$). The fast component may be due to activation of low-threshold Kv1.1/1.2 channels (Dodson et al., 2003), and the slow component is probably due to the passive discharge of the membrane potential, since this slow τ_{decay} is comparable to the τ_m (11.4 ms at P15–P16) (Fig. 2a–c). Together, these results indicate that a TTX-sensitive Na^+ current, and not depolarization per se, generates the DAP. We conclude that the calyx of Held AP waveform is generated by a primary Na^+ influx event during the upstroke of the spike and a secondary Na^+ influx event that causes the DAP. This secondary Na^+ influx may also have important consequences for energy consumption in axons and nerve terminals (Alle et al., 2009; Carter and Bean, 2009).

To test whether local Na^+ spike after currents close to the calyx and its axonal heminode play a role in DAP generation, we recorded using a low concentration of the permanently charged lidocaine analog QX-314 in the pipette solution (0.2–0.3 mM). Shortly after whole-cell break-in, calyx spikes were followed by a DAP with a fast rising phase (Fig. 3c, black trace). During dialysis of QX-314, the amplitude of the DAP decreased markedly. After

5 min of QX-314 dialysis, the DAP, measured from the fAHP to the DAP peak, decreased from 8.0 ± 1.1 to 0.2 ± 0.1 mV ($n = 8$) (Fig. 3c,f), while the amplitude of the AP was only slightly decreased to $94 \pm 6\%$ ($n = 8$). In addition, the internal application of QX-314 significantly hyperpolarized the calyx of Held resting membrane potential by -4 mV (from -80 ± 0.6 to -84 ± 0.7 mV; $n = 4$; $p < 0.01$), suggesting a tonic Na^+ channel activation at rest (Huang and Trussell, 2008). Moreover, this inhibition of the DAP with QX-314 was reversible. Patching the same calyx terminal a second time using a pipette without QX-314 restored the original DAP amplitude, and the adjusted resting membrane potential moved back to the original -80 mV. In addition, a second spike, which had failed to fire with QX-314 pretreatment, was revived after washing out of QX-314 ($n = 3$) (Fig. 3d). Na^+ currents close to the calyx terminal thus generate the DAP.

By contrast, application of the Na^+ channel activator veratridine (1 μM) significantly increased the DAP amplitude from 6 ± 1.8 to 14 ± 1.3 mV and slowed the decay of the DAP (Fig. 3e,f) ($n = 3$; $p < 0.01$). The progressive increase in the DAP amplitude with veratridine eventually resulted in repetitive spontaneous spiking in response to a single afferent fiber stimulation (Fig. 3e, blue trace). Veratridine increases the open time of Na^+ channels, slows their inactivation rate (Garber and Miller, 1987), and shifts the activation curve to the left (Ulbricht, 1998). This suggests that the decay rate of the DAP may be influenced by the inactivation rate of the underlying Na^+ channels that generate the DAP. Furthermore, an abnormally large DAP amplitude produces aber-

Figure 3. Presynaptic Na⁺ current generates the rising phase of the DAP. *a*, Spikes evoked by afferent fiber stimulation in the absence (control; black trace) and presence of external TTX (20 nM, gray traces during onset of application and red traces end of application). For a pair of APs, with 4 ms stimulus interval, removing the DAP (arrowhead) delayed the latency to the peak of the second AP (red arrow). *b*, After the complete inhibition of voltage-dependent Na⁺ currents with 1 μM TTX, brief current injections [1.5 ms duration, 0.5 nA (red trace) or 1 nA (black trace)] depolarized the calyx to 0 mV (red) or +30 mV (black). Note the absence of a DAP after the larger spike, whereas the smaller spike has a monotonically decaying DAP with no hump. *c*, P13 calyx AP recorded with a pipette solution containing a low concentration of QX-314 (0.2 mM), a Na⁺ channel blocker. The black trace was obtained within a few seconds after break-in, and the red trace was recorded after 7 min of whole-cell recording. After 3 min of QX-314 dialysis, the rising phase of the DAP was significantly inhibited, although spike amplitude was little changed (gray trace). *d*, The calyx of Held was dialyzed for 5 min with QX-314 (0.3 mM), and the stunted rise of the DAP was recorded (red trace). The recording pipette was then removed, and the same calyx was patched again using normal internal pipette solution without QX-314. The normal rise and amplitude of the AP spike and DAP recovered after 7 min of dialysis, which removed QX-314 from the calyx through the patch pipette. Note the broader AP spike with QX-314 (inset). The removal of QX-314 also recovered the second AP, which failed to fire in the presence of QX-314. *e*, Veratridine (an activator of Na⁺ channels; 1 μM) increased significantly the amplitude of the DAP after 1 min (red trace) and elicited repetitive spontaneous spiking in response to a single afferent fiber shock (blue trace; recorded 10 s after red trace). *f*, Summary of DAP amplitude in the presence of TTX (20–50 nM, $n = 5$), QX-314 (0.2–0.3 mM, $n = 7$), and veratridine (1 μM, blue triangles, $n = 3$). Data were analyzed with paired t test (** $p < 0.01$, *** $p < 0.001$).

rant spiking and results in hyperexcitability at the calyx of Held (Dodson and Forsythe, 2004).

I_{NaR} and I_{NaP} at the calyx of Held

To identify the specific type of Na⁺ current responsible for the DAP, we first performed voltage-clamp recordings of calyx Na⁺ currents. A step-depolarization from the holding potential of −90 to +30 mV generated a large and rapidly inactivating I_{NaT} (Leão et al., 2005). Upon repolarization to potentials ranging from −20 to −70 mV, we observed a voltage- and time-dependent current with biophysical properties similar to the I_{NaR} present in Purkinje cells, cerebellar granule cells, vestibular neurons, and MNTB principal cells (Fig. 4*a,b*) (Raman and Bean, 1997; Leão et al., 2006; Magistretti et al., 2006; Gittis and du Lac, 2008). The current–voltage (I–V) relationship for I_{NaR} exhibited a peak of -408 ± 63 pA at ~ -30 mV in P9–P12 calyces ($n = 16$; no liquid-junction potential correction) (Fig. 4*c*). The decay kinetics of I_{NaR} was voltage dependent, with a single exponential τ_{decay} ranging from 5 to 40 ms (Fig. 4*b,d*). In addition, we also observed a I_{NaP} that remained after the decay of I_{NaR} (Fig. 4*a*) (Huang and Trussell, 2008). We also obtained outside-out patches from the calyx terminal that contained well clamped I_{NaT} s ($n = 3$; data not shown) (Leão et al., 2005), but these did not show I_{NaR} or I_{NaP} currents (see also Scott et al., 2010).

Tetrodotoxin at high concentration (1 μM) blocked all three currents described above (I_{NaT} , I_{NaR} , and I_{NaP}), confirming their identity as Na⁺ currents (Fig. 5*a,b*). In contrast, internal QX-314 (0.3 mM) blocked the persistent and resurgent currents but spared 60% of the transient current after 10 min of dialysis of the calyx (supplemental Fig. 3*a*, available at www.jneurosci.org as supplemental material). Bath application of veratridine (5 μM) increased the peak amplitude of I_{NaR} from -273 ± 44 pA to -400 ± 62 pA ($n = 4$) and slowed the inactivation decay of I_{NaR} , whereas the peak of I_{NaT} was not significantly changed (-6.0 ± 0.5 nA versus -5.6 ± 0.8 nA; $n = 4$; $p > 0.1$; paired t test) (supplemental Fig. 3*b*, available at www.jneurosci.org as supplemental material). Low concentrations of TTX (50 nM) also blocked the I_{NaR} and I_{NaP} of more mature P15–P16 calyces ($n = 3$) (supplemental Fig. 4, available at www.jneurosci.org as supplemental material). Together, these results indicate that the calyx I_{NaR} and/or I_{NaP} , both of which are blocked by low doses of TTX and QX-314, and are enhanced by veratridine, underlie the upstroke phase of the DAP.

Developmental increase of calyx I_{NaR}

The rising phase of the DAP became significantly faster from P5 to P15 (Fig. 2*a*), a period when Na⁺ currents undergo developmental changes at the calyx of Held (Leão et al., 2005). To help distinguish the individual contributions of I_{NaR} and I_{NaP} in gen-

erating the DAP, we examined changes in I_{NaR} and I_{NaP} amplitudes during the first two postnatal weeks. We found that I_{NaR} increased in amplitude during the developmental period (Fig. 5*a,b*). The I-V curve of I_{NaR} showed an inverted-bell shape at P5–P6 and P10–P12, and a peak value at -30 mV (Fig. 5*c*). Immature calyces (P5–P6) exhibited significantly smaller inward peak I_{NaR} measured at -30 mV (-86 ± 10.6 pA, $n = 3$) than P10–P12 calyces (-313 ± 71 pA, $n = 5$; $p < 0.02$, Student's t test). An I_{NaP} remained as a noninactivating current after full decay of I_{NaR} , and it slightly increased from -335 ± 103 pA at P5–P6 ($n = 4$) to -533 ± 88 pA at P10–P12, but this increase was not significant (I_{NaP} measured at -10 mV; $n = 5$; $p = 0.38$, Student's t test) (Fig. 5*c*). I_{NaR} and I_{NaP} had different voltage dependences, since I_{NaR} and I_{NaP} displayed two distinct peaks in their I-V curves (at ~ -30 and -10 mV, respectively) (Fig. 5*c,d*). We thus conclude that I_{NaR} and I_{NaP} are distinct sodium currents. Furthermore, the faster rise of DAP amplitudes observed with more mature calyces correlates well with the large increase in I_{NaR} during development.

Computer modeling of the calyx DAP

The pharmacological compounds available to us are not selective for I_{NaR} or I_{NaP} . To obtain further insight into the Na^+ conductance that generates the DAP, we thus performed computer simulations to test for the specific role of persistent and resurgent Na^+ currents. Conductances included in our modeling were passive (leak) conductance; HT and LT potassium conductances; and transient, persistent, and resurgent sodium currents (Fig. 6, black lines) (Dodson and Forsythe, 2004; Kaczmarek et al., 2005). In this manner, we could investigate the effect of eliminating persistent current (Fig. 6, blue lines) or eliminating resurgent current (Fig. 6, red lines) on the DAP. Full details of the model and its parameters are given in Materials and Methods and in supplemental material (available at www.jneurosci.org as supplemental material).

Model sodium currents (Fig. 6*a*) were generated by turning off the other conductances and performing linear leak subtraction to remove capacitive currents. The simulated voltage-clamp protocol to elicit mixed resurgent and persistent Na^+ current was the same as that used for the recordings shown in Figure 4*a*: a step depolarization to $+30$ mV for 5 ms that generated transient Na^+ current followed by a repolarizing step from -20 to -70 mV in 10 mV steps. Simulated I_{NaR} showed a peak in the bell-shaped I-V relationship near -40 mV, and the decay of I_{NaR} also showed the same voltage dependency as the experimental data (simulation results not shown). Importantly, the rate of inactivation of the transient Na^+ current was significantly faster with an I_{NaR} present in the simulations (Fig. 6*a*), as expected from the extra effect of the inactivation binding particle used to generate I_{NaR} (Raman and Bean, 2001). To generate an I-V relationship for the persistent Na^+ current, we used a slow voltage ramp (20 mV/s) from a potential of -90 mV. This I-V relationship showed current activation at -70 mV (no liquid junction potential correction) and a

Figure 4. I_{NaR} in the calyx of Held. *a*, I_{NaT} and I_{NaR} were elicited in a P10 calyx terminal by a 10 ms step depolarization to $+30$ mV, followed by step repolarizations to -70 , -60 , -50 (gray), -40 (green), -30 (blue), and -20 mV (red) as indicated in the voltage-clamp protocol (top). Note that after the decay of I_{NaR} , I_{NaP} remains activated. *b*, Individual current traces for repolarizations to -70 and -30 mV are shown together with single exponential decay fits. The I_{NaT} peaks have been truncated. Note the fast deactivating tail currents (arrowheads) before the full activation of I_{NaR} . *c*, The current-voltage dependence (I-V curve) for peak I_{NaR} for P9–P12 calyces. The peak I_{NaR} was measured from the end of I_{NaT} to the peak of I_{NaR} . *d*, The τ_{decay} of I_{NaR} as a function of test membrane potentials. I_{NaR} decayed more slowly at more depolarized potentials.

peak near -20 to -10 mV, as we observed experimentally (Fig. 5*d*; simulation results not shown).

Action potentials were generated in the computer model by current injection (4 nA for 0.2 ms) into the distal end of the axon. Four Na^+ current scenarios were studied: normal resurgent and persistent current (black traces), resurgent current without persistent current (blue traces), no resurgent current with normal persistent current (solid red traces), and no resurgent current with two times higher persistent current (broken green traces). Without I_{NaR} , we obtained a calyx action potential with no DAP (Fig. 6*b*). In addition, note that the half-width of the action potential was broader without I_{NaR} (as is also the case experimentally with 0.3 mM QX-314) (Fig. 3*c,d*). Removal of I_{NaP} had very little effect on the DAP (blue traces), and even increasing I_{NaP} two times over experimentally recorded values did not rescue the DAP in the absence of I_{NaR} . These simulations thus capture the main basic features of our previous experimental results, and they show that I_{NaP} by itself does not generate the DAP. They thus represent a proof of principle for our main hypothesis that I_{NaR} generates the DAP.

We next explored the effect of varying the amplitude of I_{NaP} (from 2.4 to 5.6 times the measured values) on the fAHP or DAP (Fig. 6*c*), while the time constant of deactivation was kept constant at 0.24 ms (close to the value measured in P16 calyces). In the presence of I_{NaR} , a 2.4-fold to 3.2-fold higher density of I_{NaP} increased the peak amplitude of the DAP (measured relative to the fAHP) only slightly, whereas a 4.2-fold to 5.6-fold increase in I_{NaP} was necessary to produce a significantly larger DAP with a prolonged time course (Fig. 6*c*). However, without I_{NaR} we observed that increasing persistent current removed the fAHP but did not produce a rebounding DAP “hump,” as observed experimentally (Fig. 6*d*). We conclude that persistent current in the calyx of Held is unlikely to generate the DAP, although it may modulate its amplitude and kinetics.

Figure 5. The calyx I_{NaR} increases during postnatal development. **a**, I_{NaT} , I_{NaP} , and I_{NaR} were evoked at a P5 calyx by a short depolarization to +30 mV followed by a step repolarization to -30 mV. TTX (1 μM) completely inhibited I_{NaT} , I_{NaP} , and I_{NaR} in the calyx of Held. The bottom trace is the TTX-sensitive current obtained by subtraction. The large I_{NaT} peak has been truncated. **b**, I_{NaT} , I_{NaP} , and I_{NaR} at a P10 calyx. TTX (1 μM) completely inhibited I_{NaP} , I_{NaR} , and I_{NaT} . **c**, I-V curve for peak I_{NaR} at P5–P6 (black) and P10–P12 calyxes (red). All sodium currents were leak subtracted (p/5) and TTX subtracted. There is a significant increase in I_{NaR} between the two age groups at the peak of I_{NaR} near -30 mV. **d**, I-V curve for the I_{NaP} for P5–P6 calyxes (black) and P10–P12 calyxes (red). There is a slight increase in I_{NaP} between the two different age groups at the peak of I_{NaP} near -10 mV, but this was not statistically significant ($p = 0.38$, Student's t test). Membrane potentials were not corrected for a 10 mV liquid junction potential. Persistent sodium current thus starts to activate at about -80 mV and may contribute to the resting membrane potential.

Figure 6. Computer model of the role of I_{NaR} and I_{NaP} in the generation of the DAP. **a**, Model Na^+ currents in four scenarios: control (black); after removal of persistent current (blue); after removal of resurgent current (red); and after removal of resurgent current with two times higher persistent current (green). **b**, Model action potentials in the same scenarios as described in **a**. Note that the DAP is normal without persistent current, but the DAP is absent without resurgent current. The action potential waveform is also broader without resurgent current. **c**, Effect of four different densities of persistent current on the DAP (relative to that measured in the calyx of Held). Black trace, $2.4 \times I_{NaP}$; blue trace, $3.2 \times I_{NaP}$; red trace, $4.2 \times I_{NaP}$; and green trace, $5.6 \times I_{NaP}$. Top, Persistent Na^+ tail currents evoked by steps from 0 to -80 mV. Bottom, Resulting DAP waveforms. The deactivation time constant (τ) was fixed at $\tau = 0.24$ ms. **d**, Same as in **c** except without resurgent Na^+ current. No DAP hump was generated. **e**, The deactivation time constant of the persistent Na^+ currents from 0 to -80 mV was varied (black, 0.2 ms; blue, 0.3 ms; red, 0.5 ms; green, 0.9 ms; and purple, 1.6 ms). **f**, Model action potentials with only persistent Na^+ currents in the same scenarios as described in **e**. **g**, **h**, Representative experimental data traces of deactivation of calyx Na^+ currents from -30 to -80 mV in the P16 calyx (**g**) and P10 calyx (**h**). Single exponential fits are shown to the data (red). The more mature P16 calyx had a faster τ .

The persistent current used in the simulation shown in Figure 6*a–d* has fast deactivation kinetics (0.24 ms at -80 mV). However, we reasoned that an I_{NaP} with sufficiently slow deactivation kinetics could contribute to the DAP since during membrane repolarization following an AP the increased driving force for Na^+ could result in a depolarizing drive similar to that caused by resurgent Na^+ current. To explore this, we tested several different models in which the time constant for deactivation at -80 mV was varied from 0.2 to 1.6 ms (Fig. 6*e*). As the kinetics of persistent current gating were slowed, the fAHP was reduced, and for very slow time constants approaching 1 ms a DAP with a hump was indeed generated (Fig. 6*f*). However, in the calyx of Held, the experimentally measured persistent Na^+ current deactivated with a fast time constant of 0.27 ± 0.07 ms (P16; $n = 4$) or 0.6 ± 0.1 ms (P10; $n = 3$) (Fig. 6*g–h*). We therefore conclude that the I_{NaP} in the calyx of Held deactivates too rapidly to generate a significant DAP.

$\beta 4$ subunit peptides induce I_{NaR} in immature calyces

To further test for a causal relationship between I_{NaR} and the DAP, we next used a portion of the $\beta 4$ sodium channel subunit to induce I_{NaR} in immature calyces. A leading model for how I_{NaR} can arise is through the voltage-dependent binding of a blocking particle (Raman and Bean, 2001), which may originate from the $\beta 4$ subunit to the $Na_v1.6$ channel (Grieco et al., 2005). According to this model, when the $\beta 4$ subunit particle binds it both blocks current flow and also prevents the normal “classical” inactivation process from occurring. On repolarization, and dissociation of the $\beta 4$ subunit particle, the channel conducts again until normal deactivation or closure occurs. Interestingly, very high levels of $\beta 4$ subunit mRNA are found throughout the adult rat brainstem (Yu et al., 2003), but the level of $\beta 4$ subunit protein in the developing rat calyx of Held is unknown. If protein levels follow mRNA, an increased expression of $\beta 4$ subunits during early postnatal development could explain the observed developmental increases in I_{NaR} and the DAP amplitude. Unfortunately, due to a lack of reliable antibodies, we have not been able to confirm a developmental increase in protein levels of the $\beta 4$ subunit in rat MNTB.

We thus tested whether a peptide that mimics the putative action of $\beta 4$ subunits on Na^+ channels could generate I_{NaR} in immature P5–P6 calyces, which have little resurgent Na^+ current (Fig. 7*a,b*). In recordings without peptide, the size of I_{NaR} in P5 calyces was relatively small and stable during whole-cell recording (control) (Fig. 7*c*). When $100 \mu M$ $\beta 4_{peptide}$ (KKLITFILKKTREK, Grieco et al., 2005) (Fig. 7*b*, red trace) was included in the pipette solution, I_{NaR} increased 10-fold, from 23 ± 8 pA at 2 min to 227 ± 53 pA after 8 min of intracellular dialysis of the peptide ($n = 5$, $p < 0.001$) (Fig. 7*c*); whereas, the increase of I_{NaP} with $\beta 4_{peptide}$ was not significant (from 32 ± 11 pA to 53 ± 20 pA after 8 min of dialysis; $n = 5$, $p = 0.11$). In the presence of the $\beta 4_{peptide}$,

Figure 7. Induction of I_{NaR} by dialysis of a $\beta 4$ peptide into P5–P6 calyces of Held. *a*, An immature calyx terminal (P5) expresses a very small amplitude I_{NaR} which was recorded with repolarizing step pulses from $+30$ to -20 mV. Inset, The P5 calyx of Held (arrowhead) and attached axon were filled with Alexa-555 via the patch pipette for morphological identification. *b*, I_{NaR} was recorded with control internal solution (black trace) or with an internal solution that contained $\beta 4_{peptide}$ ($100 \mu M$; amino acid sequence from the $\beta 4$ subunit of Na^+ channels, red trace) or with a peptide with a scrambled sequence of $\beta 4_{peptide}$ (blue trace). Inset, Expanded traces of the I_{NaR} shown from the shadowed box. *c*, The peak amplitude of I_{NaR} at -30 mV progressively increased during whole-cell dialysis of $\beta 4_{peptide}$ ($100 \mu M$; colored dots) in five individual P5–P6 calyx recordings. However, no significant change in peak amplitude of I_{NaR} was detected with control internal solution (black dots; $n = 5$). *d*, Current–voltage relationship of I_{NaR} recorded without $\beta 4_{peptide}$ (control, black; $n = 5$), with $\beta 4_{peptide}$ (red; $n = 8$), or with scrambled $\beta 4_{peptide}$ (blue; $n = 5$) for P5–P6 calyx terminals.

the I–V relation of I_{NaR} was bell shaped and similar to that observed in more mature calyces ($n = 8$) (Fig. 7*d*).

To confirm that the effects produced by the $\beta 4_{peptide}$ depended on the specific amino acid sequence, we tested the effect of a scrambled peptide on I_{NaR} in which charge was conserved but the clusters of hydrophobic and positively charged residues were disrupted ($\beta 4_{scramble}$: KIKIRFKTKTLELK) (Grieco et al., 2005) (Fig. 7*b,d*, blue trace). In the presence of $\beta 4_{scramble}$, the I_{NaR} did not significantly increase during 15 min of dialysis, and its I–V relationship remained similar to control ($n = 5$) (Fig. 7*d*). These data indicate that the $\beta 4_{peptide}$ can significantly increase I_{NaR} in immature P5–P6 calyces of Held.

$\beta 4$ subunit peptides increase the DAP amplitude and firing reliability

Next, we investigated the effect of $\beta 4_{peptide}$ dialysis on the AP waveform of immature P5–P6 calyces. AP spikes were evoked once every 10 s by afferent fiber stimulation, and $\beta 4_{peptide}$ ($100 \mu M$) was dialyzed into the calyx via the patch pipette. The amplitude of the DAP in the presence of $\beta 4_{peptide}$ was significantly increased from 2.2 ± 0.2 to 4.0 ± 0.5 mV after 10–15 min of dialysis (P5–P6; $n = 12$, $p < 0.01$; DAP amplitude was measured from the fAHP) (Fig. 8*a1*). The potential reached at the peak of the DAP measured from the resting membrane potential was also increased by intracellular $\beta 4_{peptide}$ dialysis from -67 ± 0.4 to -63 ± 0.8 mV ($n = 12$, $p < 0.01$) (see supplemental Fig. 5*a*, available at www.jneurosci.org as supplemental material). The $\beta 4_{peptide}$ depolarized the average calyx adjusted resting membrane potential from -80 ± 0.4 to -76 ± 0.5 mV and signifi-

cantly shortened the spike latency from 1.6 ± 0.2 to 1.3 ± 0.1 ms ($n = 12$, $p < 0.001$) (Fig. 8b). In addition, the 10 to 90% rise time of the spike was reduced from 328 ± 16.9 to 273 ± 14.3 μ s ($n = 9$; $p < 0.001$) (Fig. 8b). Similar effects on AP spike latency and rise time are also observed in postsynaptic MNTB neurons after the dynamic clamp incorporation of I_{NaR} into normal MNTB neurons (Leão et al., 2006).

The increase in DAP amplitude by the β_4 peptide was not due to membrane depolarization because depolarizing the calyx by 4 mV via current injection in control calyces lacking β_4 peptide had no effect on the amplitude of DAP ($n = 3$) (Fig. 8a2). In three recordings, we also observed the diffusion of the β_4 peptide using tetramethylrhodamine-tagged β_4 peptide and found that it diffused along the axon for at least 100–200 μ m in the 7–10 min after break-in to whole-cell recording mode (supplemental Fig. 5b, available at www.jneurosci.org as supplemental material). This indicates that the β_4 peptide reaches the whole extent of the heminode region of the axon where $Na_v1.6$ channels are primarily located (Leão et al., 2005). To test whether the effects of the

β_4 peptide are due to changes in K^+ current induced by the peptide, we recorded calyx K^+ currents with pipettes containing β_4 peptide. Insertion of β_4 peptide into P5 calyces had no significant effect on the amplitude of voltage-dependent K^+ currents after 10 min of β_4 peptide dialysis (K^+ currents at 0 mV were 1.9 ± 0.46 nA in control and 1.8 ± 0.59 nA with β_4 peptide; $n = 3$, $p > 0.05$; data not shown). Thus, we conclude that an increase of I_{NaR} by β_4 peptide dialysis enhances the amplitude of the DAP and reduces the AP latency.

We next examined the consequences of increased I_{NaR} and DAP for high-frequency firing by recording from P5–P6 calyces before and after β_4 peptide dialysis. To reduce calyx-to-calyx variability (Sonntag et al., 2009), we first recorded spike trains (100–200 Hz) in control without β_4 peptide. We then removed the patch pipette and repatched the same calyx with a pipette containing 100 μ M β_4 peptide. P5 calyces fired trains of 50 spikes without failure at 100 Hz but showed several failures at 200 Hz. The success rate (number of spikes per total of 50 stimuli) was smaller in control than after dialysis with β_4 peptide. At 133 Hz (stimulus interval, 7.5 ms), several control P5 calyces (6 of 10) exhibited spike failures during the late phase of the stimulus train. After dialysis with β_4 peptide, these same calyces showed no failures at 133 Hz (Fig. 8c, black and red traces, respectively). The success rate at 133 Hz increased from $92 \pm 3.5\%$ to 100% with β_4 peptide ($n = 10$; $p < 0.05$).

To control for the possibility that the depolarization of ~ 4 mV caused by the peptide could have effects on the calyx firing fidelity, we performed a control experiment without β_4 peptide. Most P5 calyces could not follow 200 Hz stimulus trains

for 250 ms, showing a large number of AP failures (e.g., 25 failures of 50 stimuli) (Fig. 8d). Constant current injection (~ 30 pA) depolarized the calyx resting potential from -80 to -70 mV; nevertheless, the calyx still had the same number of AP failures at 200 Hz. In the same calyx, however, the dialysis of 100 μ M β_4 peptide reduced AP failures in the early phase of the train. The success rate increased from $59 \pm 3.3\%$ to $71 \pm 3.9\%$ with β_4 peptide at 200 Hz ($n = 10$; $p < 0.01$) (Fig. 8d). Together, our results suggest that I_{NaR} induced by β_4 peptide increases peak DAP amplitude, reduces the spike latency, and leads to an increase in firing reliability at high stimulation frequencies.

Discussion

We have uncovered the presence of a presynaptic I_{NaR} in a mammalian nerve terminal whose action potential waveform has a prominent DAP. Direct calyx of Held recordings show that the I_{NaR} amplitude is upregulated during the first 2 weeks of postnatal development. Introducing a small β_4 peptide into an immature calyx mimicked this developmental increase in I_{NaR} and

expanded the range of reliable firing of the immature terminals. We thus propose that a developmental increase of I_{NaR} in auditory calyx of Held nerve terminals promotes reliable high-frequency firing, a critical feature for encoding signals for sound localization (Yin, 2002; Grothe, 2003).

Local Na^+ currents mediate the presynaptic DAP

The mechanism that generates the DAP at CNS bouton-type nerve terminals and axons has not been clearly established because their small size makes direct recordings very difficult. At the calyx of Held, a local puff application of TTX on the calyx terminal reduced the spike amplitude but did not affect the DAP amplitude measured from the adjusted resting membrane potential, suggesting that the DAP may originate in the axon (Borst et al., 1995). More recently, a local puff application of riluzole, a non-specific blocker of Na^+ channels, reduced the DAP and AP spike amplitude (Paradiso and Wu, 2009). However, riluzole also blocks Ca^{2+} and K^+ channels (Lamanauskas and Nistri, 2008), so the specific mechanism that generates the DAP remained uncertain. Our QX-314 dialysis experiments show that the rising phase of the DAP is caused by local Na^+ current activation because dialysis of 0.3 mM QX-314, a blocker of I_{NaR} and I_{NaP} (supplemental Fig. 3a, available at www.jneurosci.org as supplemental material), reversibly blocked the rising phase of the DAP (Fig. 3c,d). Moreover, a low concentration of TTX (20 nM) also blocked the DAP, although it had a relatively small effect on AP spike amplitude (Fig. 3a). Importantly, the DAP was observed in calyces with short axon stumps after a brief current injection (Fig. 1b; supplemental Fig. 1, available at www.jneurosci.org as supplemental material). So, a long intact axon was not necessary for generating the DAP. Instead, we propose that local Na^+ currents in the calyx pre-terminal heminode region generate the DAP.

DAP is generated predominantly by I_{NaR} in the calyx terminals

Is the DAP generated by I_{NaR} , I_{NaP} , or both types of Na^+ currents? Here, we present three separate pieces of evidence that I_{NaR} is sufficient to produce a DAP. First, our modeling studies suggest that I_{NaP} contributes to the DAP only if its deactivation kinetics is slowed significantly to values near 1.36 ms, whereas we measured an I_{NaP} time constant for deactivation of ~ 0.25 ms in P15–P16 calyces (Fig. 6g). Similarly, adult cerebellar Purkinje cells have an I_{NaP} time constant for deactivation of 0.2 ms (Kay et al., 1998). Second, the rising phase of the DAP became progressively faster and larger during development from P5 to P15 (Fig. 2a). Likewise, peak I_{NaR} amplitudes increased significantly from P5 to P12 (Fig. 5c; supplemental Fig. 4, P15, available at www.jneurosci.org as supplemental material), while the developmental increase in I_{NaP} was less significant (Fig. 5d). A resurgent current of ~ 100 pA at nearly -70 mV (the peak of the fAHP) would depolarize P15–P16 calyces with an input resistance of ~ 150 M Ω by ~ 15 mV. By contrast, the immature P5–P6 calyx has a higher input resistance of ~ 300 M Ω , so a much smaller I_{NaR} of ~ 20 pA could generate a DAP of ~ 6 mV. An increase in I_{NaR} may thus be necessary to overcome the low input resistance of mature calyces, whereas an unchecked increase in I_{NaP} may generate spontaneous firing (Raman and Bean, 1997). Third, dialysis of the $\beta 4$ peptide increased the DAP amplitude in immature P5–P6 calyces (Fig. 8a1; supplemental Fig. 5, available at www.jneurosci.org as supplemental material). Because $\beta 4$ peptide increases significantly I_{NaR} amplitude (Fig. 7b), but does not significantly change the average I_{NaP} amplitude, we suggest that the DAP is caused mainly by I_{NaR} activation. Recently, the $\beta 4$ peptide has been shown to also in-

crease I_{NaP} in transfected cells expressing $Na_v1.1$ channels (Aman et al., 2009). However, the calyx of Held heminode expresses a high density of $Na_v1.6$ channels (Leão et al., 2005), and high levels of $\beta 4$ subunit mRNA are found in adult rat brainstem (Yu et al., 2003). Interestingly, the coexpression of $\beta 4$ subunits with $Na_v1.6$ channels is not sufficient to reconstitute I_{NaR} in transfected cells (Chen et al., 2008). Additional proteins beyond $Na_v1.6 \alpha$ and $\beta 4$ subunits may thus be required for induction of I_{NaR} .

DAP kinetics and repetitive firing

The peak amplitude and time course of the DAP influences the fidelity of repetitive firing at high frequencies (Fig. 2e). Since the passive properties of the axon and its myelin sheath contribute to the kinetics of the DAP at peripheral axons (Barrett and Barrett, 1982; Bowe et al., 1987; David et al., 1995; Lin, 2008), the lack of myelin on immature calyx axons at P5–P6 could contribute to the slower decay of DAP at immature calyces, whereas the faster DAP decay observed in P15–P16 calyces may arise in part from a more compact myelin sheath that promotes a faster axonal τ_m (Fig. 2c). During repetitive firing, DAP summation in the immature calyx causes a large depolarizing plateau that probably leads to a reduction in spike amplitudes, due to an accumulation of Na^+ channels in the inactivated state, and eventual spike failures (Fig. 2e, arrow). Thus, the faster DAP decay kinetics and decrease in membrane time constant of more mature calyces help them to fire spikes at higher frequencies without failures.

What is the mechanism for reduced failure rates with the $\beta 4$ peptide (Fig. 8d)? The $\beta 4$ peptide increases the DAP amplitude (Fig. 8a1), and the DAP helps to avoid AP failures (Fig. 1c). In addition, we propose that the $\beta 4$ peptide increases the availability of Na^+ channels for opening by reducing the number of channels in the inactivated state. Studies at the nodes of Ranvier and calyx of Held indicate that at a resting membrane potential of -80 mV $\sim 20\%$ of the Na^+ channels are inactivated (Schwarz et al., 1995; Nakamura and Takahashi, 2007). A large surplus of Na^+ channels that are available for opening is necessary for the reliable firing of spikes at high frequencies (as shown by studies using low doses of TTX; Madeja, 2000). By blocking the “classical inactivation” pathway for Na^+ channels the $\beta 4$ -peptide may effectively increase the number of channels that can be opened during a stimulus train. Larger Na^+ currents due to increased availability of channels in the calyx during a stimulus train will thus lead to a reduced number of failures, as well as a reduction in the spike latency and a faster AP rise time. The presence of I_{NaR} in calyx terminals thus promotes an acceleration of firing and greater spiking reliability (Fig. 8c,d).

I_{NaR} contributes to faster Na^+ channel inactivation and recovery

Previous studies have suggested various specializations that are acquired during early development that aid the calyx of Held in its remarkable capacity for high-frequency firing (Schneggenburger and Forsythe, 2006). One is the faster rate of Na^+ channel inactivation of the more mature calyx, and another is an increase of Kv3 channel density during development (Leão et al., 2005; Nakamura and Takahashi, 2007). Both will promote a shorter AP waveform (Rudy and McBain, 2001). Indeed, the more mature calyx of Held has a shorter AP waveform and a faster recovery rate from Na^+ channel inactivation than other CNS synaptic boutons (Jackson and Zhang, 1995; Engel and Jonas, 2005). The presence of I_{NaR} in a neuron promotes faster Na^+ channel inactivation and faster recovery from inactivation, and thus facilitates repetitive

firing (Raman and Bean, 1997). Our computer simulations also support a role of I_{NaR} in maintaining AP half-widths short via fast Na^+ channel inactivation (Fig. 6*a,b*). We thus propose that the faster recovery from inactivation of Na^+ channels in the more mature calyx of Held may be due in part to the more copious expression of I_{NaR} .

Functional implications of I_{NaR} for the auditory system

Sound-evoked signals must be conveyed through the ascending auditory pathways at high speed and low jitter, and with great reliability (Carr et al., 2001; Yin, 2002). A short AP waveform is thus critical for the transfer of high-frequency auditory signals. Our computer simulations show that the presence of a relatively large I_{NaR} aids in the production of short APs (Fig. 6*b*). Moreover, I_{NaR} also produces a fast rising DAP that accelerates the firing of a subsequent AP (Fig. 1*c*), and thus the precise timing of APs during a spike train (Fig. 8*c,d*). Postsynaptic MNTB neurons also exhibit a DAP (Johnston et al., 2009), and a developmental increase in I_{NaR} which improves their ability to fire at higher frequencies, and I_{NaR} expression is altered by deafness (Leão et al., 2006). Of course, speed, precision, and reliability of spiking are also augmented through the myelination of axons, which increases during early postnatal development (Vabnick and Shrager, 1998). Resurgent Na^+ currents in nerve terminals and axons may thus be yet another specialization of auditory circuits that promotes fast, precise, and resilient action potential signaling.

References

- Afshari FS, Ptak K, Khaliq ZM, Grieco TM, Slater NT, McCrimmon DR, Raman IM (2004) Resurgent Na^+ currents in four classes of neurons of the cerebellum. *J Neurophysiol* 92:2831–2843.
- Ahern GP, Hsu SF, Klyachko VA, Jackson MB (2000) Induction of persistent sodium current by exogenous and endogenous nitric oxide. *J Biol Chem* 275:28810–28815.
- Alle H, Roth A, Geiger JR (2009) Energy-efficient action potentials in hippocampal mossy fibers. *Science* 325:1405–1408.
- Aman TK, Grieco-Calub TM, Chen C, Rusconi R, Slat EA, Isom LL, Raman IM (2009) Regulation of persistent Na^+ current by interactions between β subunits of voltage-gated Na^+ channels. *J Neurosci* 29:2027–2042.
- Baker MD (2005) Ion currents and axonal oscillators: a possible biophysical basis for positive signs and symptoms in multiple sclerosis. In: *Multiple sclerosis as a neuronal disease* (Waxman SG, ed), pp 131–144. New York: Elsevier Academic.
- Barrett EF, Barrett JN (1982) Intracellular recording from vertebrate myelinated axons: mechanism of the depolarizing afterpotential. *J Physiol* 323:117–144.
- Bean BP (2007) The action potential in mammalian central neurons. *Nat Rev Neurosci* 8:451–465.
- Borst JG, Helmchen F, Sakmann B (1995) Pre- and postsynaptic whole-cell recordings in the medial nucleus of the trapezoid body of the rat. *J Physiol* 489:825–840.
- Bowe CM, Kocsis JD, Waxman SG (1987) The association of the supernormal period and the depolarizing afterpotential in myelinated frog and rat sciatic nerve. *Neuroscience* 21:585–593.
- Burke D, Howells J, Trevillion L, McNulty PA, Jankelowitz SK, Kiernan MC (2009) Threshold behaviour of human axons explored using subthreshold perturbations to membrane potential. *J Physiol* 587:491–504.
- Carr CE, Soares D, Parameshwaran S, Perney T (2001) Evolution and development of time coding systems. *Curr Opin Neurobiol* 11:727–733.
- Carter BC, Bean BP (2009) Sodium entry during action potentials of mammalian neurons: incomplete inactivation and reduced metabolic efficiency in fast-spiking neurons. *Neuron* 64:898–909.
- Chen Y, Yu FH, Sharp EM, Beacham D, Scheuer T, Catterall WA (2008) Functional properties and differential neuromodulation of $\text{Na}_v1.6$ channels. *Mol Cell Neurosci* 38:607–615.
- Crill WE (1996) Persistent sodium current in mammalian central neurons. *Annu Rev Physiol* 58:349–362.
- David G, Modney B, Scappaticci KA, Barrett JN, Barrett EF (1995) Electrical and morphological factors influencing the depolarizing after-potential in rat and lizard myelinated axons. *J Physiol* 489:141–157.
- Debanne D (2004) Information processing in the axon. *Nat Rev Neurosci* 5:304–316.
- Dodson PD, Forsythe ID (2004) Presynaptic K^+ channels: electrifying regulators of synaptic terminal excitability. *Trends Neurosci* 27:210–217.
- Dodson PD, Billups B, Rusznák Z, Szűcs G, Barker MC, Forsythe ID (2003) Presynaptic rat $\text{K}_v1.2$ channels suppress synaptic terminal hyperexcitability following action potential invasion. *J Physiol* 550:27–33.
- Engel D, Jonas P (2005) Presynaptic action potential amplification by voltage-gated Na^+ channels in hippocampal mossy fiber boutons. *Neuron* 45:405–417.
- Garber SS, Miller C (1987) Single Na^+ channels activated by veratridine and batrachotoxin. *J Gen Physiol* 89:459–480.
- Geiger JR, Jonas P (2000) Dynamic control of presynaptic Ca^{2+} inflow by fast-inactivating K^+ channels in hippocampal mossy fiber boutons. *Neuron* 28:927–939.
- Gittis AH, du Lac S (2008) Similar properties of transient, persistent, and resurgent Na currents in GABAergic and non-GABAergic vestibular nucleus neurons. *J Neurophysiol* 99:2060–2065.
- Grieco TM, Malhotra JD, Chen C, Isom LL, Raman IM (2005) Open-channel block by the cytoplasmic tail of sodium channel $\beta 4$ as a mechanism for resurgent sodium current. *Neuron* 45:233–244.
- Grothe B (2003) New roles for synaptic inhibition in sound localization. *Nat Rev Neurosci* 4:540–550.
- Huang H, Trussell LO (2008) Control of presynaptic function by a persistent Na^+ current. *Neuron* 60:975–979.
- Jackson MB, Zhang SJ (1995) Action potential propagation and propagation block by GABA in rat posterior pituitary nerve terminals. *J Physiol* 483:597–611.
- Johnston J, Griffin SJ, Baker C, Skrzypiec A, Chernova T, Forsythe ID (2008) Initial segment $\text{Kv}2.2$ channels mediate a slow delayed rectifier and maintain high frequency action potential firing in MNTB neurons. *J Physiol* 586:3493–3509.
- Johnston J, Postlethwaite M, Forsythe ID (2009) The impact of synaptic conductance on action potential waveform: evoking realistic action potentials with a simulated synaptic conductance. *J Neurosci Methods* 183:158–164.
- Kaczmarek LK, Bhattacharjee A, Desai R, Gan L, Song P, von Hehn CA, Whim MD, Yang B (2005) Regulation of the timing of MNTB neurons by short-term and long-term modulation of potassium channels. *Hear Res* 206:133–145.
- Kandler K, Friauf E (1993) Pre- and postnatal development of efferent connections of the cochlear nucleus in the rat. *J Comp Neurol* 328:161–184.
- Kay AR, Sugimori M, Llinás R (1998) Kinetic and stochastic properties of a persistent sodium current in mature guinea pig cerebellar Purkinje cells. *J Neurophysiol* 80:1167–1179.
- Klyachko VA, Ahern GP, Jackson MB (2001) cGMP-mediated facilitation in nerve terminals by enhancement of the spike afterhyperpolarization. *Neuron* 31:1015–1025.
- Koishi R, Xu H, Ren D, Navarro B, Spiller BW, Shi Q, Clapham DE (2004) A superfamily of voltage-gated sodium channels in bacteria. *J Biol Chem* 279:9532–9538.
- Lamanauskas N, Nistri A (2008) Riluzole blocks persistent Na^+ and Ca^{2+} currents and modulates release of glutamate via presynaptic NMDA receptors on neonatal rat hypoglossal motoneurons in vitro. *Eur J Neurosci* 27:2501–2514.
- Leão RM, Von Gersdorff H (2002) Noradrenaline increases high-frequency firing at the calyx of Held synapse during development by inhibiting glutamate release. *J Neurophysiol* 87:2297–2306.
- Leão RM, Kushmerick C, Pinaud R, Renden R, Li GL, Taschenberger H, Spirou G, Levinson SR, von Gersdorff H (2005) Presynaptic Na^+ channels: locus, development, and recovery from inactivation at a high-fidelity synapse. *J Neurosci* 25:3724–3738.
- Leão RN, Naves MM, Leão KE, Walmsley B (2006) Altered sodium currents in auditory neurons of congenitally deaf mice. *Eur J Neurosci* 24:1137–1146.
- Lin JW (2008) Electrophysiological events recorded at presynaptic terminals of the crayfish neuromuscular junction with a voltage indicator. *J Physiol* 586:4935–4950.
- Madeja M (2000) Do neurons have a reserve of sodium channels for the

- generation of action potentials? A study on acutely isolated CA1 neurons from the guinea-pig hippocampus. *Eur J Neurosci* 12:1–7.
- Magistretti J, Castelli L, Forti L, D'Angelo E (2006) Kinetic and functional analysis of transient, persistent, and resurgent sodium currents in rat cerebellar granule cells in situ: an electrophysiological and modeling study. *J Physiol* 573:83–106.
- Manis PB, Marx SO (1991) Outward currents in isolated ventral cochlear nucleus neurons. *J Neurosci* 11:2865–2880.
- McIntyre CC, Richardson AG, Grill WM (2002) Modeling the excitability of mammalian nerve fibers: influence of afterpotentials on the recovery cycle. *J Neurophysiol* 87:995–1006.
- Nakamura Y, Takahashi T (2007) Developmental changes in potassium currents at the rat calyx of Held presynaptic terminal. *J Physiol* 581:1101–1112.
- Paradiso K, Wu LG (2009) Small voltage changes at nerve terminals travel up axons to affect action potential initiation. *Nat Neurosci* 12:541–543.
- Raman IM, Bean BP (1997) Resurgent sodium current and action potential formation in dissociated cerebellar Purkinje neurons. *J Neurosci* 17:4517–4526.
- Raman IM, Bean BP (2001) Inactivation and recovery of sodium currents in cerebellar Purkinje neurons: evidence for two mechanisms. *Biophys J* 80:729–737.
- Rodríguez-Contreras A, van Hoesen JS, Habets RL, Locher H, Borst JG (2008) Dynamic development of the calyx of Held synapse. *Proc Natl Acad Sci U S A* 105:5603–5608.
- Rudy B, McBain CJ (2001) K_v3 channels: voltage-gated K^+ channels designed for high-frequency repetitive firing. *Trends Neurosci* 24:517–526.
- Schneggenburger R, Forsythe ID (2006) The calyx of Held. *Cell Tissue Res* 326:311–337.
- Schwarz JR, Reid G, Bostock H (1995) Action potentials and membrane currents in the human node of Ranvier. *Pflügers Arch* 430:283–292.
- Scott LL, Mathews PJ, Golding NL (2010) Perisomatic voltage-gated sodium channels actively maintain linear synaptic integration in principal neurons of the medial superior olive. *J Neurosci* 30:2039–2050.
- Sonntag M, Englitz B, Kopp-Scheinpflug C, Rübsamen R (2009) Early postnatal development of spontaneous and acoustically evoked discharge activity of principal cells of the MNTB: an *in vivo* study in mice. *J Neurosci* 29:9510–9520.
- Taschenberger H, von Gersdorff H (2000) Fine-tuning an auditory synapse for speed and fidelity: developmental changes in presynaptic waveform, EPSC kinetics, and synaptic plasticity. *J Neurosci* 20:9162–9173.
- Ulbricht W (1998) Effects of veratridine on sodium currents and fluxes. *Rev Physiol Biochem Pharmacol* 133:1–54.
- Vabnick I, Shrager P (1998) Ion channel redistribution and function during development of the myelinated axon. *J Neurobiol* 37:80–96.
- Wang LY, Kaczmarek LK (1998) High-frequency firing helps replenish the readily releasable pool of synaptic vesicles. *Nature* 394:384–388.
- Wang LY, Gan L, Forsythe ID, Kaczmarek LK (1998) Contribution of the $K_v3.1$ potassium channel to high-frequency firing in mouse auditory neurons. *J Physiol* 509:183–194.
- Wu SH, Kelly JB (1993) Response of neurons in the lateral superior olive and medial nucleus of the trapezoid body to repetitive stimulation: intracellular and extracellular recordings from mouse brain slice. *Hear Res* 68:189–201.
- Yin TCT (2002) Neural mechanisms of encoding binaural localization cues in the auditory brainstem. In: *Integrative functions in the mammalian auditory pathway* (Oertel D, Fay RR, Popper AN, eds), pp 99–159. New York: Springer-Verlag.
- Yu FH, Westenbroek RE, Silos-Santiago I, McCormick KA, Lawson D, Ge P, Ferriera H, Lilly J, DiStefano PS, Catterall WA, Scheuer T, Curtis R (2003) Sodium channel β_4 , a new disulfide-linked auxiliary subunit with similarity to β_2 . *J Neurosci* 23:7577–7585.