

Selection of Alternative G_i-mediated Signaling Pathways at the Dopamine D₂ Receptor by Protein Kinase C

Vincenzo Di Marzo,^a Daniel Vial, Pierre Sokoloff, Jean-Charles Schwartz, and Daniele Piomelli

Unité de Neurobiologie et Pharmacologie de l'INSERM, Paris 75014, France

Evidence indicates that a single membrane receptor subtype may be responsible for the generation of multiple intracellular signals, but mechanisms allowing for the selection of a specific effector pathway have not yet been documented. In neurons and other cells, the stimulation of dopamine D₂ receptors produces, via G-protein activation, a spectrum of intracellular responses including inhibition of adenylyl cyclase activity, modulation of K⁺ currents, and potentiation of Ca²⁺-evoked arachidonic acid (AA) release. In this study, we report that, in Chinese hamster ovary cells, stimulation of protein kinase C (PKC) directs the preferential coupling of transfected D₂ receptors from inhibition of adenylyl cyclase to potentiation of AA release, two responses mediated by G_i. The switch between these two signaling systems is accompanied by marked changes in their GTP sensitivities, indicating that it may result from the phosphorylation of component(s) of the receptor–G-protein complex. Brain PKC activity is enhanced by neurotransmitters and by neuronal depolarization. Thus, the ability of this protein kinase to remodel signaling pathways at the D₂ receptor may regulate these G_i-mediated responses in an activity-dependent manner, and represent a novel form of synaptic plasticity.

[Key words: Chinese hamster ovary fibroblasts, transfected cells, second messengers, arachidonic acid, phospholipase A₂, cAMP, adenylyl cyclase]

Communication between neurons requires that the actions of neurotransmitters on their target cells be both specific and modifiable, to allow for plastic changes to occur in response to environmental demands. To achieve specificity, many neurotransmitters interact with distinct subtypes of membrane receptors coupled to specific intracellular second messengers. For example, dopamine, a major modulatory transmitter in the brain, exerts its neural actions through stimulation of at least five receptor subtypes, discretely distributed throughout the CNS, and functionally coupled either to G_s-mediated activation of adenylyl cyclase (“D₁ type”) or to G_i-mediated cyclase inhibition (“D₂ type”) (Sibley and Monsma, 1992; Schwartz et al., 1992).

Increasing evidence indicates, however, that a single receptor subtype may be linked to the formation of multiple, parallel intracellular signals. This spectrum of heterogeneous effectors may be produced through activation of one or several transducing G-proteins (for review, see Spiegel, 1992). For example, in anterior pituitary cells, two different G-proteins, G_o and G_{i3}, link D₂ receptors to modulation of Ca²⁺ and K⁺ channel opening, respectively (Lledo et al., 1992). Likewise, in Chinese hamster ovary (CHO) cells transfected with D₂ receptor cDNA [CHO(D₂)], D₂ receptor stimulation causes both inhibition of adenylyl cyclase activity and facilitation of evoked release of arachidonic acid (AA) via pertussis toxin-sensitive G_i-proteins (Felder et al., 1991; Kanterman et al., 1991; Piomelli et al., 1991).

The functional redundancy in intracellular signaling underscored by these findings raises several important questions. Are all the signals driven by a single receptor subtype equally operative under all circumstances? Or rather, are these signals subject to cellular mechanisms of integration and regulation? Can the spectrum of second messengers in a cell be remodeled, for example, by changing the functional weight of one pathway with respect to others, to adapt to its varying needs? And if this is the case, could such a remodeling at the second messenger level affect the response to receptor stimulation? Interest in such questions lies in the possibility that, like ion channels and neurosecretory mechanisms, the pathways of transmembrane signaling may represent a potential substrate for plastic changes in neural cells.

Protein kinase C (PKC) serves important regulatory functions in the CNS, where it is thought to participate in the modulation of ion channel activity, receptor desensitization and transmitter release (Nishizuka, 1988). Using CHO(D₂) cells, we have now examined the role played by this protein kinase in controlling parallel intracellular signaling at the D₂ receptor. We report that activation of PKC switches the coupling of D₂ receptors from inhibition of adenylyl cyclase toward facilitation of AA release, possibly by phosphorylating component(s) of the receptor–G_i-protein complex. Because brain PKC activity is stimulated by neurotransmitters and neuronal depolarization (Nishizuka, 1988), the results suggest that remodeling of transmembrane signaling pattern by this protein kinase, demonstrated here in an heterologous expression system, may modify in an activity-dependent manner physiological responses to D₂ receptor stimulation.

Materials and Methods

Cell culture and transfections. CHO cells were transfected with rat D₂₍₄₄₄₎ receptor (also termed D_{2L}) cDNA or with human D₂₍₄₁₅₎ receptor (also termed D_{2S}) cDNA as previously described (Giros et al., 1989). CHO

Received Dec. 11, 1992; revised May 3, 1993; accepted May 20, 1993.

D.P. was supported in part by a Young Investigator Award from the National Alliance for Research on Schizophrenia and Depression (NARSAD), and V.D., by a fellowship from the European Molecular Biology Organization. We thank Dr. J.-A. Girault for critical reading of the manuscript.

Correspondence should be addressed to Daniele Piomelli, Unité de Neurobiologie et Pharmacologie de l'INSERM, 2ter Rue d'Alésia, Paris 75014, France.

^a Present address: Istituto per la Chimica di Molecole d'Interesse Biologico, CNR, via Toiano 6, Arco Felice, Napoli 80072, Italy.

Copyright © 1993 Society for Neuroscience 0270-6474/93/134846-08\$05.00/0

cells expressing rat $D_{2(444)}$ receptor are referred to in the text as CHO(D_2); those expressing $D_{2(415)}$ receptor, as CHO($D_{2(415)}$). Transfected clones were maintained in monolayer culture in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% fetal calf serum (GIBCO) at 37°C in 5% CO₂. CHO(D_2) expressed 1.3×10^5 receptors/cell, and CHO($D_{2(415)}$), 1.5×10^5 receptors/cell.

Receptor binding. CHO(D_2) cells were incubated in DMEM containing 0.25% trypsin (GIBCO) for 5 min at room temperature, and the reaction was stopped by adding 10 ml of supplemented DMEM. Detached cells were collected by centrifugation and washed in 15 mM Tris-HCl, pH 7.4, containing 5 mM MgCl₂ and 10 mM sodium pyrophosphate, to inhibit protein phosphatase activity. After cell disruption, membranes were isolated by centrifugation (50,000 × *g* for 15 min) in the same buffer. Binding was determined in 50 mM Tris-HCl, pH 7.4, containing (mM) NaCl, 120; KCl, 5; CaCl₂, 2; MgCl₂, 2; ascorbic acid, 0.28; and 8-hydroxyquinoline, 0.1. Curves were analyzed by computerized nonlinear regression using a one- or two-site model (Martres et al., 1984).

³H-AA release. Cells (24-well plates) were labeled by incubation with ³H-AA (Amersham; 200–220 Ci/mmol, 0.25 mCi/ml) in DMEM (1 ml) containing 0.2% bovine serum albumin (BSA) for 2 hr at 37°C. To eliminate unincorporated radioactivity, cells were washed with 1 ml of DMEM plus BSA, before incubating them for 30 min at 37°C in 1 ml of DMEM, containing final concentrations of the appropriate drugs. Analysis by high-performance liquid chromatography revealed that, in BSA-containing incubations, free ³H-AA constituted more than 90% of the released radioactivity in both control and stimulated samples (Di Marzo and Piomelli, 1992). To prevent dopamine oxidation, incubations with the drug were carried out in the presence of ascorbic acid (0.1%), which did not affect either basal or stimulated ³H-AA release (not shown). In some experiments, cells were incubated with pertussis toxin (200 ng/ml) for 4 hr at 37°C in 1 ml of DMEM. Incubation with the toxin did not affect ³H-AA labeling of phospholipids (not shown). ³H-AA release was determined by liquid scintillation counting of samples (0.5 ml) of the incubation medium.

cAMP formation. Cells (96-well plates) were incubated for 10 min in 0.15 ml in DMEM containing isobutylmethylxanthine (IBMX) (0.1 mM), a phosphodiesterase inhibitor, plus forskolin (10 μM) and final concentrations of the appropriate drugs. After extraction in 0.1 M HCl (0.1 ml), sonication and neutralization of the tissue extracts, cAMP concentrations were determined using a radioimmunoassay kit (Amersham), following the manufacturer's instructions.

Measurements of intracellular free Ca²⁺ concentration. Cells were suspended by treatment with trypsin (see above), and incubated for 60 min at 37°C of DMEM containing BSA (0.2%) and fura-2 acetoxyethyl ester (1 μM). After washing, cells were resuspended in HEPES-buffered Krebs' solution (containing, in mmol/liter, NaCl, 125; KCl, 5; KH₂PO₄ and MgSO₄, 1.2; CaCl₂, 2; glucose, 6; HEPES/NaOH buffer, pH 7.4) and Ca²⁺ levels were determined by using a Hitachi F-2000 fluorescence spectrophotometer (Gryniewicz et al., 1985).

Permeabilized cells. CHO(D_2) cells in 24-well plates were washed twice with 1 ml of Spinner MEM (with Earl's salts, without glutamine; GIBCO) with 0.2% BSA, and incubated in 1 ml of the same medium containing 50 μg/ml saponin plus final concentrations of drugs. At least 90% of the cells became permeable to trypan blue within 5 min of saponin treatment. For ³H-AA release experiments, cells were previously labeled for 2 hr as described above and unincorporated radioactivity eliminated by washing.

Statistical analysis. Results are expressed as mean ± SEM of *n* experiments. EC₅₀ and IC₅₀ values were calculated by nonlinear regression analysis, using data obtained from at least three separate experiments. Statistical comparisons between means ± SEM were carried out using analysis of variance, unless otherwise indicated.

Materials. Ca²⁺ ionophore A23187, dopamine, saponin, 4-β-phorbol-12-myristate-13-acetate (PMA), 4-α-phorbol, bovine serum albumin (essentially fatty acid-free), pertussis toxin, staurosporine, forskolin, and isobutylmethylxanthine were from Sigma (France). 1-*O*-octadecyl-2-*O*-methyl-*sn*-glycero-3-phosphorylcholine (OMPC) was obtained from Biomol Research Laboratories (Plymouth Meeting, PA); guanosine-5'-*O*-(3-thiotriphosphate) (GTP-γ-S), from Boehringer Mannheim (France); fura-2 acetoxyethyl ester, from Molecular Probes (Eugene, OR); and Rp-adenosine-cAMP monophosphothioate (Rp-cAMPS), from Research Biochemicals (Natick, MA). Quinpirole was a generous gift of Eli Lilly (Indianapolis, IN), and raclopride, of Astra (Södertälje, Sweden). Verrucosin B was purified from the marine mollusk *Doris verrucosa* as described (Cimino et al., 1988).

Results

Phorbol ester switches preferential D₂ receptor signaling from inhibition of adenylyl cyclase to facilitation of AA release

In CHO(D_2) cells, dopamine enhanced the release of ³H-AA stimulated by agents that elevate intracellular Ca²⁺, such as the Ca²⁺ ionophore A23187 (Fig. 1*a*), and inhibited the accumulation of cAMP stimulated by forskolin (Fig. 1*b*). Concentrations of dopamine that produced half-maximal potentiation (EC₅₀) of ³H-AA release and half-maximal inhibition (IC₅₀) of cAMP accumulation were 24 nM and 5 nM, respectively. When cells were stimulated with dopamine in the presence of the PKC activator PMA (100 nM), the resulting potentiation of A23187-evoked ³H-AA release was much greater than with dopamine alone (Fig. 1*a*). Incubation with PMA affected both the maximal response to dopamine (from 300% to 700% of A23187-evoked ³H-AA release) and its EC₅₀ (from 24 nM to 8 nM). In contrast with this facilitating action on AA release, treatment with PMA decreased the inhibition of adenylyl cyclase activity caused by stimulation of D_2 receptors (Fig. 1*b*). In the presence of phorbol ester, maximal dopamine-induced inhibition of cAMP accumulation was reduced from 70% to 50% of control and its IC₅₀ was increased from 5 nM to 30 nM. PMA was as potent in enhancing dopamine potentiation of ³H-AA release (EC₅₀ = 8 nM PMA) as in preventing its inhibition of adenylyl cyclase (EC₅₀ = 5 nM) (Fig. 1*c*).

In unstimulated CHO(D_2) cells, PMA had no effect on either cAMP formation or ³H-AA release (not shown). Likewise, levels of cAMP measured in the presence of PMA (1 μM) plus forskolin (10 μM) were 107 ± 11% of those found with forskolin alone (mean ± SEM, *n* = 4), indicating that the phorbol ester exerted no direct action on adenylyl cyclase activity. By contrast, PMA enhanced release of ³H-AA evoked by Ca²⁺ ionophore A23187 in the absence of D_2 receptor stimulation (Figs. 1*a*, 2*A*). The result confirms previous reports showing that phorbol esters potentiate evoked ³H-AA release from CHO cells, likely by stimulating phosphorylation of cytosolic phospholipase A₂ (cPLA₂) (Felder et al., 1991; Kanterman et al., 1991; Lin et al., 1992). However, even at maximal concentrations of drug (0.1–1 μM), the effect of PMA on A23187-evoked ³H-AA release was markedly weaker than that produced by stimulating D_2 receptors with quinpirole, a D_2 receptor agonist (Fig. 2), or with dopamine (not shown). Furthermore, when PMA and quinpirole were combined, the resulting ³H-AA release was much greater than the sum of the effects of the two drugs used separately (Fig. 2). The greater-than-additive release produced by concomitant activation of D_2 receptors and PKC suggests that the two stimuli may act through independent pathways to cause a synergistic facilitation of cPLA₂ activity.

The effects of PMA on D₂ receptor signaling are mediated by PKC

The actions of PMA on D_2 receptor signaling were produced through activation of PKC. In agreement, we observed that (1) PMA enhanced ³H-AA release and prevented adenylyl cyclase inhibition with similar potencies (Fig. 1*c*); (2) the inactive analog 4-α-phorbol-12,13-didecanoate was ineffective (Table 1); (3) the effects of PMA on both cAMP accumulation and ³H-AA release were prevented by the PKC inhibitors staurosporine (0.5 μM) and OMPC (50 μM) (Helfman et al., 1983); (4) a non-phorbol ester PKC activator, the marine diterpene verrucosin B (De Petrocellis et al., 1991), enhanced ³H-AA release to an extent





Figure 1. Effects of the PKC activator PMA on intracellular signaling in CHO cells expressing the D₂ receptor subtype. *a*, PMA potentiates D₂-dependent facilitation of ³H-AA release. CHO(D₂) cells, labeled by incubation with ³H-AA, were incubated for 30 min with the Ca²⁺ ionophore A23187 (2 μM), plus dopamine at the indicated concentrations, in the absence (open circles) or in the presence (solid circles) of PMA (0.1 μM). Results are expressed as percentage of the release evoked by stimulation with A23187 (2 μM) (297 ± 55 cpm/well, *n* = 6). *b*, PMA impairs D₂-dependent inhibition of cAMP accumulation. Cells were incubated for 20 min with the adenylyl cyclase activator forskolin (10 μM), plus dopamine, in the absence (open circles) or presence (solid circles) of PMA (0.1 μM, 10 min preincubation). Results are expressed as percentage inhibition of forskolin-stimulated cAMP level (51 ± 9 fmol/well, *n* = 6). *c*, Concentration-dependent effects of PMA on both D₂-dependent facilitation of ³H-AA release (open squares) and inhibition of cAMP accumulation (solid circles). In both cases, results are expressed as percentage of the response elicited by dopamine (50 nM).

similar to PMA (Table 1). In the absence of PMA, staurosporine did not affect D₂ receptor-dependent inhibition of adenylyl cyclase (Table 1). By contrast, the potentiation of ³H-AA release produced by D₂ receptor stimulation was reduced by PKC inhibitors (Table 1), confirming that activity of this protein kinase is an important requirement in the receptor-dependent activation of cPLA₂ (Felder et al., 1991; Kanterman et al., 1991; Lin et al., 1992).

Application of 8-bromo-cAMP, a membrane-permeant cAMP analog, enhances D₂ receptor-dependent potentiation of ³H-AA release in CHO(D₂) cells (Piomelli et al., 1991). To examine whether enhancement of ³H-AA release by PMA resulted from increased cAMP levels (subsequent to its effect on D₂ receptor-induced inhibition of adenylyl cyclase), we stimulated cells in the presence of Rp-cAMPS, a selective cAMP antagonist (Rothermel et al., 1984). Rp-cAMPS (20 μM) did not affect ³H-AA release produced by PMA plus quinpirole, a D₂ receptor agonist, suggesting that cAMP does not participate in this response (Table 1).

PMA has no effect on D₂ receptor binding

In membranes prepared from rat striatum, phosphorylation by purified PKC was shown to reduce the modulatory effect of GTP-γ-S on ³H-spiroperone binding, suggesting that PKC-mediated mechanisms may regulate striatal D₂ receptor function

(Rogue et al., 1990). Therefore, we examined the effects of PMA on D₂ receptor binding in CHO(D₂) cells, using the highly selective D₂ receptor ligand ¹²⁵I-iodosulpride (Martres et al., 1985). Treating cells with PMA (1 μM) had no effect either on the displacement of ¹²⁵I-iodosulpride binding by dopamine, measured subsequently in isolated membranes, or on the modulation of this binding by GTP-γ-S. *B*_{max} values were (in pmol/mg protein) 1.36 ± 0.12 in control and 1.37 ± 0.13 in PMA-treated cells. Likewise, EC₅₀ values for GTP-γ-S in shifting the agonist competition curve from complex to monophasic remained unchanged (14 ± 3 nM in control and 19 ± 8 nM in PMA-treated cells, *n* = 4 or 5).

PKC enables the productive coupling of D₂ receptors to cPLA₂ in CHO cells expressing the D₂₍₄₁₅₎ receptor subtype

The effects of PMA on D₂-dependent potentiation of cPLA₂ and inhibition of adenylyl cyclase activities were further examined using a clonal line of CHO cells expressing the human D₂₍₄₁₅₎ receptor, CHO(D₂₍₄₁₅₎). This isoform, derived from alternative splicing of the D₂ receptor gene transcript (Schwartz et al., 1992), also termed D_{2S}, differs from the D₂₍₄₄₄₎ (here referred to as D₂) in that it lacks a segment of 29 amino acid residues located within the third putative cytoplasmic loop of the protein, a domain that is thought to participate in G-protein coupling and in effector regulation (Dohlman et al., 1991).




Figure 2. Effects of PMA on stimulated ³H-AA release from CHO(D₂) cells. *A*, Concentration-dependent effects of PMA alone (open squares), PMA in the presence of A23187 (open circles), and PMA in the presence of A23187 plus quinpirole, a D₂ receptor agonist (200 nM) (solid circles). *B*, Effects of PMA (1 μM) on the release of ³H-AA evoked by A23187 (0.5–5 μM) or by A23187 plus quinpirole (25 nM).

Cell lines expressing the two splicing variants at similar levels (see Materials and Methods) showed no marked difference either in their endogenous cPLA₂, adenylyl cyclase, and PKC activities, or in their ability to respond to the activation of a constitutive, G_i-coupled receptor. In agreement, release of ³H-AA stimulated by A23187 and accumulation of cAMP stimulated by forskolin were similar in the two cell lines (not shown). Likewise, no significant difference was seen between these cell lines in the ability of PMA to potentiate A23187-evoked ³H-AA release: in three experiments, potentiations by PMA (50 nM) were 177 ± 22% of A23187-evoked release in CHO(D₂) cells

Table 1. The effects of PMA on D₂ receptor signaling are mediated by PKC

Treatment	³ H-AA release (% of A23187 plus quinpirole)	Inhibition of cAMP accumulation (% of quinpirole)
Quinpirole	100	100
+ stauro	83 ± 2	109 ± 9
+ PMA	252 ± 19*	71 ± 5*
+ 4-α-phorbol	105 ± 8	97 ± 1
+ PMA/stauro	61 ± 4*	92 ± 5
+ PMA/OMPC	130 ± 5	ND
+ verrucosin B	289 ± 20*	ND
+ verrucosin B/stauro	89 ± 4	ND
+ PMA/Rp-cAMPs	284 ± 22*	ND

For experiments on ³H-AA release, prelabeled CHO(D₂) cells were incubated for 30 min with A23187 (4 μM) and quinpirole (1 μM) plus various combinations of the following drugs: PMA (0.5 μM); staurosporine (stauro) (0.5 μM); OMPC (50 μM); verrucosin B (1 μM); Rp-cAMPs (20 μM). For experiments on cAMP accumulation, cells were incubated for 20 min with quinpirole (0.1 μM) plus the drugs mentioned above. Results are expressed as percentage ± SEM of the effect produced by quinpirole in 3–10 separate experiments. ND, not determined.

* Statistically different from A23187 plus quinpirole.

and 149 ± 6% in CHO(D₂₍₄₁₅₎) cells. Furthermore, the two lines did not differ in their response to the application of extracellular ATP, which elevates intracellular Ca²⁺ levels and stimulates cPLA₂ activity by interacting with constitutive type 2 purinergic receptors, coupled to a G_i-protein (Gupta et al., 1990). ATP was equally potent (EC₅₀ = 5 μM) in elevating Ca²⁺ and in evoking ³H-AA release in both CHO(D₂) and CHO(D₂₍₄₁₅₎) cells.

By contrast, stimulation of CHO(D₂₍₄₁₅₎) cells with dopamine had little effect on A23187-evoked ³H-AA release and only weakly inhibited forskolin-stimulated adenylyl cyclase activity (Fig. 3). Similar results were obtained in four of five CHO(D₂₍₄₁₅₎) cell lines (not shown), suggesting that the productive coupling of transfected D₂₍₄₁₅₎ receptors to endogenous cPLA₂ and adenylyl cyclase activities was inefficient in these cells. However, when CHO(D₂₍₄₁₅₎) cells were stimulated with a combination of dopamine plus PMA (0.5 μM), release of ³H-AA was enhanced to an extent comparable to that produced by the two agents in CHO(D₂) cells (Fig. 3a). Maximal concentrations of PMA (0.1–1 μM) increased ³H-AA release by about sevenfold over dopamine alone. As seen with CHO(D₂) cells, PMA diminished dopamine-induced inhibition of adenylyl cyclase in CHO(D₂₍₄₁₅₎) cells (Fig. 3b). The results, showing that treatment with a PKC activator fully restores D₂ receptor-dependent potentiation of ³H-AA release, suggest that stimulated PKC activity enables the productive coupling between D₂₍₄₁₅₎ receptors and cPLA₂ in CHO(D₂₍₄₁₅₎) cells. They also support the idea that, where coupling to cPLA₂ is already functional, as in CHO(D₂) cells, PKC reinforces it, resulting in a much greater receptor-dependent facilitation of Ca²⁺-evoked AA release.

PMA exerts opposing actions on the sensitivities of adenylyl cyclase and cPLA₂ to GTP-γ-S

To examine further the actions of PKC on D₂ receptor signaling, we used stably permeabilized CHO(D₂) cells, previously labeled by incubation with ³H-AA. When the cells were exposed to an extracellular medium containing saponin (50 μg/ml) and no added Ca²⁺, they released ³H-AA in the medium in a Ca²⁺-




Figure 3. Effects of PMA on cAMP accumulation and ³H-AA release from CHO cells expressing the D₂₍₄₁₅₎ receptor isoform, an alternative splicing variant of the D₂ receptor. *a*, PMA enables D₂-dependent facilitation of ³H-AA release in CHO(D₂₍₄₁₅₎) cells. Cells were labeled by incubation with ³H-AA, and incubated for 30 min with A23187 (2 μM) plus dopamine, in the absence (open circles) or in the presence (solid circles) of PMA (0.1 μM). Results are expressed as percentage of release evoked by stimulation with A23187 (255 ± 17 cpm/well, *n* = 19). *b*, Weak D₂-dependent inhibition of cAMP accumulation in CHO(D₂₍₄₁₅₎) cells, and its prevention by PMA. Cells were incubated for 20 min with the adenylyl cyclase activator forskolin (10 μM), plus dopamine, in the absence (open circles) or in the presence (solid circles) of PMA (0.1 μM, 10 min preincubation). Results are expressed as percentage inhibition of forskolin-stimulated cAMP level (99 ± 12 fmol/well, *n* = 3).

dependent manner, as shown by the ability of EGTA (5 mM) to inhibit the release (from 1669 ± 86 to 1209 ± 42 cpm/well; *p* < 0.05 by paired *t* test, *n* = 12). In additional experiments, quinpirole (5 μM) was found to produce a small but significant increase in ³H-AA release, which was markedly enhanced by coapplication of GTP-γ-S (100 μM), a nonhydrolyzable GTP analog, and inhibited by treatment with either raclopride, a D₂ receptor antagonist (5 μM) (Table 2), EGTA (5 mM), or pertussis toxin (200 ng/ml; 4 hr) (not shown). Alone, GTP-γ-S had no effect on ³H-AA release (Table 2).

The ability of GTP-γ-S to synergize with D₂ receptors in potentiating cPLA₂ activity in a pertussis toxin-sensitive manner confirms that this response is transduced via a G_i-protein (Felder et al., 1991; Piomelli et al., 1991). To determine the effects of PMA on this synergistic interaction, saponin-permeabilized CHO(D₂) cells were stimulated with GTP-γ-S in the

Table 2. Effects of quinpirole and GTP-γ-S on Ca²⁺-evoked ³H-AA release from permeabilized CHO(D₂) cells

Treatment	³ H-AA release (cpm/well)
Control	2092 ± 136
Quinpirole (5 μM)	3089 ± 386*
Quinpirole + raclopride (5 μM)	2204 ± 402
GTP-γ-S (100 μM)	2284 ± 289
GTP-γ-S + raclopride	2205 ± 480
GTP-γ-S + quinpirole	3875 ± 457*

Cells were prelabeled with ³H-AA and stably permeabilized by incubation in a medium containing saponin (50 μg/ml) and no added Ca²⁺, plus the appropriate drugs (30 min). ³H-AA release, measured in samples of the incubation media, is expressed as cpm/well ± SEM from a representative experiment carried out in quadruplicate. Similar results were obtained in six additional experiments.

* Statistically different from control.

presence of a concentration of dopamine (1 μM), which alone had little or no effect on ³H-AA release. In control incubations, GTP-γ-S synergized with 1 μM dopamine to cause ³H-AA release only at concentrations greater than 50 μM. However, the inclusion of PMA (50 nM) in the incubation medium resulted both in increased GTP-γ-S-induced ³H-AA release and in enhanced sensitivity to the nucleotide analog (Fig. 4*a*). Basal, Ca²⁺-dependent release of ³H-AA was also increased by PMA (Fig. 4*a*), whereas Ca²⁺-independent release (in the presence of 5 mM EGTA) remained unaffected (not shown).

Next, we used saponin-permeabilized CHO(D₂) cells to investigate the actions of PMA on D₂ receptor-dependent inhibition of adenylyl cyclase activity. Forskolin-stimulated release of cAMP in the medium of permeabilized cells was inhibited weakly by dopamine alone (1 μM) (44% inhibition) and by GTP-γ-S (100 μM) (49% inhibition) or, more effectively, by a combination of dopamine plus GTP-γ-S (Fig. 4*b*). Inclusion of PMA (100 nM) in the incubations prevented the ability of low concentrations of GTP-γ-S to inhibit adenylyl cyclase activity (Fig. 4*b*). In the presence of phorbol ester, the concentration of extracellular GTP-γ-S necessary to produce maximal inhibition of cAMP accumulation was increased from 10 μM to 100 μM (Fig. 4*b*).

Together, the results obtained with permeabilized CHO(D₂) cells suggest that activation of PKC may exert opposing actions on the coupling efficacies of D₂ receptors with cPLA₂ and adenylyl cyclase by modifying the sensitivities of these signaling pathways to intracellular GTP. These changes in coupling strength may, in turn, underlie the ability of PKC, observed in intact CHO(D₂) cells, to direct the preferential signaling at D₂ receptors from inhibition of adenylyl cyclase to facilitation of ³H-AA release.

Discussion

Our experiments suggest that PKC activity may control effector coupling at the D₂ receptor by switching it from inhibition of adenylyl cyclase, prevalent in unstimulated cells, to facilitation of AA release, prevalent in cells where PKC has been activated. By remodeling the pattern of intracellular second messengers generated at the D₂ receptor, such an activity-dependent switch may modify the physiological actions of dopamine in select cells, and potentially represent, if operative in neurons, a novel form of short-term synaptic plasticity. Here we will discuss, first, the possible mechanisms underlying the regulation by PKC of D₂

receptor signaling and, next, its possible relevance and implications for the CNS.

Regulation of cPLA₂ activity by G_i-coupled receptors and by protein phosphorylation

We have shown here that (1) in intact CHO(D₂) cells, stimulation of PKC enhances the D₂ receptor-dependent facilitation of AA release; (2) in permeabilized cells, this enhancement is accompanied by a reduction in the concentration threshold of the GTP analog GTP-γ-S necessary to potentiate AA release; and (3) stimulation of PKC unmasks the coupling of D₂₍₄₁₅₎ receptors to cPLA₂ in a cell line in which this coupling is normally impaired. This impairment was likely caused by inadequate recognition between transfected D₂₍₄₁₅₎ receptor and endogenous G-protein, and does not appear to be consistently linked to the expression of this shorter receptor isoform. In agreement, one of five cell lines tested showed a normal AA response (not shown).

Recent studies on CHO cells have greatly advanced our understanding of the molecular mechanisms underlying receptor-operated release of AA from membrane phospholipids. It is now recognized that some membrane receptors (e.g., P₂-purinergic), acting through a transducing G_i-protein, can induce the Ca²⁺-dependent translocation of an AA-specific cPLA₂ from the cytosol to the cell membrane, and its subsequent activation (Gupta et al., 1990; Lin et al., 1992). These events are accompanied by increased phosphorylation of cPLA₂ on a serine residue, and can be prevented by treatment with PKC inhibitors, suggesting that the protein kinase plays an important role in enabling activation of cPLA₂ (Kanterman et al., 1991; Lin et al., 1992).

Furthermore, heterologous expression of D₂ receptors in CHO cells has led to the demonstration of an additional mechanism involved in regulating cPLA₂ activity. In transfected CHO cells, D₂ receptor stimulation potently facilitates release of AA evoked either by stimulating constitutive P₂ receptors or by applying Ca²⁺ ionophores, but has no effect on basal release (Kanterman et al., 1991; Piomelli et al., 1991). This permissive action of D₂ receptors on cPLA₂ activity, which is also seen with other transfected G_i-coupled receptors, such as α₂-adrenergic and m₂-muscarinic (Felder et al., 1991), is transduced by a G_i-protein. In agreement, the response (1) is prevented by treatment with pertussis toxin (Felder et al., 1991; Piomelli et al., 1991) and (2) is potentiated in permeabilized cells by coapplication of GTP-γ-S (present results). Although the identity of the transducing G-protein is not known, this is unlikely to be a G_o, because these proteins are not expressed in CHO cells (Gupta et al., 1990). In addition, differences in the sensitivity to pertussis toxin suggest that the transducing G_i may be distinct from the G_i involved in coupling the D₂ receptor to inhibition of adenylyl cyclase activity (Felder et al., 1991).

In summary, three factors appear to contribute to receptor-stimulated AA release in CHO cells: first, the state of phosphorylation of cPLA₂, which may play an enabling role in the enzyme translocation and activation processes; second, the levels of intracellular Ca²⁺, which may signal the turning on and off of these processes; and third, the presence of an activated G_i-protein, which may exert a permissive action on cPLA₂ after translocation of the phospholipase to the membrane, allowing expression of its full catalytic activity. Our findings suggest that PKC may act to reinforce the productive G_i-mediated coupling between D₂ receptors and cPLA₂. Potential sites of action for




Figure 4. Effects of PMA on the sensitivity of D₂ receptor-dependent responses to GTP-γ-S in permeabilized CHO(D₂) cells. *a*, PMA enhances the sensitivity to GTP-γ-S of D₂-dependent facilitation of ³H-AA release. Cells were labeled by incubation with ³H-AA, and incubated (30 min) in a permeabilizing medium containing saponin (50 μg/ml), plus final concentrations of dopamine (1 μM) and GTP-γ-S (0–100 μM) in the absence (open circles) or in the presence (solid circles) of PMA (50 nM). Results are from one experiment performed in triplicate (representative of six) and are expressed as percentage of maximal ³H-AA release after subtraction of Ca²⁺-independent release (determined in the presence of 5 mM EGTA). In the experiment shown, maximal ³H-AA release (obtained in the presence of dopamine, PMA, plus 50 μM GTP-γ-S) was, before background subtraction, 1390 ± 35 cpm/well. Background release (measured in the presence of 5 mM EGTA) was 1107 ± 51 cpm/well. *b*, PMA decreases the sensitivity to GTP-γ-S of D₂-mediated inhibition of cAMP accumulation. Cells were incubated for 20 min in a permeabilizing medium containing saponin plus final concentrations of dopamine (1 μM) and GTP-γ-S (0–100 μM), in the absence (open circles) or in the presence (solid circles) of PMA (100 nM). IBMX (0.1 mM) and forskolin (10 μM) were added to all incubations. Results are from one experiment performed in triplicate (representative of three) and are expressed as percentage of maximal inhibition of cAMP formation. Maximal inhibition (obtained in the presence of dopamine plus 100 μM GTP-γ-S) was 60% of forskolin alone, which produced a four- to sixfold increase in cAMP levels. In control cells, measured cAMP levels were 2.4–6.3 fmol/well, and in the presence of forskolin, 9–31.7 fmol/well.

the protein kinase include the D₂ receptor itself, which contains a consensus sequence for PKC phosphorylation (Bunzow et al., 1988); the transducing G-protein, which may be a PKC substrate, as shown for G₁₂ and G_z (Bushfield et al., 1990; Lounsbury et al., 1991); or the G-protein-regulating protein phosducin (Bauer et al., 1992).

PKC regulates D₂ receptor-dependent inhibition of cAMP accumulation

We find that stimulation of PKC reduces the inhibitory action of D₂ receptor agonists on adenylyl cyclase activity, and that this effect is accompanied by a decreased ability of GTP- γ -S to inhibit the cyclase in permeabilized cells. The findings suggest that PKC decreases the GTP sensitivity of the G_i subunit linked to adenylyl cyclase, producing, in turn, a weakened coupling of D₂ receptors with this effector system. In agreement with this possibility, phorbol esters were previously found to reduce receptor-operated inhibition of adenylyl cyclase in platelets and hepatocytes, to stimulate the phosphorylation of α G₁₂, and to reduce its sensitivity to GTP analogs (Jakobs et al., 1985; Katada et al., 1985; Bushfield et al., 1990). A similar mechanism may be operative in CHO cells. If this is the case, it will be interesting to examine whether PKC-mediated phosphorylation of α G₁₂ may also play a role in the enhancement of cPLA₂ activity produced by D₂ receptor stimulation. Following phosphorylation of α G₁₂, a greater number of D₂ receptors, which are thought to be preferentially coupled to this subunit (Senogles et al., 1990), might become free to interact with a distinct, cPLA₂-linked G-protein, whose existence was suggested by the work of Felder et al. (1991). Alternatively, phosphorylated α G₁₂ may interact better with cPLA₂ than with adenylyl cyclase. In either case, a single molecular event might affect, concomitantly and in opposite directions, the D₂-dependent regulation of both adenylyl cyclase and cPLA₂ activities. This intriguing possibility may find indirect support in two observations: (1) the identical potencies of PMA in enhancing cPLA₂ potentiation and in decreasing adenylyl cyclase inhibition and (2) the quantitatively similar, but opposing shifts produced by PMA on the sensitivities of D₂-dependent responses to GTP- γ -S.

Such a mechanism may also provide a possible explanation for the inability of PMA to affect the GTP- γ -S modulation of dopamine binding. Conceivably, no effect of PMA on this binding parameter should be seen if the linkage of D₂ receptors with G₁₂ is not affected, or if this linkage is switched onto another G-protein. A corollary to this mechanism is that PMA-induced modifications in adenylyl cyclase and cPLA₂ responses should be correlated: any given change produced on the modulation of cPLA₂ activity should result from a corresponding change in adenylyl cyclase. However, this prediction is not supported by the results obtained with CHO(D₂₍₄₁₅₎) cells, which show that activation of PKC may enhance D₂-dependent facilitation of cPLA₂ markedly, even when the adenylyl cyclase response is only slightly affected (Fig. 3).

Alternative hypothetical models may implicate multiple sites of action for PKC, for example, the concomitant phosphorylation of α G₁₂, leading to decreased cyclase inhibition, and of phosducin, leading to desinhibition of other G-proteins (Bauer et al., 1992). An essential step to allow the testing of these different possibilities will be to determine the nature of the G-protein involved in mediating the actions of D₂ receptors on cPLA₂ activity.

Possible physiological function

Although the relevance of our findings to the CNS is not yet known, several lines of evidence suggest that coupling mechanisms similar to those observed in transfected CHO cells may be found in cells constitutively expressing D₂ receptors, including neurons. G_i-mediated cyclase inhibition is a well-established response produced by D₂ receptor occupation in various brain areas (Schwartz et al., 1992). Likewise, stimulation of D₂ receptors was shown to enhance release and metabolism of AA in neural (Hillier et al., 1976) and kidney cells (Huo et al., 1991), while activation of this signaling pathway may participate in certain neuronal D₂ responses, such as stimulation of somatostatin release from hypothalamic nerve terminals (Junier et al., 1990).

AA and its metabolites are thought to serve important functions in the nervous system, where they may act both as intracellular second messengers and as local mediators, regulating K⁺ channel opening (Piomelli et al., 1987a,b; Belardetti et al., 1989; Schweitzer et al., 1990), protein kinase activity (McPhail et al., 1984; Piomelli et al., 1989), neurotransmitter release (Linch and Voss, 1990; Freeman et al., 1991), and synaptic plasticity (for review, see Fazeli, 1992). The ability of PKC to amplify D₂ receptor signaling via AA release at the expenses of cAMP inhibition may result in the recruitment of a distinct spectrum of biologically active mediators in cells in which PKC activity has been elevated. Through such a mechanism, the physiological responses to dopamine could be regulated dynamically within a neuron by stimuli promoting PKC activation, which include membrane depolarization and Ca²⁺-mobilizing neurotransmitters (e.g., glutamate acting at metabotropic receptors) (Schoepp et al., 1990).

References

- Bauer PH, Müller S, Puzicha M, Pippig S, Obermaier B, Helmreich EJM, Lohse M (1992) Phosducin is a protein kinase A-regulated G-protein regulator. *Nature* 358:73–76.
- Belardetti F, Campbell WB, Falck JR, Demontis G, Rosolowski M (1989) Products of heme-catalyzed transformation of the arachidonate derivative 12-HPETE open S-type K⁺ channels in *Aplysia*. *Neuron* 3:497–505.
- Bunzow JR, Van Tol HHM, Grandy DK, Albert P, Salon J, Christie M, Machida CA, Neve KA, Civelli O (1988) Cloning and expression of a rat D₂ dopamine receptor cDNA. *Nature* 336:783–787.
- Bushfield M, Murphy GJ, Lavan BE, Parker PJ, Hruby VJ, Milligan G, Houslay MD (1990) Hormonal regulation of G₁₂ α -subunit in intact hepatocytes. *Biochem J* 268:449–457.
- Cimino G, Gavagnin M, Sodano G, Puliti R, Mattia CA, Mazzarella L (1988) Verrucosin-A and -B, ichthyotoxic diterpenoid acid glycerides with a new carbon skeleton from dorid nudibranch *Doris verrucosa*. *Tetrahedron* 44:2301–2310.
- De Petrocellis L, Di Marzo V, Arca B, Gavagnin M, Minei R, Cimino G (1991) The effect of diterpenoid diacylglycerols on tentacle regeneration in *Hydra vulgaris*. *Comp Biochem Physiol [C]* 100:603–607.
- Di Marzo V, Piomelli D (1992) Participation of prostaglandin E₂ in dopamine D₂ receptor-dependent potentiation of arachidonic acid release. *J Neurochem* 59:379–382.
- Dohlman HG, Thorner J, Caron MG, Lefkowitz RJ (1991) Model systems for the study of seven-transmembrane segment receptors. *Annu Rev Biochem* 60:653–688.
- Fazeli MS (1992) Synaptic plasticity: on the trail of the retrograde messenger. *Trends Neurosci* 15:115–117.
- Felder CC, Williams HL, Axelrod J (1991) A transduction pathway associated with receptors coupled to the inhibitory guanine nucleotide binding protein G_i, that amplifies ATP-mediated arachidonic acid release. *Proc Natl Acad Sci USA* 88:6477–6480.
- Freeman EJ, Damron DS, Terrian DM, Dorman RV (1991) 12-Lipox-

- xygenase products attenuate the glutamate release and Ca^{2+} accumulation evoked by depolarization of hippocampal mossy fiber nerve endings. *J Neurochem* 56:1079–1082.
- Giros B, Sokoloff P, Martres MP, Riou JF, Emorine LJ, Schwartz JC (1989) Alternative splicing directs the expression of two dopamine D_2 receptors isoforms. *Nature* 342:923–926.
- Grynkiwicz G, Poenie M, Tsien RY (1985) A new generation of Ca^{2+} indicators with greatly improved fluorescence properties. *J Biol Chem* 260:3440–3450.
- Gupta SK, Diez E, Heasley LE, Osawa S, Johnson G (1990) A G protein mutant that inhibits thrombin and purinergic receptor activation of phospholipase A_2 . *Science* 249:662–666.
- Helfman DM, Barnes KC, Kinkade JM, Vogler WR, Shoji M, Kuo JF (1983) Phospholipid-sensitive Ca^{2+} -dependent protein phosphorylation system in various types of leukemia cells from human patients and in human leukemia cell lines HL60 and K5623, and its inhibition by alkyl-lysophospholipid. *Cancer Res* 43:2955–2961.
- Hillier K, Roberts PJ, Woollard PM (1976) Catecholamine-stimulated prostaglandin biosynthesis in rat brain synaptosomes. *Br J Pharmacol* 58:426P–427P.
- Huo T, Ye MQ, Healy DP (1991) Characterization of a dopamine receptor (DA_{2k}) in the kidney inner medulla. *Proc Natl Acad Sci USA* 88:3170–3174.
- Jakobs KH, Bauer S, Watanabe Y (1985) Modulation of adenylate cyclase of human platelets by phorbol ester. *Eur J Biochem* 151:425–430.
- Junier MP, Dray F, Blair I, Capdevila J, Dishman E, Falck JR, Ojeda S (1990) Epoxygenase products of arachidonic acid are endogenous constituents of the hypothalamus involved in D_2 receptor-mediated, dopamine-induced release of somatostatin. *Endocrinology* 126:1534–1540.
- Kanterman RY, Mahan LC, Briley EM, Monsma FJ, Sibley DR, Axelrod J, Felder CC (1991) Transfected D_2 receptors mediate the potentiation of arachidonic acid release in Chinese hamster ovary cells. *Mol Pharmacol* 39:364–369.
- Katada T, Gilman AG, Watanabe Y, Bauer S, Jakobs KH (1985) Protein kinase C phosphorylates the inhibitory guanine-nucleotide-binding regulatory component and apparently suppresses its function in hormonal inhibition of adenylate cyclase. *Eur J Biochem* 151:431–437.
- Lin L-L, Lin A-Y, Knopf JL (1992) Cytosolic phospholipase A_2 is coupled to hormonally regulated release of arachidonic acid. *Proc Natl Acad Sci USA* 89:6147–6151.
- Linch MA, Voss KL (1990) Arachidonic acid increases inositol phospholipid metabolism and glutamate release in synaptosomes prepared from hippocampal tissue. *J Neurochem* 55:215–221.
- Lledo PM, Homburger V, Bockaert J, Vincent JD (1992) Differential G protein-mediated coupling of D_2 dopamine receptors to K^+ and Ca^{2+} currents in rat anterior pituitary cells. *Neuron* 8:455–463.
- Lounsbury KM, Casey PJ, Brass LF, Manning DR (1991) Phosphorylation of G_i in human platelets. Selectivity and site of modification. *J Biol Chem* 266:22051–22056.
- Martres MP, Sokoloff P, Delandre M, Schwartz JC, Protais P, Costentin J (1984) Selection of dopamine antagonists discriminating various behavioral responses and radioligand binding sites. *Naunyn-Schmiedeberg Arch Pharmacol* 325:102–115.
- Martres MP, Bouthenet ML, Salès N, Sokoloff P, Schwartz JC (1985) Widespread distribution of brain dopamine receptors evidenced with (^{125}I)iodosulpride, a highly selective ligand. *Science* 228:752–755.
- McPhail LC, Clayton CC, Snyderman R (1984) A potential second messenger role for unsaturated fatty acids: activation of Ca^{2+} -dependent protein kinase. *Science* 224:622–625.
- Nishizuka Y (1988) The molecular heterogeneity of protein kinase C and its implication for cellular regulation. *Nature* 334:661–665.
- Piomelli D, Volterra A, Dale N, Siegelbaum SA, Kandel ER, Schwartz JH, Belardetti F (1987a) Lipoxygenase metabolites of arachidonic acid as second messengers for presynaptic inhibition in *Aplysia* sensory cells. *Nature* 328:38–43.
- Piomelli D, Shapiro E, Feinmark SJ, Schwartz JH (1987b) Metabolites of arachidonic acid in the nervous system of *Aplysia*: possible mediators of synaptic modulation. *J Neurosci* 7:3675–3686.
- Piomelli D, Wang JKT, Sihra TS, Nairn AC, Czernik AJ, Greengard P (1989) Inhibition of Ca^{2+} /calmodulin-dependent protein kinase II by arachidonic acid and its metabolites. *Proc Natl Acad Sci USA* 86:8550–8554.
- Piomelli D, Pilon C, Giros B, Sokoloff P, Martres M-P, Schwartz J-C (1991) Dopamine activation of the arachidonic acid cascade as a basis for D_1/D_2 receptor synergism. *Nature* 353:164–167.
- Rogue P, Zwiller J, Malviya AN, Vincendon G (1990) Phosphorylation by protein kinase C modulates agonist binding to striatal dopamine D_2 receptors. *Biochem Int* 22:575–582.
- Rothermel JD, Jastorff B, Parker Botelho LH (1984) Inhibition of glucagon-induced glycogenolysis in isolated rat hepatocytes by the R_p diastereoisomer of adenosine cyclic 3'5'-phosphorothioate. *J Biol Chem* 259:8151–8155.
- Schoepp D, Bockaert J, Sladeczek F (1990) Pharmacological and functional characteristics of metabotropic excitatory amino acid receptors. *Trends Pharmacol Sci* 11:508–515.
- Schwartz JC, Giros B, Martres MP, Sokoloff P (1992) The dopamine receptor family: molecular biology and pharmacology. *Semin Neurosci* 4:99–108.
- Schweitzer P, Madamba S, Siggins GR (1990) Arachidonic acid metabolites as mediators of somatostatin-induced increase of neuronal M-current. *Nature* 346:464–467.
- Senogles SE, Spiegel AM, Padrell E, Iyengar R, Caron MG (1990) Specificity of receptor-G protein interactions. Discrimination of G_i subtypes by the D_2 dopamine receptor in a reconstituted system. *J Biol Chem* 265:4507–4514.
- Sibley DR, Monsma FJ (1992) Molecular biology of dopamine receptors. *Trends Pharmacol Sci* 13:61–69.
- Spiegel AM (1992) G proteins in cellular control. *Curr Opin Cell Biol* 4:203–211.