

# Coordinated Gating of TRP-Dependent Channels in Rhabdomeral Membranes from *Drosophila* Retinas

Joan E. Haab,<sup>1</sup> Cecilia Vergara,<sup>2</sup> Juan Bacigalupo,<sup>2</sup> and Peter M. O'Day<sup>1</sup>

<sup>1</sup>Institute of Neuroscience and Department of Biology, University of Oregon, Eugene, Oregon 97403, and <sup>2</sup>Department of Biology, Facultad de Ciencias, University of Chile, Santiago, Chile

Using a newly developed dissociation procedure, we isolated the specialized rhabdomeral membranes from *Drosophila* retinal photoreceptors. From these membranes, we have recorded spontaneous active currents in excised patch, voltage-clamp recordings. We observed rapid opening events that closely resembled those ascribed to one class of light-activated channels, TRP. All activity exhibited Ba<sup>2+</sup> permeability, little voltage dependence, and sensitivity to La<sup>3+</sup> block. Mutational analysis indicated that the spontaneous activity present in these membranes was TRP-dependent. Excised patches from wild-type rhab-

domeral membranes exhibited a wide range of conductance amplitudes. In addition, large conductance events exhibited many conductance levels in the open state. Block of activity by La<sup>3+</sup> both developed and recovered in a stepwise manner. Our results indicate that TRP-dependent channels have a small unitary conductance and that many channels can be gated coordinately.

**Key words:** *Drosophila*; TRP; phototransduction; channel; barium; rhabdomere

Phototransduction in *Drosophila* retinal photoreceptor cells involves G-protein-mediated biochemical cascades initiated by the activation of the receptor rhodopsin by light and culminating in an electrical response generated by the opening of many TRP and TRP-related plasma membrane ion channels (for review, see Montell, 1999). Many channels open in response to the activation of a single receptor protein, indicating that amplification of the signal occurs at some step within the biochemical cascades. Amplification and its modulation are essential features of signaling systems, but their underlying mechanisms are unknown in *Drosophila* phototransduction.

Recent studies indicate that there is little or no amplification within the first three enzymatic steps of the phototransduction pathway (Scott and Zuker, 1998). One light-activated rhodopsin molecule activates a single heterotrimeric G-protein (DG<sub>q</sub>), which in turn activates a single phospholipase C molecule (PLC). Activation of PLC is thought to lead to the gating of plasma membrane channels. Three channel polypeptides have been identified and implicated in *Drosophila* phototransduction, TRP, TRPL, and TRP- $\gamma$  (Phillips et al., 1992; Niemeyer et al., 1996; Reuss et al., 1997; Xu et al., 1997, 2000). However, the assembly of these subunits *in vivo* and the gating mechanisms of the channels are not well understood.

TRP or TRP-related channel proteins are found in a wide variety of organisms and tissue types (for review, see Putney, 1999; Harteneck et al., 2000). They are generally thought to be store-operated channels, important in Ca<sup>2+</sup> regulation and gated through a poorly understood mechanism linked to the depletion of internal Ca<sup>2+</sup> stores. Evidence from the heterologous expression of both the *Drosophila* TRP and TRPL proteins has suggested that the gating of these channels is likewise linked to internal store

depletion (Vaca et al., 1994; Dong et al., 1995; Petersen et al., 1995; Yagodin et al., 1998). Recent evidence, however, suggests that they may be gated directly by PLC-generated diacylglycerol (DAG) or by a metabolic byproduct of DAG (Chyb et al., 1999).

Ideally, one would like to study these channels in cell-attached patches, in intact phototransducing cells, but the complicated cellular morphology has precluded this type of recording. To approach these channels directly, we have developed a novel preparation with which to isolate the rhabdomeral membranes, in which TRP- and TRPL-dependent channels are localized. Here we report the first patch-clamp recordings from TRP-dependent channels in their native membranes. We find that groups of channels open and close in a concerted manner. This coordinated gating may contribute to the generation of amplification within the phototransduction cascade.

## MATERIALS AND METHODS

**Solutions.** Bath solution contained (in mM): 120 NMDG, 15 tetraethylammonium chloride, 10 *N*-tris(hydroxymethyl)methyl-2-aminoethanesulfonic acid buffer, 25 L-proline, and 10 barium chloride (Sigma, St. Louis, MO), pH 7.2. Pipette solution was identical to bath solution, except that the concentration of BaCl<sub>2</sub> was only 1 mM.

**Membrane preparation.** Corneal tissue from the eyes of five adult flies (red- or white-eyed Oregon R, mutants *trp*<sup>p343</sup>, *trpl*<sup>p302</sup>, *trp*<sup>p343</sup>, and *trpl*<sup>p302</sup>, courtesy of Dr. Charles Zuker, University of California, San Diego CA; and *trp*<sup>CM</sup>, courtesy of Dr. William Pak, Purdue University, Lafayette, IN) was dissected into an ~50  $\mu$ l bubble of bath solution on a dissecting dish. This solution was transferred by pipette into a small-volume Eppendorf tube containing ~0.5 ml of bath solution and a small amount of 0.1-mm-diameter zirconia-silicone beads (BioSpec Products, Bartlesville, OK). The tube was shaken for 20 sec at 2500 rpm using a Mini-Bead Beater (BioSpec Products) such that the beads sheared the plasma membrane and cytosol from the rhabdomeral membranes. The solution was transferred to the recording chamber by pipette, leaving the beads settled at the bottom of the tube. The cell fragments were allowed to settle to the bottom of the dish mounted to the microscope stage for several minutes before we began recording.

**Recording.** Recording pipettes were pulled from borosilicate glass using a laser puller (Sutter Instruments, Novato, CA). All recording electrodes had a tip resistance of 25–40 M $\Omega$ , measured in bath solution. Viewing and recording conditions were the same as those described previously (Bacigalupo et al., 1995). Suitable rhabdomeral membranes exhibited a smooth surface and were free of attached plasma membrane fragments. Membranes were selected by eye and sampled using a patch recording pipette. Patches were excised by pulling the pipette away from the rhabdomeral fragment, which was stuck to the bottom of the recording chamber. All data were acquired in excised patch voltage-clamp recordings using Axopatch 200A patch clamp (Axon Instruments, Foster City, CA). Currents were filtered at 2 kHz (eight-pole Bessel filter), acquired digitally (Digi-

Received May 31, 2000; revised July 14, 2000; accepted July 17, 2000.

This work was supported by the American Heart Association, Oregon Affiliate, and by National Science Foundation Grant INT-9604977 in collaboration with Fundacion Andes/Conicyt Grants Fundecyt-1990938 and 1981053 and Grant P99-031 Program ICM Mideplan. J.E.H. was supported by National Institutes of Health predoctoral training Grant 5-T32-GM07257. We gratefully acknowledge Drs. Judith Eisen, William Roberts, Roger Hardie, Craig Montell, and Enrico Nasi for critical reading of earlier versions of this manuscript, Drs. William Pak and Charles Zuker for mutant fly strains, Drs. Charles Zuker and Craig Montell for anti-TRP antibodies, and Dr. Nancy Olea for electron microscopy sample preparation.

Correspondence should be addressed to Dr. Peter O'Day, Institute of Neuroscience, University of Oregon, Eugene, OR 97403. E-mail: oday@uoneuro.uoregon.edu.

Copyright © 2000 Society for Neuroscience 0270-6474/00/207193-06\$15.00/0

Data 1200 interface; Axon Instruments), and stored on computer disk. Applications of  $\text{La}^{3+}$  were made using a four-bore glass panpipe spritzer (50–80  $\mu\text{m}$ ) and a microperfusion system (Goodman and Art, 1996).

**Analysis.** Offline analysis was performed using pClamp 6.0 (Axon Instruments). Gaussian fits to all-points histograms were performed using Pstat (Axon Instruments) and either the Simplex least squares or the Marquardt method. Best fits were chosen by comparing the residuals for each. Calculations of open probability ( $P_{\text{open}}$ ) were made with the assumption that opening events could be analyzed as unitary events.  $P_{\text{open}}$  values were calculated using Fetchan and Pstat (Axon Instruments). Records were initially examined in Fetchan. Opening and closing events were counted using the half-maximal conductance of the smallest discernable event as the cutoff criteria. For records in which the events were very small, we altered the cutoff criteria to three-fourths maximal conductance of the smallest events, to reduce the possibility of counting closed state noise as open events. The results of these counts were analyzed in pStat to determine the mean open probability for each entire trace.

**Electron microscopy.** For transmission electron microscopic images, two preparations of 100 eyes each were dissected from adult Oregon red-eyed flies into bath solution. One preparation was left intact, whereas the second was subjected to dissociation by shaking. Dissociated cells were centrifuged at  $13,000 \times g$  for 20 min at  $4^\circ\text{C}$  and fixed in 3% glutaraldehyde in 0.1 M phosphate buffer, pH 7.0, for 30 min. Samples were suspended in 2% agar in an Eppendorf tube, and post-fixation was done in 1% osmium tetroxide for 1 hr. Inclusion into Epon 812 was done by following a previously described technique (Suzuki et al., 1993), and the samples were observed in a Zeiss (Thornwood, NY) EM 109 transmission electron microscope at the University of Chile School of Medicine (Santiago, Chile).

**Western blot analysis.** Samples for Western blots were prepared by homogenizing the tissue collected from four dissected eyes and denaturing proteins by boiling for 1 min. Samples were run on SDS-polyacrylamide gels, transferred to nitrocellulose, and blotted according to standard techniques (Sambrook et al., 1989), using anti-TRP antibodies generously provided by Drs. Craig Montell (Johns Hopkins University, Baltimore, MD) and Charles Zuker (University of California, San Diego, CA).


## RESULTS

In *Drosophila* retinal photoreceptors, the light-activated channels reside in a specialized region of tightly packed microvilli known as the rhabdomere, which runs the length of each photoreceptor cell. All of the known enzymes involved in phototransduction are localized to this region. In the *Drosophila* retina, photoreceptor cells are bound together in an eight-cell bundle, called an ommatidium, with the rhabdomere of each cell oriented toward the center of the bundle (for review, see Ready, 1989; Fig. 1*A,C*). Thus, in an intact ommatidium normally used for whole-cell patch-clamp recording, the rhabdomeres are inaccessible to the recording pipette.

To study the light-activated channels directly, we developed an isolation procedure that exposes the rhabdomeral membranes in which they reside. Using a mechanical dissociation technique, we stripped away the exposed plasma membrane from the ommatidia, leaving the rhabdomeral membranes exposed (Fig. 1*B,D*). We examined membrane fragments using light microscopy and selected rhabdomeres with smooth membrane surfaces without apparent plasma membrane for patch recording.

We used a simplified recording solution to maximize the likelihood of observing light-activated channels. This solution substituted  $\text{Ba}^{2+}$  for  $\text{Ca}^{2+}$  as the only permeable cation.  $\text{Ca}^{2+}$  has been shown to have modulatory effects on transduction (Hardie and Minke, 1994b, 1995; Hardie, 1995; Scott et al., 1997) and commonly affects the open probabilities of some cation-permeable channels (Yue et al., 1990; Marunaka et al., 1992). Barium, in contrast, permeates light-dependent channels (Reuss et al., 1997), but it does not seem to mediate secondary modulatory effects. Monovalent cations were replaced with nonpermeant NMDG (Hardie and Minke, 1992).

More than 95% of the patches from wild-type membranes exhibited spontaneous channel activity within 1–2 min of excision, if the patch was held at voltages between  $-70$  and  $-100$  or  $+70$  and  $+100$  mV. The majority of the excised patches from wild-type membranes exhibited large-conductance, burst-like behavior, often quickly followed by a dramatic decrease in seal resistance. We have not included data from these patches in the analysis presented



**Figure 1.** *A, B*, Light micrographs with Nomarski differential interference optics show an intact ommatidium (*A*) and an isolated rhabdomeral membrane (*B*) under standard recording conditions. A recording pipette is visible at the bottom of *A*, attached to the plasma membrane of a single cell within the intact ommatidium. In the isolated rhabdomeral membranes shown in *B*, the plasma membranes normally present in intact cells have been stripped away. *C, D*, Transmitting electron micrographs show cross sections of an intact ommatidium (*C*) and an isolated rhabdomeral membrane (*D*). The electron-dense regions in the center of the intact ommatidium are the rhabdomeral membranes.


here. Some patches ( $n = 12$ ), however, made a transition from this erratic kinetic behavior to more stereotypical single-channel behavior and maintained a high seal resistance (8–20 G $\Omega$ ). Data from these patch recordings will be discussed in this paper.

We recorded a wide distribution of spontaneously active conductances in excised patches from wild-type rhabdomeral membranes (Fig. 2*A*). All of the conductances that we recorded, regardless of size, were permeable to barium, exhibited little or no voltage dependence, and were sensitive to block by  $\text{La}^{3+}$ , reminiscent of TRP-dependent light-sensitive channels observed in whole-cell recordings (Hardie and Mojet, 1995).

We observed no sustained spontaneous activity in patches excised from the double null mutant  $trp^{p343}; trpl^{p302}$  ( $n = 14$ ) or the single null mutant  $trp^{p343}$  ( $n = 6$ ). Neither did we observe the high-frequency burst-like behavior observed initially in wild-type patches. In some patches ( $n = 5$  of 14,  $trp^{p343}; trpl^{p302}$ ) we did see channel-like events. However, in every case, the duration of this activity was extremely brief, generally a single burst lasting between 5 and 15 sec within 10–15 min of recording time and containing only a few events. The patches otherwise remained silent. The few channel-like events we observed in this mutant always occurred at a single holding potential and totaled less than a dozen events, making it difficult to determine single-channel characteristics. This contrasts sharply with wild-type patches in which we observed prolonged (10–40 min) spontaneous activity in  $\sim 95\%$  of patches.

We also examined patches from the hypomorphic allele  $trp^{CM}$ , in which a small amount of TRP protein is present (as identified by Western blot analysis; data not shown). In patches from this mutant we did observe channel activity but found a distribution of much smaller conductance values (Fig. 2*B*).

For comparison, we normalized all of the data from all holding voltages by converting current to conductance and generated all-points conductance histograms for both wild-type and  $trp^{CM}$  data.




**Figure 2.** *A*, Ten individual events selected from 10 different excised patches from wild-type isolated rhabdomeral membranes, plotted in descending order of conductance size. Open state level is upward in each case.  $Ba^{2+}$  in the bath was 10 mM;  $Ba^{2+}$  in the pipette was 1 mM. Conductance values were calculated as  $g = [I/(V_{\text{hold}} - E_{\text{rev}})]$ , where  $g$  is conductance in picosiemens,  $I$  is measured current amplitude, and  $E_{\text{rev}}$  is the reversal potential for each patch, as determined for each individual patch by the slope of the current plotted for multiple holding voltages. All conductances were permeable to barium as determined by the reversal potential of the activity ( $E_{\text{rev}} = 24 \pm 6$  mV;  $n = 12$ ). The calculated equilibrium potential for  $Ba^{2+}$ , the only permeant ion in our solutions, was +29 mV. *B*, Eight individual conductance events selected from eight different excised patches taken from *trp<sup>CM</sup>* isolated rhabdomeral membranes, plotted in descending order of conductance size. *C*, *D*, All-points conductance histograms show marked differences in the distribution of conductances observed among all wild-type (*C*) and *trp<sup>CM</sup>* (*D*) excised patches. Conductance values were determined separately for each patch by dividing the measured current by the driving force. All data from each patch were binned cumulatively, and the resultant histograms were normalized to the zero peak (the largest peak in each histogram) to highlight the relative contribution of each subsequent open state. Wild type,  $n = 12$ ; *trp<sup>CM</sup>*,  $n = 8$ .

(Fig. 2*C,D*). This allowed us to examine the relative frequency of occurrence of each conductance size. The *trp<sup>CM</sup>* histogram shows primarily small conductance values compared with those of wild type, presumably reflecting the residual TRP channels expressed by this allele.

The lack of channel activity in the *trp* null mutant and the smaller conductance events observed in the *trp* hypomorph compared with those observed in wild-type patches suggested that the activity we observed in our patches is TRP-dependent. The prevalence of much smaller conductance values observed in the *trp<sup>CM</sup>* allele further suggests that the unitary conductance of individual TRP channels is small.


A key feature of the activity observed in wild-type patches is the frequent appearance of multiple conductance levels while the channel is open (Fig. 3). If, as our data suggest, all of the activity we observed is TRP-dependent and the unitary conductance of these channels is small, the large conductance events in wild-type patches must be composed of small unitary conductance, TRP-dependent channels.



**Figure 3.** Multiple conductance levels are clearly discernable during large conductance events. *Top*, *A*, ~36 sec of continuously sampled data trace.  $V_{\text{hold}} = -70$  mV. *Top*, *B–F*, Examples illustrating multiple conductance levels that occurred within the longer trace. Traces have been enlarged from the indicated regions of *A*. *Bottom*, *A–F*, Histograms corresponding to the data traces in *top*, *A–F*. Histogram *A* was generated from a larger sample of data than that represented in data panel *A* (*top*). Histograms *B–F* were generated using only the same-letter data samples. When the data samples include only a single large opening event, the all-points current histograms show distinct variability in amplitudes of opening events (histograms *B–F*).

To examine this possibility, we tested the hypothesis that pharmacological inhibition of wild-type channel activity would reduce the number of active channels and reveal smaller-amplitude events, which would be expected if many channels contribute to each large event. One micromolar  $La^{3+}$  quickly and completely blocked all activity, and this block was at least partially reversible (Fig. 4*A*). The block developed in a stepwise manner (Fig. 4*B*). The recovery of the activity occurred in similar stepwise opening events that initially flickered open and closed and then remained open (Fig. 4*C,D*). These steps occurred primarily sequentially and were not

**Figure 4.**  $\text{La}^{3+}$  block and recovery occurred in a stepwise manner. A large conductance event with very long open times was blocked completely by 1 mM  $\text{LaCl}_3$ . This block occurred in a stepwise manner (B). After several minutes of continuous wash, the activity began to reappear, occurring again in small stepwise events that were quite variable in amplitude (C, D). After many such small steps open, the events appear to become concerted, opening and closing as a single unit (E). The dashed line at the bottom of each trace represents the closed level. Two stretches of the data have been removed for clarity (at slash marks) because of the long periods of inactivity after the application of the blocker and while it remained open near the end of the top trace. The current trace begins slightly before the application of  $\text{LaCl}_3$  (bar). Subsequent wash of the patch with bath solution continued for the duration of the trace.  $V_{\text{hold}} = -40$  mV.



uniform in amplitude, nor did they correspond to the amplitude of the steps observed in the initial block. After several such stepwise events, the now large current closed nearly to zero in a single fast transition (Fig. 4E). A large conductance event dominated the record thereafter (Fig. 4A,E, latter part of trace). These results indicate that large events are not unitary channel openings but are made up of smaller conductance channels. The rapidity of transitions exhibited by the conductance in the latter part of Figure 4A suggests that the smaller channels open and close in a coordinated manner.

When we examined the action of  $10 \mu\text{M}$   $\text{La}^{3+}$ , which blocks activity much more slowly than 1 mM  $\text{La}^{3+}$ , we found a similar phenomenon (Fig. 5). Ten micromolar  $\text{La}^{3+}$  decreased the amplitudes of the conductance events without significantly changing their calculated  $P_{\text{open}}$  values (Fig. 5A,B). After initiating the wash-off of  $\text{La}^{3+}$ , we recorded conductance events of smaller amplitudes than those of the preblock events. Nevertheless,  $P_{\text{open}}$  recovered almost completely. This suggests that  $\text{La}^{3+}$  block did not occur through the obstruction of a single pore, but instead it may block many individual channels that underlie the large conductance.

## DISCUSSION

The spontaneous activity recorded from excised wild-type patches exhibited strikingly similar single-channel characteristics, despite the range of amplitudes observed from patch to patch. All were permeable to the divalent cation  $\text{Ba}^{2+}$ , were sensitive to  $\text{La}^{3+}$  block, and exhibited little or no voltage dependence. The lack of activity in rhabdomeral membranes from both *trp* null alleles and *trp*<sup>-</sup>; *trpl*<sup>-</sup> double null alleles and the reduced activity observed in the *trp*<sup>CM</sup> hypomorph suggest that the spontaneous activity we observe in wild-type and *trp*<sup>CM</sup> rhabdomeres is composed of TRP-dependent channels. Taken together, these data indicate that the spontaneous activity we observed in excised patches from isolated rhabdomeral membranes was dependent on the presence of the TRP protein.

Furthermore, these data suggest that little if any spontaneous TRPL-dependent current was present in these patches under our recording conditions. Excised patch recordings from the null mutant *trpl*<sup>p302</sup> were nearly indistinguishable from those from wild type (data not shown). It may be that TRPL-dependent channels are present in these membranes but simply do not activate spontaneously. Alternatively, these data do not address the question of whether TRPL polypeptides can be components of TRP-dependent channels through heterologous assembly.

The spontaneous activity we observed in these patches is similar to a phenomenon that develops in intact, functioning photoreceptor cells under whole-cell recording conditions (Hardie and Minke, 1994a). The rundown current (RDC) is a sustained inward current that develops minutes after the whole-cell configuration is achieved and is accompanied by a dramatic loss in light sensitivity. Although its origin is not understood, the RDC is generally thought to be caused by the dissociation of the light-activated TRP-dependent


channels from the rest of the phototransduction machinery, because it does not occur in *trp*<sup>-</sup> flies (Hardie and Minke, 1994a). Spontaneous activation of TRP channels from human has been observed in heterologous expression systems (Yamada et al., 2000).

The preponderance of small amplitude events observed in the hypomorphic *trp*<sup>CM</sup> allele suggests that the unitary conductance of individual channels is small (Fig. 2). We conclude that the large conductance events (as large as 144 pS) arise from the coordinated gating of many of these small conductance channels. There are two lines of evidence to support this idea. First, large conductance events consistently exhibit many open-state conductance transitions that are quite variable in amplitude (Fig. 3). Second, the amplitudes of large conductance events can be broken down into smaller events pharmacologically.

Determining the number of channels that open coordinately in an event requires an estimate of the single-channel conductance. Although they are rare, we have observed clear individual events of 4 pS as transitions from the closed state. Transitions of 8 pS can frequently be discerned within the open state, which contains higher-frequency noise than that of the closed state. If the unitary channel conductance is 4 pS, this would mean that 144 pS, approximately the largest wild-type conductance we measure, would correspond to 36 channels opening coordinately. This is close to the number of channels estimated to be open at the peak of a quantum bump, a single photon response (Henderson et al., 2000). Wild-type flies exhibit average bump peak amplitudes of  $\sim 12$  pA when voltage-clamped to  $-80$  mV (Cook et al., 2000). Assuming the reversal potential for each patch ( $E_{\text{rev}} = 9$  mV (Reuss et al., 1997)), this translates to  $\sim 35$  channels of 4 pS unitary conductance open at the quantum bump peak. Thus, it is possible that coordinated gating of TRP-dependent channels contributes to quantum bump formation.

In Figure 4, during the washout of  $\text{La}^{3+}$ , small unitary events initially appeared sequentially and then began to gate in a concerted manner. The transitions at the end of this trace are larger than any previous individual opening event (Fig. 4A,E, latter part of trace). These large transitions exhibit several conductance amplitudes, suggesting that coordination does not require a fixed number of elementary components. This is also the case in Figure 5. Both in the presence of a lower concentration of  $\text{La}^{3+}$  ( $10 \mu\text{M}$ ) and after wash, the amplitudes of the current were distinct from those before block by  $\text{La}^{3+}$ .

Coordinated gating may provide a mechanism for the generation or regulation of gain that appears to occur late in the phototransduction cascade. The idea that a variable number of TRP-dependent channels can be gated together to form a single unitary event provides an interesting model for regulating gain. Although the underlying gating mechanisms of these channels remains unclear, the TRP protein has been shown to be part of a supramolecular signaling assembly in the rhabdomeral membrane (Huber et al., 1996; Shieh and Zhu, 1996; Chevesich et al., 1997; Tsunoda et



**Figure 5.**  $\text{La}^{3+}$  reversibly blocked channel activity. We found that 1 mM  $\text{La}^{3+}$  blocked all conductances completely within seconds. Block by micromolar concentrations of  $\text{La}^{3+}$  was much slower. *A*, Current activity is shown before the application of the blocker. The all-points current histogram (right) was generated from the sample data (left). The  $P_{(\text{open})}$  shown in the top right corner of the histogram plot is the average  $P_{(\text{open})}$  calculated from the data shown on the left. Calculations of  $P_{(\text{open})}$  were made as described in Materials and Methods. *B*, Example of the activity seen during the application of 10  $\mu\text{M}$   $\text{LaCl}_3$ . The application of  $\text{LaCl}_3$  occurred at the beginning of the trace and continued through the entire trace. *C*, Data are shown for a patch exposed to 1 mM  $\text{LaCl}_3$ , after which activity was almost completely blocked. *D*, Activity that began to recover several seconds after we removed  $\text{LaCl}_3$ .  $V_{\text{hold}} = -20$  mV.

al., 1997). These assemblies might play a role in coordinated gating and possibly induce amplification and its regulation.

Finally, although many single-channel characteristics of TRP and TRPL channels appear to depend heavily on the environment in which they are placed, coordinated gating may explain at least one puzzling phenomenon. Coordinated gating may underlie the large range of unitary conductance values reported for the light-activated conductance (10–20 pS), the light-activated conductance in the absence of TRPL (4 pS), the RDC (1.5–4.5 pS), and TRP channels expressed in heterologous expression systems (Vaca et al., 1994; Hardie and Minke, 1994a; Hardie and Mojet, 1995; Hardie et al., 1997; Reuss et al., 1997). Although isolation procedure and recording conditions certainly do not provide an environment in which the channels normally find themselves, our approach never-

theless opens new technological means by which to gain a more complete understanding of channel function.

## REFERENCES

- Bacigalupo J, Bautista DM, Brink DL, Hetzer JF, O'Day PM (1995) Cyclic-GMP enhances light-induced excitation and induces membrane currents in *Drosophila* retinal photoreceptors. *J Neurosci* 15:7196–7200.
- Chevesich J, Kreuz AJ, Montell C (1997) Requirement for the PDZ domain protein, INAD, for localization of the TRP store-operated channel to a signaling complex. *Neuron* 18:95–105.
- Chyb S, Raghu P, Hardie RC (1999) Polyunsaturated fatty acids activate the *Drosophila* light-sensitive channels TRP and TRPL. *Nature* 397:255–259.
- Cook B, Bar-Yaacov M, Ben-Ami HC, Goldstein RE, Paroush Z, Selinger Z, Minke B (2000) Phospholipase C and termination of G-protein-mediated signalling *in vivo*. *Nat Cell Biol* 2:296–301.
- Dong Y, Kunze DL, Vaca L, Schilling WP (1995) Ins(1,4,5)P<sub>3</sub> activates *Drosophila* cation channel Trpl in recombinant baculovirus-infected Sf9 insect cells. *Am J Physiol* 269:C1332–C1339.
- Goodman MB, Art JJ (1996) Positive feedback by a potassium-selective inward rectifier enhances tuning in vertebrate hair cells. *Biophys J* 71:430–442.
- Hardie RC (1995) Photolysis of caged  $\text{Ca}^{2+}$  facilitates and inactivates but does not directly excite light-sensitive channels in *Drosophila* photoreceptors. *J Neurosci* 15:889–902.
- Hardie RC, Minke B (1992) The trp gene is essential for a light-activated  $\text{Ca}^{2+}$  channel in *Drosophila* photoreceptors. *Neuron* 8:643–651.
- Hardie RC, Minke B (1994a) Spontaneous activation of light-sensitive channels in *Drosophila* photoreceptors. *J Gen Physiol* 103:389–407.
- Hardie RC, Minke B (1994b) Calcium-dependent inactivation of light-sensitive channels in *Drosophila* photoreceptors. *J Gen Physiol* 103:409–427.
- Hardie RC, Minke B (1995) Phosphoinositide-mediated phototransduction in *Drosophila* photoreceptors: the role of  $\text{Ca}^{2+}$  and trp. *Cell Calcium* 18:256–274.
- Hardie RC, Mojet MH (1995)  $\text{Mg}^{2+}$  dependent block of the light-activated and trp dependent conductance in *Drosophila* photoreceptors. *J Neurophysiol* 74:2590–2599.
- Hardie RC, Reuss H, Lansdell SJ, Millar NS (1997) Functional equivalence of native light-sensitive channels in the *Drosophila* trp301 mutant and TRPL cation channels expressed in a stably transfected *Drosophila* cell line. *Cell Calcium* 21:431–440.
- Harteneck C, Plant T, Schultz G (2000) From worm to man: three subfamilies of TRP channels. *Trends Neurosci* 23:159–166.
- Henderson SR, Reuss H, Hardie RC (2000) Single photon responses in *Drosophila* photoreceptors and their regulation by  $\text{Ca}^{2+}$ . *J Physiol (Lond)* 524:179–194.
- Huber A, Sander P, Gobert A, Bahner M, Hermann R, Paulsen R (1996) The transient receptor potential protein (Trp), a putative store-operated  $\text{Ca}^{2+}$  channel essential for phosphoinositide-mediated photoreception, forms a signaling complex with NorpA, InaC and InaD. *EMBO J* 15:7036–7045.
- Marunaka Y, Tohda H, Hagiwara N, O'Brodvich H (1992) Cytosolic  $\text{Ca}^{2+}$ -induced modulation of ion selectivity and amiloride sensitivity of a cation channel and beta agonist action in fetal lung epithelium. *Biochem Biophys Res Commun* 187:648–656.
- Montell C (1999) Visual transduction in *Drosophila*. *Annu Rev Cell Dev Biol* 231–268.
- Niemeyer BA, Suzuki E, Scott K, Jalink K, Zuker CS (1996) The *Drosophila* light-activated conductance is composed of the two channels TRP and TRPL. *Cell* 85:651–659.
- Petersen CC, Berridge MJ, Borgese MF, Bennett DL (1995) Putative capacitative calcium entry channels: expression of *Drosophila* trp and evidence for the existence of vertebrate homologues. *Biochem J* 311:41–44.
- Phillips AM, Bull A, Kelly LE (1992) Identification of a *Drosophila* gene encoding a calmodulin-binding protein with homology to the trp phototransduction gene. *Neuron* 8:631–642.
- Putney Jr JW (1999) TRP, inositol 1,4,5-trisphosphate receptors, and capacitative calcium entry (comment). *Proc Natl Acad Sci USA* 96:14669–14671.
- Ready DF (1989) A multifaceted approach to neural development. *Trends Neurosci* 12:102–110.
- Reuss H, Mojet MH, Chyb S, Hardie RC (1997) *In vivo* analysis of the *Drosophila* light-sensitive channels, TRP and TRPL. *Neuron* 19:1249–1259.
- Sambrook J, Fritsch EF, Maniatis T (1989) Molecular cloning: a laboratory manual. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory.
- Scott K, Zuker CS (1998) Assembly of the *Drosophila* phototransduction cascade into a signalling complex shapes elementary responses. *Nature* 395:805–808.

- Scott K, Sun Y, Beckingham K, Zuker CS (1997) Calmodulin regulation of *Drosophila* light-activated channels and receptor function mediates termination of the light response in vivo. *Cell* 91:375–383.
- Shieh BH, Zhu MY (1996) Regulation of the TRP  $Ca^{2+}$  channel by INAD in *Drosophila* photoreceptors. *Neuron* 16:991–998.
- Suzuki E, Katayama E, Hirokawa K (1993) Structure of photoreceptive membranes of *Drosophila* compound eyes as studied by quick-freezing electron microscopy. *J Electron Microsc* 42:178–184.
- Tsunoda S, Sierralta J, Sun Y, Bodner R, Suzuki E, Becker A, Socolich M, Zuker CS (1997) A multivalent PDZ-domain protein assembles signaling complexes in a G-protein-coupled cascade. *Nature* 388:243–249.
- Vaca L, Sinkins WG, Hu Y, Kunze DL, Schilling WP (1994) Activation of recombinant trp by thapsigargin in Sf9 insect cells. *Am J Physiol* 267: C1501–C1505.
- Xu XS, Li H, Guggino WB, Montell C (1997) Coassembly of TRP and TRPL produces a distinct store-operated conductance. *Cell* 89:1155–1164.
- Xu X-ZS, Chien F, Butler A, Salkoff L, Montell C (2000) TRP $\gamma$ , a new *Drosophila* TRP family member, forms a regulated cation channel with TRPL. *Neuron* 26:647–657.
- Yagodin S, Hardie RC, Lansdell SJ, Millar NS, Mason WT, Sattelle DB (1998) Thapsigargin and receptor-mediated activation of *Drosophila* TRPL channels stably expressed in a *Drosophila* S2 cell line. *Cell Calcium* 23:219–228.
- Yamada H, Wakamori M, Hara Y, Takahashi Y, Konishi K, Mori Y (2000) Spontaneous single-channel activity of neuronal TRP5 channels recombinantly expressed in HEK293 cells. *Neurosci Lett* 12:111–114.
- Yue DT, Backx PH, Imredy JP (1990) Calcium-sensitive inactivation in the gating of single calcium channels. *Science* 250:1735–1738.