The Journal of Neuroscience

April 17, 2013 • Volume 33 Number 16 • www.jneurosci.org

Cover legend: Confocal projections showing ventral zebrafish olfactory bulbs at 11, 21, and 35 days after fertilization. The magnification and image settings are consistent among the three specimens shown. Olfactory glomeruli (red) were visualized with an antibody directed against synaptic vesicle protein 2 and axonal inputs from the olfactory epithelium (white) were visualized with an antibody against keyhole limpet hemocyanin. Some regions of the olfactory bulb remain virtually unchanged during this period of dramatic growth and reorganization, while alterations in other regions are influenced by olfactory experience. For more information, see the article by Braubach et al. (6905–6916).

Persons interested in becoming members of the Society for Neuroscience should contact the Membership Department, Society for Neuroscience, 1121 14th St., NW, Suite 1010, Washington, DC 20005, phone 202-962-4000.

Instructions for Authors are available at http://www.jneurosci.org/misc/itoa.shtml. Authors should refer to these Instructions online for recent changes that are made periodically.

Brief Communications Instructions for Authors are available via Internet (http://www.jneurosci.org/misc/ifa_bc.shtml).

Submissions should be submitted online using the following url: http://jneurosci.msubmit.net. Please contact the Central Office, via phone, fax, or e-mail with any questions. Our contact information is as follows: phone, 202-962-4000; fax, 202-962-4945; e-mail, jn@sfn.org.