Experience-Dependent Induction of Hippocampal Δ FosB Controls Learning

Andrew L. Eagle,¹ Paula A. Gajewski,¹ Miyoung Yang,² Megan E. Kechner,¹ Basma S. Al Masraf,¹ Pamela J. Kennedy,³ Hongbing Wang,¹ Michelle S. Mazei-Robison,¹ and Alfred J. Robison¹

¹Department of Physiology and Neuroscience Program, Michigan State University, East Lansing, Michigan 48824, ²Department of Anatomy, School of Medicine, Wonkwang University, Iksan, Jeonbuk, 570–749, South Korea, and ³Department of Psychology, University of California Los Angeles, Los Angeles, California 90095

The hippocampus (HPC) is known to play an important role in learning, a process dependent on synaptic plasticity; however, the molecular mechanisms underlying this are poorly understood. Δ FosB is a transcription factor that is induced throughout the brain by chronic exposure to drugs, stress, and variety of other stimuli and regulates synaptic plasticity and behavior in other brain regions, including the nucleus accumbens. We show here that Δ FosB is also induced in HPC CA1 and DG subfields by spatial learning and novel environmental exposure. The goal of the current study was to examine the role of Δ FosB in hippocampal-dependent learning and memory and the structural plasticity of HPC synapses. Using viral-mediated gene transfer to silence Δ FosB transcriptional activity by expressing Δ JunD (a negative modulator of Δ FosB transcriptional function) or to overexpress Δ FosB, we demonstrate that HPC Δ FosB regulates learning and memory. Specifically, Δ JunD expression in HPC impaired learning and memory on a battery of hippocampal-dependent tasks in mice. Similarly, general Δ FosB overexpression also impaired learning. Δ JunD expression in HPC did not affect anxiety or natural reward, but Δ FosB overexpression induced anxiogenic behaviors, suggesting that Δ FosB may mediate attentional gating in addition to learning. Finally, we found that overexpression of Δ FosB increases immature dendritic spines on CA1 pyramidal cells, whereas Δ JunD reduced the number of immature and mature spine types, indicating that Δ FosB may exert its behavioral effects through modulation of HPC synaptic function. Together, these results suggest collectively that Δ FosB plays a significant role in HPC cellular morphology and HPC-dependent learning and memory.

Key words: Δ FosB; dendritic spines; hippocampus; learning; transcription

Significance Statement

Consolidation of our explicit memories occurs within the hippocampus, and it is in this brain region that the molecular and cellular processes of learning have been most closely studied. We know that connections between hippocampal neurons are formed, eliminated, enhanced, and weakened during learning, and we know that some stages of this process involve alterations in the transcription of specific genes. However, the specific transcription factors involved in this process are not fully understood. Here, we demonstrate that the transcription factor Δ FosB is induced in the hippocampus by learning, regulates the shape of hippocampal synapses, and is required for memory formation, opening up a host of new possibilities for hippocampal transcriptional regulation.

Introduction

Hippocampal synaptic plasticity is critical for memory consolidation, and many immediate early genes (IEGs) engender epigenetic, transcriptional, and direct effector changes that regulate neuroplasticity and memory consolidation, including *Zif268*,

Arc, *c-Fos*, and *FosB* (Guzowski, 2002; Shepherd and Bear, 2011; Alberini and Kandel, 2015). IEGs are driven by neural activity and are induced robustly in a variety of regions in response to stress (Cullinan et al., 1995; Kovács, 2008), drugs (Hope et al., 1992; Larson et al., 2010; Guez-Barber et al., 2011), and other environmental stimuli, such as experiential events (Lanahan and Worley, 1998; Guzowski et al., 2001; Stack et al., 2010). Δ FosB is

Received May 30, 2015; revised Aug. 18, 2015; accepted Sept. 5, 2015.

Author contributions: A.L.E., M.Y., P.J.K., H.W., M.S.M.-R., and A.J.R. designed research; A.L.E., P.A.G., M.E.K., B.S.A.M., P.J.K., and A.J.R. performed research; A.L.E. and A.J.R. analyzed data; A.L.E. and A.J.R. wrote the paper.

This research was supported by Whitehall Foundation Grant 2013-08-43 (A.J.R.), Multidisciplinary Training in Environmental Toxicology Training Grant T32-ES007255 (A.L.E.), and a 2014 National Alliance for Research on Schizophrenia and Depression Young Investigator Award from the Brain and Behavior Research Foundation (A.L.E.). We thank Kenneth Moon for outstanding technical support.

The authors declare no competing financial interests.

Correspondence should be addressed to Alfred J. Robison, 3180 Biomedical Physical Sciences, 567 Wilson Road, East Lansing, MI 48824. E-mail: robiso45@msu.edu.

DOI:10.1523/JNEUROSCI.2083-15.2015

Copyright © 2015 the authors 0270-6474/15/3513773-11\$15.00/0

a transcription factor within the Fos family that is known to regulate synaptic plasticity in brain reward regions, such as the nucleus accumbens (NAc), prefrontal cortex, and ventral tegmental area (VTA; Nestler et al., 1999; Robison and Nestler, 2011). It is unique among the IEGs in that it is stably induced by chronic stimuli, such as repeated stress (Nikulina et al., 2008; Nestler, 2015), chronic antidepressant treatment (Vialou et al., 2010; Thibault et al., 2014), and repeated administration to drugs of abuse (Nestler et al., 1999; Perrotti et al., 2008; Robison and Nestler, 2011). Whereas the induction of most Fos family proteins is transient, Δ FosB has a half-life of ~ 8 d *in vivo* (Carle et al., 2007; Ulery-Reynolds et al., 2009), making it an ideal candidate mechanism for long-term changes in gene expression resulting from chronic stimuli.

In the NAc, Δ FosB is known to regulate gene transcription of key target molecules associated with cellular growth and synaptic plasticity, including GluA2 receptor (Kelz et al., 1999), calcium/calmodulin-dependent kinase II (CaMKII; Robison et al., 2013), nuclear factor- κ B (Ang et al., 2001), and cyclin-dependent protein kinase 5 (Cdk5; Bibb et al., 2001). Considering the nature of these targets, it is not surprising that Δ FosB regulates the number and morphology of NAc dendritic spines (Maze et al., 2010; Grueter et al., 2013), as well as NAc synaptic strength and the apparent number of silent synapses (Grueter et al., 2013). Although many of these gene targets are also critical to hippocampal synaptic plasticity and morphology, as well as spatial learning, the role of Δ FosB in these processes remains unknown.

Previous evidence indicates that Δ FosB is induced robustly in the hippocampus by drugs of abuse (Perrotti et al., 2008), and we reported recently that it is also induced in the hippocampus by antidepressants (Vialou et al., 2015). Importantly, Δ FosB induction in the NAc is dependent on the function of the transcription factors cAMP response element-binding protein (CREB) and serum response factor (SRF; Vialou et al., 2012), both of which are essential for hippocampal function and are induced by spatial learning (Mizuno et al., 2002; Porte et al., 2008; Alberini and Kandel, 2015). Moreover, genetic knock-out of the *FosB* gene causes malformation of the hippocampus and impairment in spatial memory (Solecki et al., 2008; Yutsudo et al., 2013). However, because *FosB* is an IEG and important for development (Redemann-Fibi et al., 1991), the effects of hippocampal Δ FosB on learning and memory are impossible to ascertain using *FosB* null mutants, because the genetic deletion of *FosB* impairs early hippocampal development and adult neurogenesis (Yutsudo et al., 2013). Therefore, we sought to use temporally and spatially discrete viral-mediated manipulation of Δ FosB expression and activity to assess the role of Δ FosB in the adult hippocampus. Thus, the goal of the present study is to characterize the induction of hippocampal Δ FosB by learning and to identify any role of Δ FosB in hippocampal cell morphology or learning and memory.

Materials and Methods

Animals. The study followed guidelines described in the *Guide for the Care and Use of Laboratory Animals*, eighth edition (Institute of Laboratory Animal Resources, 2011). Before any testing, all experimental procedures were approved by the Institutional Animal Care and Use Committee at Michigan State University or at the Mount Sinai School of Medicine. After arrival to the facility, male mice ($n = 4$ per cage) or rats ($n = 2$ per cage) were group housed for at least 3 d before experimentation in a 12 h light/dark cycle with *ad libitum* food and water. Animals were weighed and briefly handled daily before and throughout the study.

Surgery. Stereotaxic surgery was performed to inject viral vectors into the hippocampus of adult male mice. Thirty gauge needles (Hamilton Company) were bilaterally placed at the following coordi-

nates: 10° angle; -2.2 mm anteroposterior (AP); ± 2.5 mm mediolateral. Purified high-titer virus ($0.6 \mu\text{l}$) was infused separately ($0.3 \mu\text{l}/\text{infusion}$) over 3 min periods at two sites: -2.1 mm dorsoventral (DV) and -1.9 mm DV. After both infusions, the needles remained at the injection site for a 5 min after infusion to allow diffusion of the viral particles. Previously validated (Robison et al., 2013) viral vectors included the following: adeno-associated virus (AAV2) expressing GFP alone (AAV-GFP), AAV expressing GFP and Δ FosB (AAV- Δ FosB), AAV expressing GFP and Δ JunD (AAV- Δ JunD), herpes simplex virus (HSV) expressing GFP alone (HSV-GFP), HSV expressing GFP and Δ FosB (HSV- Δ FosB), or HSV expressing GFP and Δ JunD (HSV- Δ JunD). HSVs were obtained from the Massachusetts Institute of Technology Viral Core Facility, and AAVs were obtained from the University of North Carolina Vector Core.

Open-field activity. The open-field (OF) apparatus consisted of a custom-made, square white polyvinylchloride foam box ($38 \times 38 \times 35$ cm). Animals were tested for 30 min, and activity was recorded with a digital CCD camera connected to a computer running an automated video tracking software package (Clever Sys).

Novel object recognition. Novel object recognition (NOR) was assessed using a 3 d paradigm including habituation, training, and testing. Mice were first habituated to the OF for 1 h, and OF activity was recorded at this time. Twenty-four hours later, two similar objects were placed near the corners of the square apparatus, and animals were allowed to explore the apparatus for 30 min. Objects pairs consisted of metal and plastic objects, including Lego blocks, miniature action figures, spark plugs, and metal knobs. Mice did not show consistent biases for any one object over another. The following day, mice were tested for NOR. One object was removed and replaced with another dissimilar object, and mice were allowed to freely explore the apparatus for 5 min. Behavior was video recorded, and time spent in the 7 cm square corners around the objects was assessed.

Contextual fear conditioning. Contextual fear conditioning (FC) consisted of a 2 d procedure: conditioning and testing. For conditioning, mice were placed into an operant chamber equipped with a house light, metal grid floor, and footshock stimulus generator (Coulbourn Instruments) for 180 s. This was followed by the delivery of three mild electric footshocks (0.8 mA, 1 s duration), each separated by a 60 s intershock interval in which no shock was delivered. Mice remained in the chamber for another 60 s after the last delivery. Twenty-four hours after conditioning, mice were placed back into the conditioning chamber for 8 min. The percentage of time spent freezing was video scored by a blind, independent observer. Freezing was defined as the lack of skeletal movement for periods > 1 s.

Temporally dissociated passive avoidance learning paradigm. Temporally dissociated passive avoidance (TDPA) was assessed across 5 d and was performed as described previously (Zhang et al., 2008) with slight modifications. During each daily session, the mouse was placed into the lit half of a light–dark passive avoidance chamber (Coulbourn Instruments) and allowed to explore for 1 min before a door was opened to allow entry into the dark half of the chamber. After entry into the dark half, the door was closed and a 5 min dissociation period was allowed to pass before delivery of a mild electric footshock (0.8 mA, 2 s duration). The animal was returned to its home cage 30 s after the footshock. The crossover latency (in seconds) to enter the dark half after door opening was scored across daily sessions as an index of learning. Maximal latency was set at 600 s.

Elevated plus maze. Elevated plus maze (EPM) behavior was measured using a custom-built apparatus (aluminum base with polyvinylchloride arms) based on plans from ANY-maze (www.anymaze.com; Stoelting). The maze was laid out in the shape of a plus sign as viewed from above. It consisted of four gray-colored, interconnected runways (5×35 cm) with two open and two closed arms, elevated 45 cm from the ground. Closed arms were enclosed on the long sides of the runway by two 15 cm high gray walls. Animals were placed into the center of the maze facing an open arm, and behavior was video recorded as described above for 5 min. The percentage of time spent in the open arms and the percentage of entries into the open arms were quantified as assessments of anxiety-like behavior.

Two-bottle choice for sucrose preference. A standard two-bottle choice procedure was assessed across 4 d. Singly housed mice were first given access to two bottles of drinking water on the top of their home cage for 3 d to assess baseline drinking and to acclimate the mice to the bottles. Then, mice were given two-bottle choice, in which one of the bottles was replaced with a 1% sucrose solution. Sucrose solution consumed per day as a percentage of total liquid consumption was measured as an index of anhedonia.

Morris water maze. The rat Morris water maze (MWM) was performed essentially as described previously (Morris et al., 1982). The pool of water (~ 1.3 m diameter, 26°C) was made opaque with nontoxic white tempera paint, and platforms (8 cm diameter) were placed 0.3 m from the wall either 1 cm below the surface (hidden) or 2 cm above the surface (visible). Rats were given six training sessions per day for 5 consecutive days, and a probe trial was performed on day 6. If a rat failed to find the hidden platform in 60 s, it was placed onto the platform. Rats remained on the platform for 15 s to allow spatial orientation at the end of each trial. A computerized tracking system (Datawave Technologies) recorded and analyzed swim-path data, including latency to reach the platform (in seconds), time in each quadrant, and the average swim speed.

Mice were assayed in the MWM as described previously (Zhang and Wang, 2013). Briefly, behavior was measured using an automated video-tracking system (WaterMaze; Coulbourn Instruments) in a water-filled pool with the surface covered with white polypropylene beads and a platform. Procedures included visible platform training (six trials, day 1), hidden platform training (four trials per day; days 2–6), and a probe test (day 7). During visible platform training, a flag was attached to the platform and placed randomly at different locations for each trial. During hidden platform training, the flag was removed and the platform was always in the same location across trials. Trials lasted 60 s or when the mouse reached the platform, whichever came first. Intertrial intervals lasted 60 min. Trial latencies and swim speed (centimeters per second) were recorded during training. During the probe test, the platform was removed and mice were allowed to freely explore the maze for 60 s. The time spent in each quadrant was recorded.

Novel environment exposure. Mice were removed from their home cage and placed in a novel enriched environment along with their cage mates (five mice per cage) for 60 min/d for 10 consecutive days. The novel environment consisted of a round polycarbonate chamber 48 inches across with 8-inch walls containing four equal connected subdivisions, each of which contained different novel objects similar to those used for the NOR test (described above). Low lighting was used, and the experimenter stayed outside of the room to minimize stress.

Western blotting for Δ FosB. Brains were extracted rapidly on cold ice and then sliced into 1 mm sections, and the dorsal hippocampus was removed with a 12 gauge punch and frozen immediately on dry ice. Samples were then processed for SDS-PAGE and transferred to PVDF membranes for Western blotting with chemiluminescence. Blots were probed for FosB (5G4; 1:500; Cell Signaling Technology) and assayed for total protein using Swift Membrane Stain (G Biosciences). Protein was quantified using NIH ImageJ software.

Immunohistochemical assay for FosB. Animals were perfused transcardially with ice-cold PBS, followed by 10% Formalin. Brains were post-fixed 24 h in 10% Formalin, cryopreserved in 30% sucrose, and then sliced into 35 μm sections. Immunohistochemistry was performed using anti-FosB primary antibody (sc-48; 1:500; Santa Cruz Biotechnology) and visualized by 3,3'-diaminobenzidine (DAB) staining (Vector Laboratories).

Immunofluorescence for FosB and JunD. Animals were perfused and brains sliced as above. Immunofluorescence was performed using mouse anti-FosB (ab11959; 1:500; Abcam) or mouse anti-JunD (ab28837; 1:500; Abcam) and a goat anti-GFP (ab5450; 1:1000; Abcam) primary antibody and corresponding secondary antibodies (705-545-147 and 711-165-152; 1:200; Jackson ImmunoResearch). Fluorescent images were visualized on an Olympus FluoView 1000 filter-based laser scanning confocal microscope.

Spine analysis. Animals were perfused as above, and brains were sliced into 100 μm sections. Sections were mounted on slides using DPX mounting medium (Sigma-Aldrich), and GFP fluorescence was visual-

ized using an Olympus FluoView 1000 filter-based laser scanning confocal microscope. Spines were analyzed essentially as described previously (Christoffel et al., 2011). Briefly, dendritic segments 50–150 μm away from the soma were chosen randomly from HSV-infected cells that express GFP. Images were acquired on a confocal LSM 710 (Carl Zeiss) for morphological analysis using NeuronStudio with the rayburst algorithm. NeuronStudio classifies spines as thin, mushroom, or stubby based on the following values: (1) aspect ratio; (2) head/neck ratio; and (3) head diameter. Spines with a neck can be classified as either thin or mushroom, and those without a significant neck are classified as stubby. Spines with a neck are labeled as thin or mushroom based on head diameter.

Statistical analysis. All statistical analyses were performed using Prism software (GraphPad Software). Anxiety phenotypic behavior (EPM and OF activity), MWM probe test performance (in mice), contextual FC freezing, protein quantification, and dendritic spine number were analyzed using Student's *t* tests. MWM training performance (in rats) was analyzed using one-way repeated-measures (RM) ANOVA, followed by Holm–Sidak corrected *post hoc* comparisons for simple effects. Sucrose preference (percentage), NOR, MWM training performance (mice), TDPA latency, and body mass were analyzed using two-way RM ANOVAs. This was followed by Holm–Sidak-corrected *post hoc* comparisons for simple effects. Alpha criterion of 0.05 was set for all analyses.

Results

Induction of hippocampal Δ FosB by novel environmental exposure and spatial learning

A variety of stimuli have been reported to induce Δ FosB in many brain regions, including the hippocampus (Nikulina et al., 2008; Robison and Nestler, 2011; Nestler, 2015), but it is unknown whether exposure to complex environments and spatial learning, the primary function of the rodent hippocampus, induces Δ FosB in this region. Therefore, we exposed mice to a novel, enriched environment for 60 min/d for 10 d and compared the expression of Δ FosB in the hippocampus with that of animals remaining in their home cages. As expected, exposure to the novel environment significantly induced Δ FosB protein in the dorsal hippocampus (Fig. 1A; $t_{(15)} = 2.90$, $p = 0.011$).

We also aimed to ascertain whether specific spatial learning could induce Δ FosB and whether this effect was species specific. Rats or mice were trained on a standard MWM paradigm in cued learning using a visible platform, which does not require the dorsal hippocampus, or in spatial learning using a nonvisible platform, which requires the use of the hippocampus to learn orientation based on spatial cues around the maze (Vorhees and Williams, 2006). Animals readily acquired the ability to locate the hidden platform across days, showing a decrease in escape latency each day (Fig. 1B, rats: $F_{(2,122,10,61)} = 7.864$, $p = 0.0075$; significant on days 2 and 5 by RM ANOVA, followed by Holm–Sidak multiple comparison test; Fig. 1C, mice: $F_{(4,000,28,00)} = 13.92$, $p < 0.0001$; significant on days 3–5). Immunohistochemical analysis for FosB-positive (FosB⁺) expressing (FosB/ Δ FosB) cells in the hippocampus (Fig. 1D) revealed induction of FosB⁺ cells in the CA1 of spatial learning rats compared with naive animals and cued learning animals (Table 1). Because tissue was extracted 1 d after the last training, it is likely that the FosB⁺ cells represent the presence of Δ FosB, because full-length FosB expression is transient, i.e., <24 h (Carle et al., 2007). We also noted that the pattern of FosB⁺ cells in the CA1 region of spatial learning animals was sparse, with only a few very dark cells (Fig. 1D, red arrow), possibly consistent with the encoding of individual memories during learning. We also observed that both spatial and cued learning increased FosB⁺ cells in the dentate gyrus (DG) of both mice and rats (Table 1), perhaps indicating that the stress of exposure to the MWM or the novel environment of the testing chamber is sufficient to induce Δ FosB in this region. This is un-

Figure 1. Induction of Δ FosB in the hippocampus by novelty exposure and spatial learning. **A**, Quantitation and representative Western blots (inset) of Δ FosB in the hippocampus show significant increases after novel environmental exposure (red; $n = 9$) compared with mice left in their home cage (white; $n = 8$; $p < 0.05$). **B**, Representative paths of rats seeking a visible (cued) or hidden (spatial) platform in the MWM test. **C**, Escape latency from the MWM in rats (left) and mice (right) using a hidden platform significantly decreased across training days ($n = 6-8$; $p < 0.05$ compared to day 1). **D**, Representative $40\times$ images of coronal sections stained for FosB gene products from naive, cued learning, and spatial learning rats (left) and mice (right). Red arrowheads indicate FosB⁺ pyramidal cells in CA1.

surprising, because we have reported previously that chronic stress can induce Δ FosB in the DG (Vialou et al., 2014, 2015), and we saw a similar induction of Δ FosB in the rat NAc (Table 1), a region in which Δ FosB is also induced by stress (Vialou et al., 2010, 2014, 2015). Induction in the CA3 region was inconsistent between species and not correlated with swimming stress (Table 1), indicating that Δ FosB in CA3 may be unrelated to learning or stress. These findings collectively suggest that exposure to a novel environment and the process of spatial learning robustly induce Δ FosB in the hippocampus.

Transcriptional silencing of hippocampal Δ FosB impairs learning and memory

Because learning was found to induce Δ FosB, we hypothesized that silencing Δ FosB-mediated activator protein-1 transcriptional activity in the hippocampus would produce spatial learning and memory impairment. We used viral vector-mediated (AAV2 or HSV) expression of GFP and Δ FosB, GFP and the Δ FosB inhibitor Δ JunD, or GFP alone (as control), targeted to the dorsal hippocampus of mice, and examined behavior using a battery of tests that are associated with hippocampal function. Using AAV, gene transfer of GFP was detected in both pyramidal

and nonpyramidal cells in CA3 and CA1 and granule and non-granule cells in DG subregions of the hippocampus (Fig. 2A), whereas HSV expression was more clearly restricted to pyramidal and granule cells (Fig. 2A,C,D), as has been reported previously (Burger et al., 2004). We validated the specific overexpression of Δ FosB in transduced neurons by immunofluorescence (Fig. 2C) and noted that basal expression of FosB was present in pyramidal neurons and in microglia likely activated by the injection-related tissue damage, as suggested by recent findings (Nomaru et al., 2014). We also validated the viral overexpression of Δ FosB by quantitative PCR of tissue punches from injected hippocampus and found a 271.2 ± 74.5 -fold increase in Δ FosB message in tissue transduced with HSV-GFP- Δ FosB over tissue transduced with HSV-GFP alone ($t_{(13)} = 3.9$, $p = 0.0018$), which is not unexpected when using the cytomegalovirus promoter (Papadakis et al., 2004).

We specifically examined learning and memory behavior across a battery of tests including: NOR memory, contextual FC freezing behavior, TDPA learning, and spatial learning in the MWM. In the NOR task (Fig. 3A), as expected, control mice (AAV-GFP) spent more time with a novel object than a familiar one ($t_{(7)} = 2.54$, $p = 0.038$). However, mice that received AAV-GFP- Δ JunD spent equal time with both objects, indicating impairment in recognition memory ($p > 0.05$). We next assessed memory for contextual fear in the same group of mice (Fig. 3B). We found that mice expressing Δ JunD spent less time freezing compared with GFP controls ($t_{(14)} = 2.41$, $p = 0.030$), indicating impairment in memory for contextual fear.

TDPA learning (Fig. 3C) was assessed in a separate cohort of mice ($n = 16$ per group) to test context-associative fear learning using a paradigm that has been demonstrated previously to be impaired by hippocampal lesions (Zhang et al., 2008). We observed that, although both GFP and Δ JunD groups increased their latency to cross over to the footshock-associated “dark” side over time ($F_{\text{day}(4,120)} = 21.24$, $p < 0.001$), Δ JunD mice were significantly impaired ($F_{\text{group}(1,30)} = 8.58$, $p = 0.007$; $F_{\text{day} \times \text{group}(4,120)} = 5.22$, $p = 0.001$) in the formation of this association between the dark side of the apparatus and the pain of the footshock. Specifically, Δ JunD mice had significantly attenuated crossover latency at days 4 and 5 ($p = 0.001$ for both days), indicating that hippocampal Δ FosB transcriptional activity is required for passive avoidance learning.

Spatial learning was assessed in another cohort by performance in the MWM. Escape latencies decreased across days of hidden platform training for both groups (Fig. 3D; $F_{(4,56)} = 8.44$, $p < 0.001$), but only nonsignificant differences were observed between groups ($p > 0.05$). Swim speed also decreased across days for both groups (data not shown; $F_{(4,56)} = 12.3$, $p < 0.001$), but no group differences were observed ($p > 0.05$). A probe test was then conducted with the platform removed. When measured in the probe test, Δ JunD-expressing mice exhibited longer latency to find the target area where the platform had been ($t_{(14)} = 3.56$, $p = 0.003$) and spent significantly less time in the target quadrant compared with GFP control mice (Fig. 3E; $t_{(14)} = 2.21$, $p = 0.044$). In addition, Δ JunD mice had fewer crossings over the (previous) location of the hidden platform (data not shown; $t_{(14)} = 2.23$, $p = 0.043$). These data suggest that Δ FosB transcriptional activity in the hippocampus is required for spatial learning and reference memory.

Because AAV-mediated gene transfer requires up to 2 weeks to reach peak expression and thus experiments are performed 3 weeks after surgery, it is possible that the effects of AAV-mediated silencing of Δ FosB transcriptional activity are attributable to ho-

Table 1. Induction of Δ FosB by spatial and cued learning

Brain region	Control cells/mm ² (mean $\pm$ SEM) (n)	Cued cells/mm ² (mean $\pm$ SEM) (n)	Spatial cells/mm ² (mean $\pm$ SEM) (n)	F (one-way ANOVA)	p (one-way ANOVA)
Rats					
CA1	2.10 $\pm$ 0.66 (5)	2.88 $\pm$ 0.43 (4)	5.10 $\pm$ 1.00 (5)*	$F_{(2,11)} = 4.316$	0.0413
CA3	3.50 $\pm$ 0.67 (5)	6.00 $\pm$ 1.02 (4)	4.90 $\pm$ 0.91 (5)	$F_{(2,11)} = 2.038$	0.176
DG	17.70 $\pm$ 0.75 (5)	30.88 $\pm$ 4.04 (4)*	40.10 $\pm$ 4.05 (5)*	$F_{(2,11)} = 13.02$	0.0013
NAC	58.17 $\pm$ 7.15 (6)	93.42 $\pm$ 4.17 (6)*	89.83 $\pm$ 7.95 (6)*	$F_{(2,15)} = 8.582$	0.0033
Mice					
CA1	69.15 $\pm$ 11.0 (8)	92.64 $\pm$ 12.7 (8)	143.9 $\pm$ 19.5 (7)*	$F_{(2,20)} = 6.752$	0.0057
CA3	533.7 $\pm$ 60.2 (8)	844.9 $\pm$ 70.9 (8)*	736.1 $\pm$ 56.3 (7)	$F_{(2,20)} = 6.448$	0.0069
DG	1222 $\pm$ 67.4 (8)	1893 $\pm$ 83.8 (8)*	1586 $\pm$ 131.9 (7)*	$F_{(2,20)} = 13.08$	0.0002

FosB⁺ cells as visualized by DAB staining (Fig. 1) in control, cued learning, and spatial learning mice and rats by brain region. Statistics represent one-way ANOVA with Tukey's multiple comparison *post hoc* test used to compare conditions. **p* < 0.05, significant difference from the control group.

Figure 2. Viral-mediated gene transfer in the hippocampus. **A**, Representative fluorescent coronal images of the hippocampus (approximately -2.0 mm AP) 6 d after injection of HSV-GFP or 28 d after injection of AAV-GFP. **B**, Coronal figures (-1.94 and -2.46 AP; Paxinos and Franklin, 2012) showing representative center fluorescence distribution from GFP (green circles), GFP- Δ JunD (orange circles), or GFP- Δ FosB (blue circles) infusions. Immunofluorescence of CA1 neurons close to the injection site of HSV-GFP and HSV-GFP- Δ FosB (**C**) or HSV-GFP- Δ JunD (**D**) reveals specific transgene expression only in cells induced with virus.

meostatic adaptations to long-term changes in Δ FosB-mediated gene expression (Janson et al., 2001). To determine the acute effects of silencing Δ FosB transcriptional activity, we used HSV-mediated gene transfer, which begins within hours of surgery and lasts 5–7 d, allowing behavioral assessment within 2–5 d of surgery (Fig. 2). Using a cohort of mice that received hippocampal injections of HSV-GFP or HSV-GFP- Δ JunD we found that, in the NOR test, control GFP mice spent more time with a novel object (Fig. 3F; $t_{(15)} = 4.36$, $p = 0.001$), whereas HSV-GFP- Δ JunD mice did not ($p > 0.05$). Similarly, when we tested con-

Figure 3. Transcriptional silencing of hippocampal Δ FosB impairs learning and memory. **A**, NOR in mice that received stereotaxic hippocampal injections of AAV-GFP (GFP, green; $n = 8$) and AAV-GFP- Δ JunD (Δ JunD, orange; $n = 8$). The duration of time spent with a familiar object (blank bars) and a novel object (crosshatched bars) was measured. GFP mice spent significantly more time with the novel object than the familiar object ($*p < 0.05$ compared with familiar), whereas the Δ JunD mice spent an equal amount of time with both objects. **B**, Contextually fear conditioned (Context FC) freezing (seconds) in AAV-GFP ($n = 8$) and AAV-GFP- Δ JunD ($n = 8$) mice. Δ JunD significantly decreased contextual freezing ($*p < 0.05$ compared with GFP). **C**, TDPA performance was measured in mice that received AAV-GFP ($n = 16$) or AAV-GFP- Δ JunD ($n = 16$). Latency (seconds) to crossover to the shock-paired dark side of the box was measured. Δ JunD mice had significantly lower crossover latency at days 4–5 ($**p < 0.01$, $***p < 0.001$, respectively, compared with GFP). **D**, Spatial learning in the MWM was measured after AAV-GFP ($n = 8$) or AAV-GFP- Δ JunD ($n = 8$). Escape latency (seconds) did not differ between groups across training days. **E**, The probe test revealed that Δ JunD mice spent significantly less time in the target quadrant ($*p < 0.05$ compared with GFP). **F**, Mice received stereotaxic hippocampal infusions of HSV-GFP ($n = 16$) or HSV-GFP- Δ JunD ($n = 8$). GFP mice spent significantly more time with the novel object than the familiar object ($*p < 0.001$ compared with familiar), whereas the Δ JunD mice spent an equal amount of time with both objects. **G**, Context FC freezing after HSV-GFP ($n = 8$) and HSV-GFP- Δ JunD ($n = 8$). Δ JunD mice spent significantly less time freezing ($*p < 0.05$ compared with HSV-GFP). Error bars indicate mean $\pm$ SEM.

text FC behavior, we found that HSV-GFP- Δ JunD mice spent less time freezing compared with GFP mice (Fig. 3G; $t_{(14)} = 2.28$, $p = 0.039$). Thus, these data were consistent with AAV results and suggest that memory formation requires acute Δ FosB-mediated changes in hippocampal gene expression.

Overexpression of Δ FosB in hippocampus also induces impairments in learning and memory

Spatial memories must be distinct to be recalled without interference, and recent studies indicate that discrete hippocampal networks allow for storage of large quantities of uncorrelated spatial

Figure 4. Overexpression of Δ FosB in the hippocampus impairs learning and memory. **A**, NOR in mice that received stereotaxic hippocampal injections of AAV-GFP (GFP, green; $n = 14$) or AAV-GFP- Δ FosB (Δ FosB, blue; $n = 16$). GFP mice spent significantly more time with the novel object than the familiar object ($*p < 0.05$ compared with familiar), whereas the Δ FosB mice spent an equal amount of time with both objects. **B**, Contextually fear conditioned (Context FC) freezing (seconds) in AAV-GFP ($n = 8$) and AAV-GFP- Δ FosB ($n = 8$) mice. Δ FosB mice had significantly greater contextual freezing ($*p < 0.05$ compared with GFP). **C**, TDPA performance was measured in mice that received AAV-GFP ($n = 15$) or AAV-GFP- Δ FosB ($n = 16$). Δ FosB mice had significantly lower crossover latency at days 4–5 ($*p < 0.05$, $**p < 0.01$, respectively, compared with GFP). **D**, Spatial learning in the MWM was measured after AAV-GFP ($n = 7$) or AAV-GFP- Δ FosB ($n = 7$). Escape latency (seconds) was significantly decreased by Δ FosB at days 3–4 ($**p < 0.01$ for both days compared with GFP). **E**, Probe test quadrant time revealed that Δ FosB mice spent significantly less time in the target quadrant ($*p < 0.05$ compared with GFP). Error bars indicate mean $\pm$ SEM.

information (Moser et al., 2015). In fact, in rodents, a spatial location or memory can be linked directly to the plasticity of single cells or small groups of cells within the hippocampal-entorhinal circuit (Moser et al., 2015), suggesting that changes in gene expression mediating that plasticity must be equally discrete. Therefore, we examined how indiscriminate viral overexpression of Δ FosB in hippocampus would affect learning and memory. We again used AAV to overexpress Δ FosB (or GFP as control) in the hippocampus of mice, and mice were tested across the same battery of learning tasks. As expected, control GFP mice expressing only GFP spent more time with a novel object (Fig. 4A; $t_{(26)} = 2.89$, $p = 0.008$), but mice overexpressing Δ FosB were significantly impaired in recognition of the novel object ($p > 0.05$). In context FC, Δ FosB-overexpressing mice spent significantly less time freezing than GFP control mice (Fig. 4B; $t_{(14)} = 2.32$, $p = 0.036$). These results indicate that overexpression of Δ FosB, similar to silencing Δ FosB, impairs recognition and contextual fear memory.

In TDPA, we replicated findings from the Δ JunD experiments (Fig. 4C). Both GFP control and Δ FosB-overexpressing mice increased their crossover latency ($F_{\text{day}(4,84)} = 17.27$, $p < 0.001$). However, Δ FosB overexpression produced impairment, i.e., reduction, in crossover latency ($F_{\text{group}(1,21)} = 4.41$, $p = 0.048$; $F_{\text{day} \times \text{group}(4,84)} = 3.06$, $p = 0.021$), specifically at days 4 ($p = 0.038$) and 5 ($p = 0.010$). The effects of hippocampal Δ FosB overexpression on spatial learning were assessed in a separate cohort in the MWM. Escape latency (Fig. 4D) decreased in the

GFP control group ($F_{\text{group}(1,12)} = 7.00$, $p = 0.021$), with significant differences between GFP controls and Δ FosB overexpressing mice at days 3 and 4 ($F_{\text{day} \times \text{group}(4,48)} = 4.39$, $p = 0.004$; $p = 0.002$ for day 3; $p = 0.005$ for day 4). Additionally, although trial speed decreased across days for both groups (data not shown; $F_{\text{day}(4,48)} = 22.31$, $p < 0.001$), Δ FosB-overexpressing mice were significantly slower ($F_{\text{group}(1,12)} = 7.53$, $p = 0.018$), specifically at days 2–4 compared with GFP controls ($F_{\text{day} \times \text{group}(4,48)} = 4.05$, $p = 0.007$; $p < 0.05$ for days 2–4). During the probe test, no differences were observed in latency to target ($p > 0.05$), but Δ FosB-overexpressing mice spent significantly less time in the target quadrant (Fig. 4E; $t_{(12)} = 2.67$, $p = 0.021$). No differences were observed between groups in the number of crossings over the (previous) location of the hidden platform (data not shown; $p > 0.05$). These findings, along with the results from NOR and contextual fear memory, collectively indicate that nonselective overexpression of Δ FosB in the hippocampus impairs learning and memory. This further suggests that discrete expression of Δ FosB in cells directly encoding a given memory and only in those cells is required for proper learning.

Overexpression, but not silencing, of Δ FosB in hippocampus induces anxiety-like behavior

Δ FosB transcriptional activity in the NAc is linked to mood and anxiety (Vialou et al., 2010), and the hippocampus is associated directly with maladaptive stress responses, leading to depression and/or anxiety disorders (Sapolsky, 2001; Campbell and Macqueen, 2004; Shin and Liberzon, 2010). Therefore, we assessed whether silencing Δ FosB transcriptional activity in the hippocampus would lead to anxiety and depressive phenotypes. Mice that received AAV-GFP, AAV-GFP- Δ JunD, or AAV-GFP- Δ FosB were assayed across a battery of tests to examine anxiety-like and depressive phenotypes, including OF behavior, EPM, sucrose preference, and body weight. Thigmotaxis (e.g., avoidance of the center) behavior in the OF was found to be different between groups. Δ FosB overexpression, but not Δ JunD expression, decreased the time spent in the center of the OF (Fig. 5A; $F_{(2,29)} = 9.12$, $p < 0.001$). No differences were observed in center entries (Fig. 5B; $p > 0.05$). Activity, e.g., distance moved, was also unaffected by Δ FosB overexpression or silencing (Fig. 5C; $p > 0.05$), indicating that locomotor activity is unaltered by hippocampal Δ FosB. Similar to the OF test, behavior in the EPM showed an anxiogenic phenotype after Δ FosB overexpression. Specifically, Δ FosB overexpression, but not Δ JunD expression, decreased both the time spent in the open arms (Fig. 5D; $F_{(2,29)} = 3.59$, $p = 0.040$; GFP vs Δ FosB, $p = 0.025$) and entries into the open arms (Fig. 5E; $F_{(2,29)} = 6.64$, $p = 0.004$; GFP vs Δ FosB, $p = 0.004$). No differences were observed in activity, e.g., distance moved, in the EPM (Fig. 5F; $p > 0.05$). These results suggest collectively that indiscriminate hippocampal Δ FosB overexpression, but not silencing of Δ FosB transcriptional activity, induces anxiety-like behavior.

Anhedonia, such as decreased preference for natural rewards, and perturbations in body weight are hallmarks of clinical depression and can be modeled in mice (Nestler and Hyman, 2010). Thus, another cohort of mice was assayed for sucrose preference (percentage) using the two-bottle choice procedure to assess anhedonia. All mice (AAV-GFP, AAV-GFP- Δ JunD, and AAV-GFP- Δ FosB) increased preference for the bottle containing the 1% sucrose solution compared with baseline (Fig. 5G; $F_{\text{day}(4,120)} = 54.81$, $p < 0.001$), and no differences between groups ($p > 0.05$) were observed in baseline drinking or preference for sucrose solution. Finally, mice were

Figure 5. Overexpression, but not silencing, of Δ FosB in hippocampus produces anxiety-like behavior. **A**, OF behavior was assessed in AAV-GFP (green; $n = 16$), AAV-GFP- Δ JunD (orange; $n = 8$), and AAV-GFP- Δ FosB (blue; $n = 8$) mice. Center zone time (percentage) was reduced significantly by Δ FosB ($*p < 0.001$ compared with GFP), but no differences were observed between GFP and Δ JunD. **B**, Center zone entries were not different between groups. **C**, Locomotor activity in an OF, measured by distance moved (meters), was not different between groups. **D**, Open arm time (percentage) in an EPM was decreased in mice that received Δ FosB ($*p < 0.05$ compared with GFP). **E**, Open arm entries were also reduced in mice that received Δ FosB ($**p < 0.01$ compared with GFP). **F**, Distance moved (meters) in the EPM was not different between groups. **G**, Preference for sucrose (percentage) increased in all groups compared with a sucrose-free baseline ($***p < 0.001$ compared with baseline; **B**). No group differences were observed in sucrose preference. **H**, Body weight (grams) was measured across weeks after AAV surgery ($n = 15$ –16 per group). Although mice steadily increased total body weight, no differences were observed between groups (dotted lines represent 95% confidence interval). Error bars indicate mean $\pm$ SEM.

assayed for changes in body weight, and no group differences were observed (Fig. 5H; $p > 0.05$). Thus, neither silencing nor overexpression of hippocampal Δ FosB induces a depressive (anhedonic) phenotype, nor does it produce body weight perturbations.

Hippocampal Δ FosB controls dendritic spine morphology

Δ FosB induces changes in cellular physiology and morphology in NAc D₁ medium spiny neurons (Grueter et al., 2013). In particular, NAc Δ FosB induction increases the number of silent synapses along with a corresponding increase in immature dendritic spines. Therefore, we used HSV-mediated overexpression of GFP, GFP- Δ FosB, or GFP- Δ JunD in the mouse hippocampus to determine the effects of Δ FosB on hippocampal dendritic spine morphology. We found that overexpression of Δ FosB caused an

Figure 6. Δ FosB regulates hippocampal spine formation. Mice received hippocampal infusions of HSV-GFP, HSV-GFP- Δ FosB, or HSV-GFP- Δ JunD and dendritic spine analysis was conducted on hippocampal CA1 pyramidal neurons. **A**, Representative dendrites showing spines from GFP⁺ hippocampal CA1 neurons. **B**, Quantitation of spines in CA1 neurons shows Δ FosB (blue; $n = 30$) had significantly more total spines, thin spines, and stubby spines compared with GFP alone (gray; $n = 30$; $*p < 0.001$), with no significant difference in mushroom spines. Δ JunD (red; $n = 19$) had significantly fewer total spines, thin spines, and mushroom spines compared with GFP alone (gray; $n = 22$; $*p < 0.001$), with no significant difference in stubby spines. Error bars indicate mean $\pm$ SEM. **C**, Model depicting experience-dependent changes in hippocampal cells under control conditions or when virally transduced with Δ JunD (red) or Δ FosB (blue). Green indicates changes in synaptic structure/function resulting from Δ FosB expression within a circuit that may underlie learning.

increase in total spines in CA1 hippocampal neurons (Fig. 6A,B; $t_{(38)} = 8.26$, $p < 0.001$). Furthermore, we found that, although Δ FosB did not increase the number of mushroom spines ($p > 0.05$), it did increase the number of thin ($t_{(38)} = 4.97$, $p < 0.001$) and stubby ($t_{(38)} = 7.20$, $p < 0.001$) spines, indicating that Δ FosB overexpression induces immature spines in hippocampal neurons. In contrast, expression of Δ JunD caused a decrease in total spines (Fig. 6A,B; $t_{(39)} = 4.09$, $p < 0.001$) that was driven by decreases in thin ($t_{(39)} = 4.18$, $p < 0.001$) and mushroom ($t_{(39)} = 3.09$, $p = 0.004$) spines but did not affect stubby spines ($p > 0.05$). These data indicate that Δ FosB regulates basal spine formation and stabilization in CA1, although levels of Δ FosB in CA1 neurons in the absence of learning were quite low (Fig. 1, Table 1). Combined with the effects of Δ FosB overexpression, this indicates that Δ FosB modulates the number and function of hippocampal synapses, a possible mechanism for its effects on learning and memory.

Discussion

We demonstrate for the first time that the transcription factor Δ FosB is important for hippocampal-dependent learning and

memory. Specifically, we observed that silencing the transcriptional activity of hippocampal Δ FosB impaired learning and memory across a battery of hippocampal-dependent memory tasks. It is likely that, during learning and exposure to novel environments, groups of hippocampal cells encoding an engram for a particular association express Δ FosB and that the expression of Δ FosB mediates downstream signaling events that prepare the cell for synaptic morphological and physiological changes associated with memory consolidation. This hypothesis is further bolstered by the fact that Δ FosB in the NAc is known to directly regulate the expression of target genes that are essential for hippocampal synaptic plasticity and learning (Guzman-Karlsson et al., 2014), such as CDK5 (Kumar et al., 2005), CaMKII, and GluA receptors (Robison and Nestler, 2011).

These candidate targets of hippocampal Δ FosB have been studied extensively in the hippocampus. For example, CDK5 regulates neural development and neurogenesis (Lagace et al., 2008; Su and Tsai, 2011), glutamatergic neurotransmission (Morabito et al., 2004), and synaptic plasticity (Bibb, 2003; Dhariwala and Rajadhyaksha, 2008; Su and Tsai, 2011). CaMKII has long been known to be a molecular substrate of synaptic plasticity and memory in the hippocampus (Shonesy et al., 2014), and its activity is necessary for long-term potentiation induction (Malinow et al., 1989), AMPA receptor trafficking and function (Malinow and Malenka, 2002), and structural remodeling of the synapse (Okamoto et al., 2007). In addition, these activities, and interactions with NMDA receptors and other postsynaptic proteins (Robison et al., 2005a; Robison et al., 2005b), may contribute to a variety of neuropsychiatric diseases (Robison, 2014). Specifically in D_1 medium spiny neurons of the NAc, Δ FosB regulates the induction of CaMKII gene expression by cocaine and that CaMKII provides a feedforward loop via its direct phosphorylation of a stability site on Δ FosB (Robison et al., 2013), which suggests a close relationship between these two molecules in addition. Δ FosB overexpression in inducible bitransgenic mice has also been shown previously to increase GluA2 levels (Kelz et al., 1999), and Δ FosB overexpression increases the number of AMPA-receptor-lacking silent synapses in D_1 medium spiny neurons of the NAc (Grueter et al., 2013). This is significant in that AMPA receptor activity and expression is related directly to the hippocampal synaptic activity and plasticity critical for memory formation (Guzman-Karlsson et al., 2014).

Despite this evidence, a gap in knowledge exists for the specific contribution of hippocampal Δ FosB to memory formation. Early reports suggested the possibility of such a contribution, because Δ FosB is expressed highly within the hippocampus and electrical stimulation of this region induces Δ FosB mRNA expression (Cole et al., 1990; Nakabeppu and Nathans, 1991). In addition, *FosB* knock-out mice display malformation of the hippocampus, reduced hippocampal neurogenesis, and spatial learning deficits (Yutsudo et al., 2013). These data indicate that *FosB* is important for development of the hippocampus and perhaps adult neurogenesis but, because of the nature of the knock-out model, do not test the acute role of *FosB* in learning. The current finding that Δ FosB regulates spine formation in CA1 hippocampal neurons suggests that Δ FosB plays an important role in synaptic plasticity, the first evidence that hippocampal Δ FosB plays a role in the function of fully differentiated hippocampal neurons. It will now be important to test whether experience-dependent (e.g., learning-induced) changes in hippocampal spine morphology require Δ FosB.

The specific mechanisms for the induction of Δ FosB by learning or novel environmental exposure are currently unknown, but

again, previous studies from the NAc provide a model. For example, CREB and SRF mediate the induction by cocaine of Δ FosB in NAc (Vialou et al., 2012) and are essential in hippocampal function and learning (Knöll and Nordheim, 2009; Alberini and Kandel, 2015). Furthermore, the induction of Δ FosB may be regulated by dopamine from the VTA, which provides a signal to hippocampal neurons relating the salience and novelty of events (Lisman and Grace, 2005). Dopamine has been shown to enhance plasticity in CA1 pyramidal neurons (Li et al., 2003) and, more recently, to mediate network-level activity and memory persistence (McNamara et al., 2014). Indeed, this may explain why drugs of abuse, such as cocaine, are able to produce strong induction of Δ FosB in hippocampus (Perrotti et al., 2008) and may also explain the strong associative memories of drug effects and drug-taking contexts.

Interestingly, overexpression of Δ FosB in the hippocampus produced impairments in learning and memory similar to those observed with inhibition of FosB transcriptional activity. Although it may be hypothesized that increased Δ FosB expression is associated with memory consolidation and learning, the current finding may be explained by an overall decrease in the signal-to-noise ratio. We hypothesize that *FosB* gene products accumulate specifically in the relatively few cells of the hippocampus encoding a particular memory, i.e., the engram (Mayford, 2014; Fig. 1 and model in Fig. 6). Thus, non-specific overexpression of Δ FosB throughout entire regions of the hippocampus masks the specific effects of Δ FosB in the engram neurons, losing their signal in the “noise” created by overexpression. The new memory circuits are not consolidated because they cannot be distinguished from the other nonrelevant circuits artificially expressing Δ FosB. New evidence is beginning to emerge that synchronous neural networks within the hippocampus may underlie memory consolidation (Patel, 2015), consistent with the view that the hippocampus encodes memories via specific networks (Abraham, 2008; Neves et al., 2008), supporting this signal-to-noise hypothesis. Alternatively, the impairment may be attributable to aberrant hippocampal network reorganization, because seizures, which are known to induce both Δ FosB (Chen et al., 1995) and its target CDK5 (Chen et al., 2000), also induce reorganization of granule cells in the DG (Parent et al., 1997). Finally, it may be that global overexpression of Δ FosB interferes with homeostatic mechanisms characteristic of hippocampal networks (Pozo and Goda, 2010). However, all of these possibilities hint that non-selective Δ FosB overexpression may be a detriment to intrahippocampal network-level activity and plasticity, which may impinge on memory consolidation.

Δ FosB overexpression also caused an increase in immature dendritic spines in the hippocampus, as has been reported previously with overexpression in the NAc (Grueter et al., 2013; Robinson et al., 2013). This may be indicative of a priming effect, in which Δ FosB accumulation causes the neuron to produce immature synapses that are ready to be consolidated rapidly into mature mushroom synapses after addition neuronal activation, i.e., new learning. This may also indicate a mechanism by which circuits that have been activated repeatedly by particularly salient stimuli may remain poised for recall, for instance, those encoding associations between drugs of abuse and the context of drug exposure that appear to underlie relapse to addiction behaviors (Crombag et al., 2008; Lasseter et al., 2010). However, it is important to note that general overexpression of Δ FosB in the hippocampus does not promote learning but rather impairs it. Thus, if Δ FosB primes synapses for rapid learning, it must require additional cell-specific signaling events to be effective.

We also found increased anxiety as a result of global Δ FosB overexpression in the hippocampus. The current findings are in line with the theory of Gray and McNaughton (2000) for hippocampal involvement in anxiety, with new optogenetic evidence indicating an important contribution of the hippocampus (for review, see Fournier and Duman, 2013). Global Δ FosB overexpression may impair attentional gating, specifically the ability to attribute salience to aversive cues. In addition, animals that attribute too much salience to nonpredictive environmental cues tend to also place high negative valence on aversive cues (Morrow et al., 2011). A clinical example of this may be observed in post-traumatic stress disorder (PTSD), in which afflicted individuals are unable to differentiate safety signals from threatening stimuli (Jovanovic et al., 2012). Safety signal encoding and contextual conditioning are regulated by the amygdala and hippocampus (Wiltgen et al., 2006; Alvarez et al., 2008; Marschner et al., 2008; Sangha et al., 2013). Therefore, the lack of safety signal discrimination in clinical PTSD may be attributed to hippocampal dysfunction and overactive amygdala activity (Pitman et al., 2012). Similarly, in the current study, Δ FosB overexpression throughout the hippocampus may prevent proper hippocampal regulation of amygdalar activity, leading to the observed increase in anxiety-like behaviors. It is interesting to note that chronic social defeat stress, one model of PTSD and anxiety, also promotes Δ FosB expression throughout hippocampus (Vialou et al., 2015). However, this hippocampal Δ FosB is unlikely to be the sole cause of anxiety in the social defeat model, because hippocampal Δ FosB also accumulates with chronic antidepressant treatment (Vialou et al., 2015), a regimen that does not produce anxiety.

The current findings indicate that learning induces Δ FosB in specific cells encoding memories in the hippocampus and suggest that this expression is essential for learning. This highlights the need for large-scale candidate gene and omics approaches to identify hippocampal Δ FosB gene targets to uncover the mechanism by which Δ FosB regulates hippocampal cell and synaptic function. This novel role for Δ FosB is particularly relevant, because it suggests that, by impairing the functions of hippocampal cells active during memory formation, we might impair a specific memory. Thus, Δ FosB and its target genes could be therapeutic targets for neuropsychiatric diseases associated with impaired memory, such as depression, PTSD, and addiction.

References

- Abraham WC (2008) Metaplasticity: tuning synapses and networks for plasticity. *Nat Rev Neurosci* 9:387. [CrossRef Medline](#)
- Alberini CM, Kandel ER (2015) The regulation of transcription in memory consolidation. *Cold Spring Harb Perspect Biol* 7:a021741. [CrossRef Medline](#)
- Alvarez RP, Biggs A, Chen G, Pine DS, Grillon C (2008) Contextual fear conditioning in humans: cortical-hippocampal and amygdala contributions. *J Neurosci* 28:6211–6219. [CrossRef Medline](#)
- Ang E, Chen J, Zagouras P, Magna H, Holland J, Schaeffer E, Nestler EJ (2001) Induction of nuclear factor- κ B in nucleus accumbens by chronic cocaine administration. *J Neurochem* 79:221–224. [CrossRef Medline](#)
- Bibb JA (2003) Role of Cdk5 in neuronal signaling, plasticity, and drug abuse. *Neurosignals* 12:191–199. [CrossRef Medline](#)
- Bibb JA, Chen J, Taylor JR, Svenningsson P, Nishi A, Snyder GL, Yan Z, Sagawa ZK, Ouimet CC, Nairn AC, Nestler EJ, Greengard P (2001) Effects of chronic exposure to cocaine are regulated by the neuronal protein Cdk5. *Nature* 410:376–380. [CrossRef Medline](#)
- Burger C, Gorbatyuk OS, Velardo MJ, Peden CS, Williams P, Zolotukhin S, Reier PJ, Mandel RJ, Muzyczka N (2004) Recombinant AAV viral vectors pseudotyped with viral capsids from serotypes 1, 2, and 5 display differential efficiency and cell tropism after delivery to different regions of the central nervous system. *Mol Ther* 10:302–317. [CrossRef Medline](#)
- Campbell S, Macqueen G (2004) The role of the hippocampus in the patho-

- physiology of major depression. *J Psychiatry Neurosci* 29:417–426. [Medline](#)
- Carle TL, Ohnishi YN, Ohnishi YH, Alibhai IN, Wilkinson MB, Kumar A, Nestler EJ (2007) Proteasome-dependent and -independent mechanisms for FosB destabilization: identification of FosB deproton domains and implications for DeltaFosB stability. *Eur J Neurosci* 25:3009–3019. [CrossRef Medline](#)
- Chen J, Nye HE, Kelz MB, Hiroi N, Nakabeppu Y, Hope BT, Nestler EJ (1995) Regulation of delta FosB and FosB-like proteins by electroconvulsive seizure and cocaine treatments. *Mol Pharmacol* 48:880–889. [Medline](#)
- Chen J, Zhang Y, Kelz MB, Steffen C, Ang ES, Zeng L, Nestler EJ (2000) Induction of cyclin-dependent kinase 5 in the hippocampus by chronic electroconvulsive seizures: role of Δ FosB. *J Neurosci* 20:8965–8971. [Medline](#)
- Christoffel DJ, Golden SA, Dumitriu D, Robison AJ, Janssen WG, Ahn HF, Krishnan V, Reyes CM, Han MH, Ables JL, Eisch AJ, Dietz DM, Ferguson D, Neve RL, Greengard P, Kim Y, Morrison JH, Russo SJ (2011) I κ B kinase regulates social defeat stress-induced synaptic and behavioral plasticity. *J Neurosci* 31:314–321. [CrossRef Medline](#)
- Cole AJ, Abu-Shakra S, Saffen DW, Baraban JM, Worley PF (1990) Rapid rise in transcription factor mRNAs in rat brain after electroshock-induced seizures. *J Neurochem* 55:1920–1927. [CrossRef Medline](#)
- Crombag HS, Bossert JM, Koya E, Shaham Y (2008) Review. Context-induced relapse to drug seeking: a review. *Philos Trans R Soc Lond B Biol Sci* 363:3233–3243. [CrossRef Medline](#)
- Cullinan WE, Herman JP, Battaglia DF, Akil H, Watson SJ (1995) Pattern and time course of immediate early gene expression in rat brain following acute stress. *Neuroscience* 64:477–505. [CrossRef Medline](#)
- Dhariwala FA, Rajadhyaksha MS (2008) An unusual member of the cdk family: Cdk5. *Cell Mol Neurobiol* 28:351–369. [CrossRef Medline](#)
- Fournier NM, Duman RS (2013) Illuminating hippocampal control of fear memory and anxiety. *Neuron* 77:803–806. [CrossRef Medline](#)
- Gray JA, McNaughton N (2000) The neuropsychology of anxiety, Ed 2. New York: Oxford UP.
- Grueter BA, Robison AJ, Neve RL, Nestler EJ, Malenka RC (2013) FosB differentially modulates nucleus accumbens direct and indirect pathway function. *Proc Natl Acad Sci U S A* 110:1923–1928. [CrossRef Medline](#)
- Guez-Barber D, Fanous S, Golden SA, Schrama R, Koya E, Stern AL, Bossert JM, Harvey BK, Picciotto MR, Hope BT (2011) FACS identifies unique cocaine-induced gene regulation in selectively activated adult striatal neurons. *J Neurosci* 31:4251–4259. [CrossRef Medline](#)
- Guzman-Karlsson MC, Meadows JP, Gavin CF, Hablitz JJ, Sweatt JD (2014) Transcriptional and epigenetic regulation of Hebbian and non-Hebbian plasticity. *Neuropharmacology* 80:3–17. [CrossRef Medline](#)
- Guzowski JF (2002) Insights into immediate-early gene function in hippocampal memory consolidation using antisense oligonucleotide and fluorescent imaging approaches. *Hippocampus* 12:86–104. [CrossRef Medline](#)
- Guzowski JF, Setlow B, Wagner EK, McGaugh JL (2001) Experience-dependent gene expression in the rat hippocampus after spatial learning: a comparison of the immediate-early genes Arc, c-fos, and zif268. *J Neurosci* 21:5089–5098. [Medline](#)
- Hope B, Kosofsky B, Hyman SE, Nestler EJ (1992) Regulation of immediate early gene expression and AP-1 binding in the rat nucleus accumbens by chronic cocaine. *Proc Natl Acad Sci U S A* 89:5764–5768. [CrossRef Medline](#)
- Institute of Laboratory Animal Resources (2011) Guide for the care and use of laboratory animals, Ed 8. Washington, DC: National Academy of Sciences.
- Janson CG, McPhee SW, Leone P, Freese A, During MJ (2001) Viral-based gene transfer to the mammalian CNS for functional genomic studies. *Trends Neurosci* 24:706–712. [CrossRef Medline](#)
- Jovanovic T, Kazama A, Bachevalier J, Davis M (2012) Impaired safety signal learning may be a biomarker of PTSD. *Neuropharmacology* 62:695–704. [CrossRef Medline](#)
- Kelz MB, Chen J, Carlezon WA Jr, Whisler K, Gilden L, Beckmann AM, Steffen C, Zhang YJ, Marotti L, Self DW, Tkatch T, Baranaukas G, Surmeier DJ, Neve RL, Duman RS, Picciotto MR, Nestler EJ (1999) Expression of the transcription factor Δ FosB in the brain controls sensitivity to cocaine. *Nature* 401:272–276. [CrossRef Medline](#)
- Knöll B, Nordheim A (2009) Functional versatility of transcription factors in the nervous system: the SRF paradigm. *Trends Neurosci* 32:432–442. [CrossRef Medline](#)
- Kovács KJ (2008) Measurement of immediate-early gene activation: c-fos and beyond. *J Neuroendocrinol* 20:665–672. [CrossRef Medline](#)
- Kumar A, Choi KH, Renthall W, Tsankova NM, Theobald DE, Truong HT, Russo SJ, Laplant Q, Sasaki TS, Whistler KN, Neve RL, Self DW, Nestler EJ (2005) Chromatin remodeling is a key mechanism underlying cocaine-induced plasticity in striatum. *Neuron* 48:303–314. [CrossRef Medline](#)
- Lagace DC, Benavides DR, Kansy JW, Mapelli M, Greengard P, Bibb JA, Eisch AJ (2008) Cdk5 is essential for adult hippocampal neurogenesis. *Proc Natl Acad Sci U S A* 105:18567–18571. [CrossRef Medline](#)
- Lanahan A, Worley P (1998) Immediate-early genes and synaptic function. *Neurobiol Learn Mem* 70:37–43. [CrossRef Medline](#)
- Larson EB, Akkentli F, Edwards S, Graham DL, Simmons DL, Alibhai IN, Nestler EJ, Self DW (2010) Striatal regulation of DeltaFosB, FosB, and cFos during cocaine self-administration and withdrawal. *J Neurochem* 115:112–122. [CrossRef Medline](#)
- Lasseter HC, Xie X, Ramirez DR, Fuchs RA (2010) Sub-region specific contribution of the ventral hippocampus to drug context-induced reinstatement of cocaine-seeking behavior in rats. *Neuroscience* 171:830–839. [CrossRef Medline](#)
- Li S, Cullen WK, Anwyl R, Rowan MJ (2003) Dopamine-dependent facilitation of LTP induction in hippocampal CA1 by exposure to spatial novelty. *Nat Neurosci* 6:526–531. [CrossRef Medline](#)
- Lisman JE, Grace AA (2005) The hippocampal-VTA loop: controlling the entry of information into long-term memory. *Neuron* 46:703–713. [CrossRef Medline](#)
- Malinow R, Malenka RC (2002) AMPA receptor trafficking and synaptic plasticity. *Annu Rev Neurosci* 25:103–126. [CrossRef Medline](#)
- Malinow R, Schulman H, Tsien RW (1989) Inhibition of postsynaptic PKC or CaMKII blocks induction but not expression of LTP. *Science* 245:862–866. [CrossRef Medline](#)
- Marschner A, Kalisch R, Vervliet B, Vansteenwegen D, Büchel C (2008) Dissociable roles for the hippocampus and the amygdala in human cued versus context fear conditioning. *J Neurosci* 28:9030–9036. [CrossRef Medline](#)
- Mayford M (2014) The search for a hippocampal engram. *Philos Trans R Soc Lond B Biol Sci* 369:20130161. [Medline](#)
- Maze I, Covington HE 3rd, Dietz DM, LaPlant Q, Renthall W, Russo SJ, Mechanic M, Mouzon E, Neve RL, Haggarty SJ, Ren Y, Sampath SC, Hurd YL, Greengard P, Tarakhovskiy A, Schaefer A, Nestler EJ (2010) Essential role of the histone methyltransferase G9a in cocaine-induced plasticity. *Science* 327:213–216. [CrossRef Medline](#)
- McNamara CG, Tejero-Cantero Á, Trouche S, Campo-Urriza N, Dupret D (2014) Dopaminergic neurons promote hippocampal reactivation and spatial memory persistence. *Nat Neurosci* 17:1658–1660. [CrossRef Medline](#)
- Mizuno M, Yamada K, Maekawa N, Saito K, Seishima M, Nabeshima T (2002) CREB phosphorylation as a molecular marker of memory processing in the hippocampus for spatial learning. *Behav Brain Res* 133:135–141. [CrossRef Medline](#)
- Morabito MA, Sheng M, Tsai LH (2004) Cyclin-dependent kinase 5 phosphorylates the N-terminal domain of the postsynaptic density protein PSD-95 in neurons. *J Neurosci* 24:865–876. [CrossRef Medline](#)
- Morris RG, Garrud P, Rawlins JN, O'Keefe J (1982) Place navigation impaired in rats with hippocampal lesions. *Nature* 297:681–683. [CrossRef Medline](#)
- Morrow JD, Maren S, Robinson TE (2011) Individual variation in the propensity to attribute incentive salience to an appetitive cue predicts the propensity to attribute motivational salience to an aversive cue. *Behav Brain Res* 220:238–243. [CrossRef Medline](#)
- Moser MB, Rowland DC, Moser EI (2015) Place cells, grid cells, and memory. *Cold Spring Harb Perspect Biol* 7:a021808. [CrossRef Medline](#)
- Nakabeppu Y, Nathans D (1991) A naturally occurring truncated form of FosB that inhibits Fos/Jun transcriptional activity. *Cell* 64:751–759. [CrossRef Medline](#)
- Nestler EJ (2015) FosB: a transcriptional regulator of stress and antidepressant responses. *Eur J Pharmacol* 753:66–72. [CrossRef Medline](#)
- Nestler EJ, Hyman SE (2010) Animal models of neuropsychiatric disorders. *Nat Neurosci* 13:1161–1169. [CrossRef Medline](#)
- Nestler EJ, Kelz MB, Chen J (1999) Δ FosB: a molecular mediator of long-term neural and behavioral plasticity. *Brain Res* 835:10–17. [CrossRef Medline](#)
- Neves G, Cooke SF, Bliss TV (2008) Synaptic plasticity, memory and the

- hippocampus: a neural network approach to causality. *Nat Rev Neurosci* 9:65–75. [CrossRef Medline](#)
- Nikulina EM, Arrillaga-Romany I, Miczek KA, Hammer RP Jr (2008) Long-lasting alteration in mesocorticolimbic structures after repeated social defeat stress in rats: time course of μ -opioid receptor mRNA and FosB/ Δ FosB immunoreactivity. *Eur J Neurosci* 27:2272–2284. [CrossRef Medline](#)
- Nomaru H, Sakumi K, Katogi A, Ohnishi YN, Kajitani K, Tsuchimoto D, Nestler EJ, Nakabeppu Y (2014) FosB gene products contribute to excitotoxic microglial activation by regulating the expression of complement C5a receptors in microglia. *Glia* 62:1284–1298. [CrossRef Medline](#)
- Okamoto K, Narayanan R, Lee SH, Murata K, Hayashi Y (2007) The role of CaMKII as an F-actin-bundling protein crucial for maintenance of dendritic spine structure. *Proc Natl Acad Sci U S A* 104:6418–6423. [CrossRef Medline](#)
- Papadakis ED, Nicklin SA, Baker AH, White SJ (2004) Promoters and control elements: designing expression cassettes for gene therapy. *Curr Gene Ther* 4:89–113. [CrossRef Medline](#)
- Parent JM, Yu TW, Leibowitz RT, Geschwind DH, Sloviter RS, Lowenstein DH (1997) Dentate granule cell neurogenesis is increased by seizures and contributes to aberrant network reorganization in the adult rat hippocampus. *J Neurosci* 17:3727–3738. [Medline](#)
- Patel J (2015) Network mechanisms underlying the initiation and generation of sharp-wave-associated ripple oscillations. *J Neurosci* 35:2323–2325. [CrossRef Medline](#)
- Paxinos G, Franklin KB (2012) The mouse brain in stereotaxic coordinates, Ed 4. San Diego: Academic.
- Perrotti LI, Weaver RR, Robison B, Renthall W, Maze I, Yazdani S, Elmore RG, Knapp DJ, Selley DE, Martin BR, Sim-Selley L, Bachtell RK, Self DW, Nestler EJ (2008) Distinct patterns of Δ FosB induction in brain by drugs of abuse. *Synapse* 62:358–369. [CrossRef Medline](#)
- Pitman RK, Rasmussen AM, Koenen KC, Shin LM, Orr SP, Gilbertson MW, Milad MR, Liberzon I (2012) Biological studies of post-traumatic stress disorder. *Nat Rev Neurosci* 13:769–787. [CrossRef Medline](#)
- Porte Y, Buhot MC, Mons NE (2008) Spatial memory in the Morris water maze and activation of cyclic AMP response element-binding (CREB) protein within the mouse hippocampus. *Learn Mem* 15:885–894. [CrossRef Medline](#)
- Pozo K, Goda Y (2010) Unraveling mechanisms of homeostatic synaptic plasticity. *Neuron* 66:337–351. [CrossRef Medline](#)
- Redemann-Fibi B, Schuermann M, Müller R (1991) Stage and tissue-specific expression of fosB during mouse development. *Differentiation* 46:43–49. [CrossRef Medline](#)
- Robison AJ (2014) Emerging role of CaMKII in neuropsychiatric disease. *Trends Neurosci* 37:653–662. [CrossRef Medline](#)
- Robison AJ, Nestler EJ (2011) Transcriptional and epigenetic mechanisms of addiction. *Nat Rev Neurosci* 12:623–637. [CrossRef Medline](#)
- Robison AJ, Bartlett RK, Bass MA, Colbran RJ (2005a) Differential modulation of Ca^{2+} /calmodulin-dependent protein kinase II activity by regulated interactions with N-methyl-D-aspartate receptor NR2B subunits and alpha-actinin. *J Biol Chem* 280:39316–39323. [CrossRef Medline](#)
- Robison AJ, Bass MA, Jiao Y, MacMillan LB, Carmody LC, Bartlett RK, Colbran RJ (2005b) Multivalent interactions of calcium/calmodulin-dependent protein kinase II with the postsynaptic density proteins NR2B, densin-180, and alpha-actinin-2. *J Biol Chem* 280:35329–35336. [CrossRef Medline](#)
- Robison AJ, Vialou V, Mazei-Robison M, Feng J, Kourrich S, Collins M, Wee S, Koob G, Turecki G, Neve R, Thomas M, Nestler EJ (2013) Behavioral and structural responses to chronic cocaine require a feedforward loop involving Δ FosB and calcium/calmodulin-dependent protein kinase II in the nucleus accumbens shell. *J Neurosci* 33:4295–4307. [CrossRef Medline](#)
- Sangha S, Chadick JZ, Janak PH (2013) Safety encoding in the basal amygdala. *J Neurosci* 33:3744–3751. [CrossRef Medline](#)
- Sapolsky RM (2001) Depression, antidepressants, and the shrinking hippocampus. *Proc Natl Acad Sci U S A* 98:12320–12322. [CrossRef Medline](#)
- Shepherd JD, Bear MF (2011) New views of Arc, a master regulator of synaptic plasticity. *Nat Neurosci* 14:279–284. [CrossRef Medline](#)
- Shin LM, Liberzon I (2010) The neurocircuitry of fear, stress, and anxiety disorders. *Neuropsychopharmacology* 35:169–191. [CrossRef Medline](#)
- Shonesy BC, Jalan-Sakrkar N, Cavenier VS, Colbran RJ (2014) CaMKII: a molecular substrate for synaptic plasticity and memory, Chap 3. In: *Progress in molecular biology and translational science* (Zafar UK, Muly EC, eds), pp 61–87. San Diego: Academic.
- Solecki W, Krowka T, Kubik J, Kaczmarek L, Przewlocki R (2008) Role of fosB in behaviours related to morphine reward and spatial memory. *Behav Brain Res* 190:212–217. [CrossRef Medline](#)
- Stack A, Carrier N, Dietz D, Hollis F, Sorenson J, Kabbaj M (2010) Sex differences in social interaction in rats: role of the immediate-early gene zif268. *Neuropsychopharmacology* 35:570–580. [CrossRef Medline](#)
- Su SC, Tsai LH (2011) Cyclin-dependent kinases in brain development and disease. *Annu Rev Cell Dev Biol* 27:465–491. [CrossRef Medline](#)
- Thibault MA, Eagle AL, Kaska S, Nestler EJ, Mazei-Robison MS, Vialou V, Robison AJ (2014) Region-specific induction of FosB isoforms in mouse brain after stress or chronic fluoxetine exposure. *Soc Neurosci Abstr* 40:426.13.
- Ulery-Reynolds PG, Castillo MA, Vialou V, Russo SJ, Nestler EJ (2009) Phosphorylation of DeltaFosB mediates its stability in vivo. *Neuroscience* 158:369–372. [CrossRef Medline](#)
- Vialou V, Robison AJ, Laplant QC, Covington HE 3rd, Dietz DM, Ohnishi YN, Mouzon E, Rush AJ 3rd, Watts EL, Wallace DL, Iniguez SD, Ohnishi YH, Steiner MA, Warren BL, Krishnan V, Bolaños CA, Neve RL, Ghose S, Berton O, Tammenga CA, Nestler EJ (2010) DeltaFosB in brain reward circuits mediates resilience to stress and antidepressant responses. *Nat Neurosci* 13:745–752. [CrossRef Medline](#)
- Vialou V, Feng J, Robison AJ, Ku SM, Ferguson D, Scobie KN, Mazei-Robison MS, Mouzon E, Nestler EJ (2012) Serum response factor and cAMP response element binding protein are both required for cocaine induction of Δ FosB. *J Neurosci* 32:7577–7584. [CrossRef Medline](#)
- Vialou V, Bagot RC, Cahill ME, Ferguson D, Robison AJ, Dietz DM, Fallon B, Mazei-Robison M, Ku SM, Harrigan E, Winstanley CA, Joshi T, Feng J, Berton O, Nestler EJ (2014) Prefrontal cortical circuit for depression- and anxiety-related behaviors mediated by cholecystokinin: role of Δ FosB. *J Neurosci* 34:3878–3887. [CrossRef Medline](#)
- Vialou V, Thibault M, Kaska S, Cooper S, Gajewski P, Eagle A, Mazei-Robison M, Nestler EJ, Robison AJ (2015) Differential induction of FosB isoforms throughout the brain by fluoxetine and chronic stress. *Neuropharmacology* 99:28–37. [CrossRef Medline](#)
- Vorhees CV, Williams MT (2006) Morris water maze: procedures for assessing spatial and related forms of learning and memory. *Nat Protoc* 1:848–858. [CrossRef Medline](#)
- Wiltgen BJ, Sanders MJ, Anagnostaras SG, Sage JR, Fanselow MS (2006) Context fear learning in the absence of the hippocampus. *J Neurosci* 26:5484–5491. [CrossRef Medline](#)
- Yutsudo N, Kamada T, Kajitani K, Nomaru H, Katogi A, Ohnishi YH, Ohnishi YN, Takase K, Sakumi K, Shigeto H, Nakabeppu Y (2013) fosB-null mice display impaired adult hippocampal neurogenesis and spontaneous epilepsy with depressive behavior. *Neuropsychopharmacology* 38:895–906. [CrossRef Medline](#)
- Zhang M, Wang H (2013) Mice overexpressing type 1 adenylyl cyclase show enhanced spatial memory flexibility in the absence of intact synaptic long-term depression. *Learn Mem* 20:352–357. [CrossRef Medline](#)
- Zhang M, Moon C, Chan GC, Yang L, Zheng F, Conti AC, Muglia L, Muglia LJ, Storm DR, Wang H (2008) Ca-stimulated type 8 adenylyl cyclase is required for rapid acquisition of novel spatial information and for working/episodic-like memory. *J Neurosci* 28:4736–4744. [CrossRef Medline](#)