

Journal Club

Editor's Note: These short reviews of recent *JNeurosci* articles, written exclusively by students or postdoctoral fellows, summarize the important findings of the paper and provide additional insight and commentary. If the authors of the highlighted article have written a response to the Journal Club, the response can be found by viewing the Journal Club at www.jneurosci.org. For more information on the format, review process, and purpose of Journal Club articles, please see <http://www.jneurosci.org/content/jneurosci-journal-club>.

ZFPM1 Necessary for Development of Serotonergic Projections Related to Anxiety and Contextual Fear Learning

Masakazu Taira and Miles J. Desforges

Neural Computation Unit, Okinawa Institute of Science and Technology Graduate University, Onna-son 904-0495, Okinawa, Japan

Review of Tikker et al.

The serotonergic system is strongly implicated in anxiety and depression, and it is the first-line target for pharmacological treatment (Tamblyn et al., 2019). Genetic factors predispose people to these disorders, but the anatomic and molecular diversity of the main serotonergic nuclei, the dorsal raphe (DR) and median raphe (MR), and their extensive projections make establishing causal links among genes, physiology, and etiology complex. Nevertheless, genome-wide screening can highlight correlations with genetic or epigenetic markers (Li et al., 2016), whose effects can then be tested through gene knockout in animal models.

A genome-wide screening by Baselmans et al. (2015) found that *Zfp1* methylation was negatively correlated with factors associated with well being. ZFPM1 is a zinc-finger cofactor that interacts with the GATA family of transcription factors and is expressed in certain serotonergic and GABAergic neurons from early development to adulthood. Therefore, Tikker et al. (2020) recently used selective gene knockout to examine the role of ZFPM1 in development and behavior.

Tikker et al. (2020) generated *Zfp1* conditional knockout (*Zfp1^{CKO}*) mice in which *Zfp1* was knocked out selectively

in rhombomere 1, an embryonic hind-brain segment in which serotonin neurons are derived. The number of serotonergic neurons was reduced in the ventrolateral DR (DRVL) and increased in the dorsal DR (DRD) in *Zfp1^{CKO}* mice. Notably, a substantial number of DRD serotonergic neurons expressed the DRVL-specific marker BCL11B, and there was a decrease in the number of serotonergic cells expressing this marker in DRVL. No change was detected in ventral DR (DRV) or B8 MR areas, and, importantly, the total number of neurons across DR nuclei was unaffected. Based on these findings, the authors hypothesized that *Zfp1* knockout disrupted cell positioning, rather than generation. The authors speculate that *Zfp1* regulates positioning by repressing the transcriptional processes that cause medial migration of serotonergic precursors, and thus knocking out *Zfp1* causes DRVL neurons to be displaced to DRD.

Notable effects of *Zfp1* knockout were also produced at DR projection sites. The density of serotonergic axons was greatly reduced in the dorsal hippocampus (dHPC), ventral hippocampus (vHPC), dorsal lateral geniculate nucleus, and superior colliculus. In contrast, the density increased in the basolateral amygdala (BLA), although less consistently than the reductions seen elsewhere.

Knockout of *Zfp1* also affected mouse behavior. In an elevated plus maze, *Zfp1^{CKO}* mice made less open arm entries than control littermates, indicating anxiety-like behavior. In a recognition task,

Zfp1^{CKO} mice spent less time than controls exploring novel objects. *Zfp1^{CKO}* mice also showed more freezing during contextual fear conditioning compared with controls, and this effect was abolished by treatment with the selective serotonin reuptake inhibitor (SSRI) fluoxetine. Tikker et al. (2020) tested for depression-like behaviors with a forced swim test but found no significant differences between control and *Zfp1^{CKO}* mice, and both groups responded equally positively to fluoxetine in this test. Overall, these results suggest that alterations in specific projections led to enhanced fear and anxiety-like responses in *Zfp1^{CKO}* mice, while behaviors dependent on spared serotonergic projections were unaffected.

The two key findings of Tikker et al. (2020) were that *Zfp1* is critically involved in normal development of the DR serotonergic system and that alterations in *Zfp1* produce anxiety-like and fear-related behaviors. The article also raised two important questions. First, how do the serotonergic axonal density changes produced by *Zfp1* knockout align with our current understanding of DRVL and DRD projections? Second, how does each of the affected areas contribute to the behavioral phenotype seen?

The reduction in hippocampal serotonergic afferents in *Zfp1^{CKO}* mice is surprising if attributed solely to the displacement of DRVL to DRD. In a previous tracing study, Muzerelle et al. (2016) explicitly noted a lack of projections from the DRVL to the hippocampus. This suggests that *Zfp1* knockout

Received Dec. 13, 2020; revised Mar. 15, 2021; accepted Mar. 18, 2021.

The authors declare no competing financial interests. Correspondence should be addressed to Masakazu Taira at masakazu.taira@oist.jp or Miles J. Desforges at miles.desforges@oist.jp. <https://doi.org/10.1523/JNEUROSCI.3125-20.2021>

Copyright © 2021 the authors

either affected a hippocampal-projecting area not examined by Tikker et al. (2020) or altered axonal targeting of neurons in the examined DR regions without affecting cell placement. Serotonergic projections to the hippocampus include dense efferents from the caudal DR (DRC) (B6), MR (B8), and some sparse efferents from DRV and DRD (Soiza-Reilly and Gaspar, 2020). Of these, only the DRC was not specifically examined by Tikker et al. (2020). It may be that *Zfp1* knockout affected the DRC, which is adjacent to the DRD, but this was missed because these neurons were included with other regions during analysis. Substantiating this possibility, it was noted that the DRVL region in which significant change was found was situated caudally. Alternatively, Tikker et al. (2020) may have limited their analysis to B7 cells, which includes the DRD, DRVL, and DRV but excludes DRC, and so did not note changes in the DRC. Complicating the DRC hypothesis, Kast et al. (2017) found that the majority of DRC serotonergic neurons project strongly to the vHPC but not the dHPC, whereas Tikker et al. (2020) found both hippocampal regions were equally affected. Still another possibility is that *Zfp1* knockout altered axonal targeting in the hippocampal projecting MR without affecting cell location. A further experiment using retrograde tracers in the hippocampus of *Zfp1*^{CKO} mice might shed light on the curious decrease in hippocampal serotonergic axon density.

The enhanced anxiety-related behavior and contextual fear conditioning reported by Tikker et al. (2020) are consistent with the known functions of the regions affected by *Zfp1* knockout. In particular, the DRVL is strongly linked to anxiety-like behaviors. The intrinsic excitability properties of DRVL neurons appear to mediate greater activation by stressful stimuli (Crawford et al., 2010). For example, one previous study showed that an elevated T-maze task increased *c-fos* expression specifically in the DRVL (Vilela-Costa et al., 2019). That same study also showed that chronic, but not acute, fluoxetine treatment increased time spent in the open arm, decreased *c-fos* expression in nonserotonergic cells, and increased *c-fos* expression in serotonergic cells. 5-HT_{1A} heteroreceptors on nonserotonergic cells were speculated to be a mechanism for this shift. The effect of fluoxetine suggests DRVL serotonergic output is anxiolytic, while DRVL nonserotonergic activity may be anxiogenic. Consequently, anxiety-like behavior may arise in *Zfp1*^{CKO} mice through both decreased serotonergic output in DRVL

projection sites and increased nonserotonergic activity in the DRVL.

Reduction in the number of serotonergic neurons in DRVL was mirrored by an increase in DRD. This increase may also be instrumental in the anxiety-like behavioral phenotype. Ren et al. (2018) found that activation of DR projections to the central amygdala (CeA) promoted anxiety-like behaviors. In addition, activity of serotonergic neurons in the CeA strongly increased after footshock (Ren et al., 2018). And inhibition of CeA neurons that express 5-HT_{2A} receptors downregulates learned fear responses (Isosaka et al., 2015). Accordingly, increased DRD serotonin output in the CeA in response to footshock may contribute to the enhanced contextual fear conditioning.

As the hippocampus is associated with contextual fear learning and anxiety-like behaviors, reduced serotonergic afferents could play a key role. In contextual fear conditioning, hippocampal neurons have been proposed to represent the contextual cues that come to elicit a conditioned response by exciting the BLA, which in turn activates the CeA to produce fear responses (Maren et al., 2013). Waider et al. (2019) showed that knockout of *Tph2*, which is necessary for serotonin production, increased activity in the dHPC along with fear responses. Conversely, Mlinar and Corradetti (2018) found that application of serotonin in the dHPC decreased activity. Therefore, *Zfp1* knockout may enhance fear responses by reducing serotonergic inhibition in the dHPC, thus enhancing activation of BLA. Notably, *Zfp1* knockout also slightly increased the number of serotonergic axons in the BLA itself. Furthermore, the BLA has been linked to neophobia (Sarowar et al., 2017), another behavioral phenotype reported by Tikker et al. (2020).

Intriguingly, chronic serotonin depletion in the hippocampus, as presumably occurred in *Zfp1*^{CKO} mice, appears to paradoxically produce anxiogenic effects similar to those of acute serotonin activation. For example, Ohmura et al. (2020) found that optogenetic activation of serotonergic terminals in vHPC was anxiogenic. This seeming contradiction may mirror the effect of SSRIs in acutely increasing synaptic serotonin, worsening anxiety and depression, but chronically rebalancing release and uptake to alleviate symptoms. One long-term effect of serotonin in the hippocampus that may contribute to improving symptoms is the upregulation of neurogenesis. Mahar et al. (2014) describe how chronic unpredictable

stress decreases both DR output and hippocampal neurogenesis, producing depression-like behavior, all of which are alleviated by chronic SSRIs. Tikker et al. (2020) performed a test for depression, the forced swim test, and they found fluoxetine-induced immobility reduction was unaffected by *Zfp1*^{CKO} mice. However, Mahar et al. (2014) discuss how fluoxetine decreases immobility independently of hippocampal neurogenesis. Therefore, the reduction in serotonergic afferents to the hippocampus in *Zfp1*^{CKO} mice may have reduced neurogenesis without influencing the forced swim test. Further investigation of depression-like behavior and hippocampal neurogenesis may be fruitful in *Zfp1*^{CKO} mice.

As a final point of interest, the role of *Zfp1* in hippocampal function may not be limited to the developmental effects discussed here. Li et al. (2016) identified *Zfp1* as a novel gene that is methylated in the hippocampus following acute stressful restraint. This suggests a dynamic role for *Zfp1* in the adult brain, which warrants future examination.

In conclusion, Tikker et al. (2020) found that *Zfp1* is required for the normal development of DRVL serotonergic neurons that project to brain regions involved in anxiety and fear learning. The behavioral phenotype found in *Zfp1*^{CKO} mice suggests that *Zfp1* may be a useful genetic or epigenetic etiologic marker for investigating anxiety-related disorders.

References

- Baselmans BM, van Dongen J, Nivard MG, Lin BD, Consortium B, Zilhao NR, Boomsma DI, Bartels M (2015) Epigenome-wide association study of wellbeing. *Twin Res Hum Genet* 18:710–719.
- Crawford LK, Craige CP, Beck SG (2010) Increased intrinsic excitability of lateral wing serotonin neurons of the dorsal raphe: a mechanism for selective activation in stress circuits. *J Neurophysiol* 103:2652–2663.
- Isosaka T, Matsuo T, Yamaguchi T, Funabiki K, Nakanishi S, Kobayakawa R, Kobayakawa K (2015) Htr2a-expressing cells in the central amygdala control the hierarchy between innate and learned fear. *Cell* 163:1153–1164.
- Kast RJ, Wu HH, Williams P, Gaspar P, Levitt P (2017) Specific connectivity and unique molecular identity of MET receptor tyrosine kinase expressing serotonergic neurons in the caudal dorsal raphe nuclei. *ACS Chem Neurosci* 8:1053–1064.
- Li S, Papale LA, Zhang Q, Madrid A, Chen L, Chopra P, Keleş S, Jin P, Alisch RS (2016) Genome-wide alterations in hippocampal 5-hydroxymethylcytosine links plasticity genes to acute stress. *Neurobiol Dis* 86:99–108.
- Mahar I, Bambico FR, Mechawar N, Nobrega JN (2014) Stress, serotonin, and hippocampal

- neurogenesis in relation to depression and antidepressant effects. *Neurosci Biobehav Rev* 38:173–192.
- Maren S, Phan KL, Liberzon I (2013) The contextual brain: implications for fear conditioning, extinction and psychopathology. *Nat Rev Neurosci* 14:417–428.
- Mlinar B, Corradetti R (2018) Differential modulation of CA1 impulse flow by endogenous serotonin along the hippocampal longitudinal axis. *Hippocampus* 28:217–225.
- Muzerelle A, Scotto-Lomassese S, Bernard JF, Soiza-Reilly M, Gaspar P (2016) Conditional anterograde tracing reveals distinct targeting of individual serotonin cell groups (B5-B9) to the forebrain and brainstem. *Brain Struct Funct* 221:535–561.
- Ohmura Y, Tsutsui-Kimura I, Sasamori H, Nebuka M, Nishitani N, Tanaka KF, Yamanaka A, Yoshioka M (2020) Different roles of distinct serotonergic pathways in anxiety-like behavior, antidepressant-like, and anti-impulsive effects. *Neuropharmacology* 167:107703.
- Ren J, Friedmann D, Xiong J, Liu CD, Ferguson BR, Weerakkody T, DeLoach KE, Ran C, Pun A, Sun Y, Weissbourd B, Neve RL, Huguenard J, Horowitz MA, Luo L (2018) Anatomically defined and functionally distinct dorsal raphe serotonin sub-systems. *Cell* 175:472–487.e20.
- Sarowar T, Grabrucker S, Boeckers TM, Grabrucker AM (2017) Object phobia and altered RhoA signaling in amygdala of mice lacking RICH2. *Front Mol Neurosci* 10:180.
- Soiza-Reilly M, Gaspar P (2020) From B1 to B9: a guide through hindbrain serotonin neurons with additional views from multidimensional characterization. In: *Handbook of behavioral neuroscience* (Müller CP, Cunningham KA, eds), pp 23–40. Amsterdam: Elsevier.
- Tamblyn R, Bates DW, Buckeridge DL, Dixon W, Forster AJ, Girard N, Haas J, Habib B, Kurteva S, Li J, Sheppard T (2019) Multinational comparison of new antidepressant use in older adults: a cohort study. *BMJ Open* 9:e027663.
- Tikka L, Casarotto P, Singh P, Biojone C, Piepponen TP, Estartús N, Seelbach A, Sridharan R, Laukkanen L, Castrén E, Partanen J (2020) Inactivation of the GATA cofactor ZFPM1 results in abnormal development of dorsal raphe serotonergic neuron subtypes and increased anxiety-like behavior. *J Neurosci* 40:8669–8682.
- Vilela-Costa HH, Spiacci A Jr, Bissolli IG, Zangrossi H Jr (2019) A shift in the activation of serotonergic and non-serotonergic neurons in the dorsal raphe lateral wings subnucleus underlies the panicolytic-like effect of fluoxetine in rats. *Mol Neurobiol* 56:6487–6500.
- Waider J, Popp S, Mlinar B, Montalbano A, Bonfiglio F, Aboagye B, Thuy E, Kern R, Thiel C, Araragi N, Svirin E, Schmitt-Böhrer AG, Corradetti R, Lowry CA, Lesch KP (2019) Serotonin deficiency increases context-dependent fear learning through modulation of hippocampal activity. *Front Neurosci* 13:245.