This Week in The Journal

Sox9 Reserves Progenitors for Making Upper-Layer Neurons

Jaime Fabra-Beser, Jessica Alves Medeiros de Araujo, Diego Marques-Coelho, Loyal A. Goff, Marcos R. Costa, et al.

(see pages 6969–6986)

Cortical excitatory neurons are generated by radial glial progenitor cells (RGCs) that divide in the ventricular zone; in mice they are generated between embryonic day 11 (E11) and E17. Newborn neurons migrate away from the ventricles to form the cortical layers sequentially, starting with layer VI. Because neurons in different cortical layers are morphologically and functionally distinct, RGCs must produce different types of neurons over time; how they achieve this remains incompletely understood. In particular, the extent to which RGCs programmed to generate upper-layer neurons are present during earlier stages of corticogenesis has been a matter of controversy. But Fabra-Beser et al. provide evidence that RGCs destined to make upper-layer neurons are present in the ventricular zone as early as E12.5.

Using *in utero* electroporation to introduce fluorescent proteins under the control of different promoters, Fabra-Beser et al. found that many RGCs labeled on E12.5 generated progeny that populated the ventricular, subventricular, and intermediate zones of the developing cortex 24 h later; the neural progeny eventually populated the lower cortical layers. In contrast, RGCs in which the *Hes5* promoter was active on E12.5 produced few progeny that exited the ventricular layer by the next day; the progeny of these cells ultimately populated upper cortical layers.

Single-cell RNA sequencing analysis suggested that the majority of RGCs electroporated on E12.5 generated postmitotic neurons within 16 h, but those in which the *Hes5* promoter was active remained as RGCs. In addition, the analysis identified 11 genes that were expressed at higher levels in *Hes5*⁺ cells than in *Hes5*[−] cells. One of these genes, *Sox9*, encodes a transcription factor previously shown to be involved in stem cell maintenance and suppression of

differentiation. Importantly, overexpressing *Sox9* in E12.5 RGCs reduced the number of actively dividing cells, increased the number of these cells remaining in the ventricular zone the next day, and ultimately increased the number of upper-layer neurons produced. Knocking down *Sox9* had the opposite effects.

These results suggest that high levels of *Sox9* expression maintain RGCs in a quiescent state at E12.5, likely holding them in reserve to generate upper-layer cortical neurons in subsequent days.

Most RGCs electroporated on E12.5 generate lower-layer (LL) neurons (left), but overexpression of *Sox9* in RGCs promotes generation of upper-layer (UL) neurons (right). See Fabra-Beser et al. for details.

Carboxypeptidase E Promotes Membrane Insertion of TrkB

Na Li (李娜), Shuai-Wen Teng (滕帅文), Ling Zhao (赵玲), Jing-Rui Li (李敬瑞), Jia-Ling Xu (许佳玲), et al.

(see pages 6987–7002)

Brain-derived neurotrophic factor (BDNF) acts throughout the nervous system to promote generation, maturation, synaptic plasticity, and survival of neurons. Although most neurons can respond to BDNF, its actions are circumscribed by regulated expression of its receptor, TrkB. TrkB is typically sequestered within neurons, but the receptors are rapidly inserted into the plasma membrane in response to specific triggers, including neural activity. Activity-induced insertion of TrkB depends on calcium-dependent activation of adenylyl cyclase, which increases cAMP levels, leading

to activation of protein kinase A (PKA) and phosphoinositide 3-kinase (PI3K; Andreska et al., 2020, *Cell Tissue Res* 382:5). These kinases may promote membrane insertion of TrkB in part by acting on specific intracellular trafficking proteins, such as carboxypeptidase E (CPE).

CPE was originally identified as an enzyme involved in converting hormones from the immature pro-form to mature forms, but it is also involved in sorting and trafficking of secretory vesicles. Li, Teng, Zhao, et al. found that CPE and TrkB interact in transfected HEK293 cells and colocalize in puncta—including transport vesicles—throughout cultured hippocampal neurons. Importantly, chemically induced long-term potentiation (cLTP) significantly increased colocalization of CPE and TrkB near the plasma membrane, and knocking down CPE attenuated the increase in surface TrkB levels that normally occurred after cLTP.

Mutation analysis identified a 5 aa sequence in the C terminus of CPE that interacts with a segment (called Box2) of the TrkB juxtamembrane domain. Deleting Box2 prevented cLTP-induced membrane insertion of TrkB, whereas adding Box2 to a TrkB isoform that normally is unresponsive to cLTP conferred such responsiveness. Furthermore, an interfering peptide targeting the TrkB-binding domain of CPE (TAT-CPE^{452–466}) reduced cLTP-induced increases in TrkB surface expression, spine density, and spine width. TAT-CPE^{452–466} also reduced BDNF-induced activation of TrkB and downstream kinases. Finally, infusing TAT-CPE^{452–466} into mouse hippocampus *in vivo* impaired acquisition and consolidation of conditioned fear memory and prevented conditioning-induced increases in CPE–TrkB interaction and TrkB phosphorylation.

These results indicate that CPE interacts with TrkB within cells and promotes its insertion into the plasma membrane during cLTP. This effect might be mediated in part by activity-induced increases in CPE–TrkB binding. Future work should investigate this and other potential mechanisms by which activity promotes CPE-mediated TrkB insertion.

This Week in The Journal was written by Teresa Esch, Ph.D.
<https://doi.org/10.1523/JNEUROSCI.41.33.2021>