This Week in The Journal

Cannabinoid Receptors and Reconsolidation of Cocaine Memories

Jessica A. Higginbotham, Rong Wang, Ben D. Richardson, Hiroko Shiina, Shi Min Tan, et al.

(see pages 613–629)

A major challenge for people fighting substance abuse disorders is resisting the urge to seek and use drugs when exposed to drug-associated objects and contexts. Such urges might be lessened by weakening learned associations between cues and drug use. One way to do this is by interfering with memory reconsolidation, a process that occurs when learned stimuli are re-encountered. Studies in animals have shown that reactivation of learned associations during such encounters results in transient destabilization of the memory. The memory is then reconsolidated via cellular and molecular processes that include gene transcription and protein synthesis. Interfering with these processes can weaken or even erase the link between stimulus and response.

Higginbotham et al. report that associations between specific contexts and cocaine self-administration can be disrupted by systemic administration of the cannabinoid type 1 receptor (CB1R) antagonist AM251. Rats were trained to press a lever to receive cocaine in a specific context, then were placed in a second context where lever-pressing had no effect. Under this regimen, rats stopped pressing levers in the unpaired chamber, but resumed lever-pressing when returned to the original context, demonstrating that they associated the original context with cocaine delivery. As expected, reactivation of this learned association during a 15 min exposure to the cocaine-paired context resulted in memory destabilization and reconsolidation, as indicated by increased expression of the immediate early gene *zif268*, phosphorylation of the NMDA-receptor subunit GluN2B, and phosphorylation the AMPA-receptor subunit GluA1 in the basolateral amygdala. Notably, these changes were blocked by administering AM251 immediately after re-exposure to the cocaine-paired context. Moreover, AM251 administration immediately after

the brief exposure led to reduced lever-pressing and reduced expression of *zif268*, GluN2B, and GluA1 when rats were returned to the cocaine-paired context 3 d later.

These results suggest that cannabinoid signaling contributes to reconsolidation of learned associations that promote cocaine seeking and that blocking this signaling reduces subsequent cocaine seeking in response to contextual cues. Blocking CB1Rs was shown to disrupt molecular changes occurring in the basolateral amygdala during reconsolidation; future studies will need to determine the relative contributions of these disruptions and effects in other brain regions to subsequent behavioral changes.

When gerbils were exposed to background noise (during time indicated by black bars under plots), MNTB neurons exhibited time-varying responses (black traces). Under normal conditions (top), a pure tone (presented during red bars) modulated MNTB responses, with louder tones (purple) producing greater modulation than quieter tones (orange). Blocking nAChRs (bottom) reduced this modulation, and thus reduced the ability of MNTB neurons to discriminate signal from noise. See Zhang et al. for details.

Cholinergic Modulation of Auditory Neurons in MNTB

Chao Zhang, Nichole L. Beebe, Brett R. Schofield, Michael Pecka, and R. Michael Burger

(see pages 674–688)

Mammals use minute differences in the intensity and time of arrival of sounds in the two ears to localize sound sources. Interaural differences in intensity and arrival time are computed by neurons in the lateral and medial nuclei of the superior olivary complex in the brainstem. These

neurons receive precisely timed excitatory input from the cochlear nucleus and inhibitory input from the medial nucleus of the trapezoid body (MNTB), another nucleus of the superior olivary complex. MNTB neurons receive strong synaptic input from the contralateral cochlear nucleus, which drives them to spike at high frequency. The presence of nicotinic acetylcholine receptors (nAChRs) throughout the superior olivary complex suggests that cholinergic signaling modulates auditory processing in these nuclei. How acetylcholine affects MNTB neuron activity has been unknown, however. Zhang et al. addressed this question by applying nAChR antagonists while recording sound-evoked responses in gerbil MNTB neurons *in vivo*.

MNTB neurons exhibit spontaneous firing in the absence of sound. For sounds above threshold intensity, the spike rate of MNTB neurons increases linearly with sound level until reaching a maximum rate, where the response saturates. Antagonists of either $\alpha 7$ or $\alpha 4\beta 2$ nAChRs reduced the responses of MNTB neurons to above-threshold sounds without affecting spontaneous activity. Furthermore, the antagonists reduced the slope of the rate-level function, and thus reduced the ability of MNTB neurons to represent differences in sound intensity. Consequently, blocking nAChRs reduced the ability of MNTB neurons to detect pure tone stimuli delivered in the presence of noise. Notably, applying the $\alpha 7$ -specific antagonist after the $\alpha 4\beta 2$ antagonist led to a further decrease in tone-evoked spiking, suggesting individual neurons expressed both receptors and the receptors activated distinct signaling pathways. Consistent with the latter hypothesis, only the $\alpha 7$ -specific antagonist increased spike latency.

These results demonstrate that cholinergic signaling helps MNTB neurons represent small differences in sound level. Therefore, acetylcholine is likely critical for accurate computation of interaural level differences used for sound localization. Future studies should investigate the molecular mechanisms by which $\alpha 7$ and $\alpha 4\beta 2$ nAChRs modulate MNTB neuron responses and determine whether cholinergic input to the MNTB varies with behavioral state.

This Week in The Journal was written by Teresa Esch, Ph.D.
<https://doi.org/10.1523/JNEUROSCI.41.4.2021>