This Week in The Journal

How to Rock a Fly to Sleep

Shahnaz Rahman Lone, Sheetal Potdar, Archana Venkataraman, Nisha Sharma, Rutvij Kulkarni, et al.

(see pages 9403–9418)

Sleep timing is influenced by many factors, including circadian rhythms, environmental conditions, and homeostatic sleep drive, which increases over time awake. The ability of gentle rocking to induce sleep in infants has been known for millennia, and rocking promotes sleep in adults as well. Remarkably, this phenomenon is not unique to humans: gentle rhythmic motion can also induce sleep in mice and even in fruit flies. Lone et al. report that *Drosophila* kept on an orbital shaker sleep more than control flies, and this effect depends on a particular type of sensory neuron.

Flies exposed to continuous orbital motion spent more time sleeping—defined as ≥ 5 min of continuous inactivity—during the light period than control flies housed nearby. In addition, more intense stimulation was needed to arouse motion-treated flies, and these flies returned to sleep more quickly than controls. Importantly, orbital motion did not affect activity levels when flies were awake, and sleep levels returned to normal the day after motion was stopped. Moreover, if orbital motion was presented only during the first 6 h of a 12 h light period, flies slept more than controls during the motion period, but slept less than controls for the remainder of the light period, suggesting sleep drive was reduced.

Orbital motion failed to induce sleep in flies lacking antennae, likely because of the loss of antennal mechanosensory (Johnston's) organs. Consistent with this, ablating neurons that express the mechanosensory receptor *Nanchung* significantly reduced motion-induced sleep, whereas activating these neurons promoted sleep. Surprisingly, although silencing *Nanchung*-expressing neurons reduced motion-induced sleep, flies exposed to morning orbital motion still slept less in the afternoon. Similarly, silencing one subset of projections from Johnston's organs to the antennal mechanosensory and motor center (AMMC) in the brain reduced motion-induced sleep in the morning without

affecting afternoon wakefulness, whereas silencing a second subset increased afternoon wakefulness without decreasing sleep in the morning.

These data suggest that orbital motion is detected by mechanosensory neurons in Johnston's organs that innervate the AMMC to regulate sleep onset and sleep drive. Previous work has shown that AMMC neurons project indirectly to sleep-control centers of the brain, which are likely responsible for inducing sleep in response to orbital motion.

Markers of endocytic compartments in mouse cortical neurons, pseudocolored according to movement rate. Overexpression of tSNARE1 slows movement into late endosomes. See Plooster et al. for details.

Linking Endosomal Transport to Schizophrenia Risk

Melissa Plooster, Guendalina Rossi, Martilias S. Farrell, Jessica C. McAfee, Jessica L. Bell, et al.

(see pages 9466–9481)

Schizophrenia is a complex disease with unclear etiology. Genetics contribute greatly to disease risk, but in most cases, the risk-conferring loci are in noncoding regions of DNA and have small, difficult-to-discern effects on brain function. One such locus is in an intronic region of *TSNARE1*. A transcriptome-wide association study linked overexpression of *TSNARE1* to schizophrenia risk, strengthening the case for this gene's involvement in the disease. Although

TSNARE1 contains a conserved DNA-binding domain and a SNARE domain, suggesting a role in fusion of lipid membranes, until now its expression and functions in the brain have been unknown.

By consulting human gene-expression databases, Plooster et al. found that *TSNARE1* is highly expressed in cortex, predominantly in excitatory neurons and endothelial cells. Three splice variants of *TSNARE1* were expressed, all of which included a Qa-type SNARE domain. This domain had greatest homology with the SNARE domain of syntaxin 12 (Stx12), a protein involved in trafficking vesicles between endosomes. Notably, tSNARE1 competed with Stx12 for incorporation into endosomal SNARE complexes. Importantly, however, $\sim 95\%$ of *TSNARE1* transcripts lacked a sequence encoding a transmembrane or membrane-interaction domain—a domain required for initiating membrane fusion.

Rodents do not express *TSNARE1*, but when GFP-tagged tSNARE1 was expressed in rodent cortical neurons, it localized to puncta in the cytoplasm and nucleus. It was enriched in the soma, but also present in axons and dendrites. Whereas tSNARE1 isoforms containing a transmembrane domain colocalized with Stx12 in multiple endosome compartments, isoforms lacking the transmembrane domain were found predominantly in late endosomes, as was most tSNARE1 in human neural progenitor and neuroblastoma cells. Notably, overexpressing tSNARE1 in rodent neurons disrupted trafficking of the dendritic transmembrane protein Nsg1, which normally colocalizes with Stx12 as it moves from the cell surface to lysosomes. Overexpression of tSNARE1 slowed entry of Nsg1 into late endosomes and lysosomes, but did not affect trafficking to recycling endosomes.

These results suggest tSNARE1 is a fusion-incompetent SNARE protein that competes with Stx12 for incorporation into SNARE complexes and thus inhibits membrane transport from early to late endosomes. Notably, another schizophrenia risk gene, SNAP25, also regulates endocytic trafficking, and it associates with tSNARE1. Therefore, disruption of endosomal transport may be an important contributor to schizophrenia risk.

This Week in The Journal was written by Teresa Esch, Ph.D.
<https://doi.org/10.1523/JNEUROSCI.41.45.2021>