This Week in The Journal

Tonic GABA Currents in Human Cortical Interneurons

Martin Field, Istvan P. Lukacs, Emily Hunter, Richard Stacey, Puneet Plaha, et al.

(see pages 9702–9719)

GABA acts on extrasynaptic GABA_A receptors (GABA_ARs) to produce tonic currents in many neurons in the brain. Such currents contribute to slow oscillations during sleep and mediate the effects of some anesthetics. They may also contribute to postpartum depression and some forms of epilepsy. Notably, tonic GABA currents have different effects in different neuron types. They sometimes hyperpolarize resting membrane potential as expected, but they can also produce shunting inhibition in the absence of hyperpolarization or even cause depolarization. A fuller understanding of how tonic GABA currents affect human neurons will be required before developing effective therapeutics that modulate these currents. As a first step, Field et al. investigated tonic GABA currents in human cortical layer 1 (L1) interneurons.

Interneurons are highly diverse, with numerous subtypes defined by morphology, innervation patterns, electrophysiological properties, and protein expression. To determine which subtypes might have tonic GABA currents, Field et al. first interrogated a human transcriptome database, asking whether GABA_AR subunits typically found in extrasynaptic receptors are expressed in subsets of interneurons defined by gene expression. This analysis suggested that tonic GABA currents occur in L1 neurogliaform cells and in rosehip cells—a type of interneuron so far described only in humans.

Next, the authors measured GABA-induced currents in L1 interneurons in brain slices obtained from people undergoing brain surgery. After clustering neurons based on electrophysiological properties, they found that tonic GABA currents tended to be larger in cells with the properties of rosehip cells (irregular spiking and a voltage sag during hyperpolarizing current steps) than in putative neurogliaform cells. In cells classified morphologically, however, rosehip cells did not have significantly larger tonic

GABA currents than other interneuron types. Instead, a group of neurons with axons that descended into L2/3 had significantly larger GABA currents than neurogliaform cells. Linear regression models indicated that both irregular spiking and having the axonal morphology of rosehip or L2/3-innervating cells were strong predictors of higher tonic GABA currents across L1 interneurons.

Despite these general trends, the amplitude of tonic GABA currents was highly variable within subclasses of L1 neurons. This illustrates the vast heterogeneity among cortical interneurons and raises the possibility that tonic GABA currents are plastic. Future work should examine this possibility and determine how these currents affect interneuron output and circuit function.

A human interneuron with soma and dendrites (red) in L1 and axon (black) descending to L2/3. Such cells typically have larger tonic GABA currents than L1 neurogliaform cells. See Field et al. for details.

A Neuron Controlling the Sequence of Courtship Behaviors

Nobuaki K. Tanaka, Takashi Hirao, Hikaru Chida, and Aki Ejima

(see pages 9732–9741)

Many complex behaviors are composed of simpler behaviors that are performed in a

particular sequence. This is true of many courtship behaviors, including those of *Drosophila*. When a male fruit fly sees and smells a female, he orients toward her and taps her with a foreleg. If gustatory receptors on the leg detect particular molecules, the male extends a wing and vibrates it to produce a song. When the female slows to listen to the song, the male licks her genitalia. If appropriate gustatory cues are present, he will copulate with her. Although many molecules, receptors, and neural pathways responsible for producing these courtship behaviors have been identified, what dictates their sequencing has been unclear. Tanaka et al. have begun to address this question by identifying a neuron involved in the transition from tapping to wing extension.

The authors focused on a lateral antennal-lobe-tract projection neuron (IPN4) because the projection patterns of these neurons suggest they can integrate olfactory, gustatory, auditory, and visual information. Intriguingly, Tanaka et al. found that the innervation patterns of these neurons are sexually dimorphic. More importantly, when synaptic output from IPN4s was blocked, males engaged in less tapping during courtship, often eliminating this step and proceeding to the wing extension step without making physical contact or receiving visual cues from the female. Conversely, increasing IPN4 activity increased tapping and delayed progression to the wing extension stage of courtship. Consequently, the number of wing extensions and copulation attempts were reduced in these flies. Finally, application of a courtship-promoting gustatory cue to male forelimbs reduced IPN4 responses to a courtship-promoting olfactory cue.

These results suggest that olfactory cues from female flies activate IPN4 in males, and this neuron suppresses wing extension. This ensures that males first engage in tapping the female, which allows them to detect gustatory cues that inhibit the activity of IPN4. This inhibition allows wing extension to proceed. The arrangement likely reduces the chance that a male will waste energy courting an inappropriate mate.

This Week in The Journal was written by Teresa Esch, Ph.D.
<https://doi.org/10.1523/JNEUROSCI.41.47.2021>