High Salt Intake Recruits Tonic Activation of NR2D Subunit-Containing Extrasynaptic NMDARs in Vasopressin Neurons

Chiranjivi Neupane,^{1,2} Ramesh Sharma,^{1,2} Yoon Hyung Pai,¹ So Yeong Lee,³ Byeong Hwa Jeon,^{1,2} Hyun-Woo Kim,^{1,2} Javier E. Stern,⁴ and Jin Bong Park^{1,2}

¹Department of Physiology, Chungnam National University, Daejeon 35015, Republic of Korea, ²Biomedicine, Brain Research Institute, College of Medicine, Chungnam National University, Daejeon 35015, Republic of Korea, ³Laboratory of Veterinary Pharmacology, College of Veterinary Medicine and Research Institute for Veterinary Science, Seoul National University, Seoul 151-742, Republic of Korea, and ⁴Neuroscience Institute, Georgia State University, Atlanta, Georgia 30302

In addition to producing a classical excitatory postsynaptic current via activation of synaptic NMDA receptors (NMDARs), glutamate in the brain also induces a tonic NMDAR current (I_{NMDA}) via activation of extrasynaptic NMDARs (eNMDARs). However, since Mg^{2+} blocks NMDARs in nondepolarized neurons, the potential contribution of eNMDARs to the overall neuronal excitatory/inhibitory (E/I) balance remains unknown. Here, we demonstrate that chronic (7 d) salt loading (SL) recruited NR2D subunit-containing NMDARs to generate an Mg^{2+} -resistant tonic I_{NMDA} in nondepolarized [V_h (holding potential) -70 mV] vasopressin (VP; but not oxytocin) supraoptic nucleus (SON) neurons in male rodents. Conversely, in euhydrated (EU) and 3 d SL mice, Mg^{2+} -resistant tonic I_{NMDA} was not observed. Pharmacological and genetic intervention of NR2D subunits blocked the Mg^{2+} -resistant tonic I_{NMDA} in VP neurons under SL conditions, while an NR2B antagonist unveiled Mg^{2+} -sensitive tonic I_{NMDA} but not Mg^{2+} -resistant tonic I_{NMDA} . In the EU group VP neurons, an Mg^{2+} -resistant tonic I_{NMDA} was not generated by increased ambient glutamate or treatment with coagonists (e.g., D-serine and glycine). Chronic SL significantly increased NR2D expression but not NR2B expression in the SON relative to the EU group or after 3 d under SL conditions. Finally, Mg^{2+} -resistant tonic I_{NMDA} selectively upregulated neuronal excitability in VP neurons under SL conditions, independent of ionotropic GABAergic input. Our results indicate that the activation of NR2D-containing NMDARs constitutes a novel mechanism that generates an Mg^{2+} -resistant tonic I_{NMDA} in nondepolarized VP neurons, thus causing an E/I balance shift in VP neurons to compensate for the hormonal demands imposed by a chronic osmotic challenge.

Key words: iMK801; NR2D; oxytocin; salt loading; tonic NMDA; vasopressin

Significance Statement

The hypothalamic supraoptic nucleus (SON) consists of two different types of magnocellular neurosecretory cells (MNCs) that synthesize and release the following two peptide hormones: vasopressin (VP), which is necessary for regulation of fluid homeostasis; and oxytocin (OT), which plays a major role in lactation and parturition. NMDA receptors (NMDARs) play important roles in shaping neuronal firing patterns and hormone release from the SON MNCs in response to various physiological challenges. Our results show that prolonged (7 d) salt loading generated a Mg^{2+} -resistant tonic NMDA current mediated by NR2D subunit-containing receptors, which efficiently activated nondepolarized VP (but not OT) neurons. Our findings support the hypothesis that NR2D subunit-containing NMDARs play an important adaptive role in adult brain in response to a sustained osmotic challenge.

Received July 6, 2020; revised Nov. 19, 2020; accepted Nov. 25, 2020.

Author contributions: C.N., R.S., Y.H.P., S.Y.L., B.H.J., H.-W.K., J.E.S., and J.B.P. designed research; C.N., R.S., and Y.H.P. performed research; C.N., R.S., and J.B.P. analyzed data; J.E.S. and J.B.P. wrote the paper.

Y.H. Pai's present address: Hallym University Sacred Heart Hospital, 22, Gwanpyeong-ro 170 beon-gil, Dongan-gu, Anyang-si, Gyeonggi-do 14068, Republic of Korea.

The authors declare no competing financial interests.

This work was supported by the National Research Foundation of Korea (Grant 2015R1D1A1A02059430), the Korean Health Technology R&D project through the Korean Health Industry Development Institute (Grant HI18C0660), and the National Institute of Neurological Disorders and Stroke/National Institutes of Health (Grant R01-NS-094640 (to J.E.S.)).

Correspondence should be addressed to Jin Bong Park at jinbong@cnu.ac.kr.

<https://doi.org/10.1523/JNEUROSCI.1742-20.2020>

Copyright © 2021 the authors

Introduction

NMDA receptors (NMDARs) are a subclass of glutamate-gated ion channels that can mediate the majority of excitatory input on neurons in the brain. At synaptic and extrasynaptic sites, NMDARs are thought to be responsible for maintaining an excitatory/inhibitory balance in two distinct modalities. Synaptic NMDARs mediate the classical transient EPSCs, whereas extrasynaptic NMDARs (eNMDARs) generate a persistent, tonic NMDAR current (I_{NMDA} ; Papouin and Oliet, 2014). NMDARs can be distinguished from other ionotropic glutamate receptors because of characteristics such as high Ca^{2+} permeability and a

strong voltage-dependent channel block caused by external Mg^{2+} (Traynelis et al., 2010). This Mg^{2+} channel block inhibits ion flux through agonist-bound open NMDARs at resting membrane potential (V_m) values, thus requiring the coincident agonists and membrane depolarization to relieve the Mg^{2+} channel block and allow ions to flow through the channel. Consequently, eNMDARs generate tonic I_{NMDA} at low concentrations of extracellular Mg^{2+} and/or with a depolarized membrane potential (Le Meur et al., 2007; Fleming et al., 2011; Hanson et al., 2019). However, to mediate tonic I_{NMDA} with the low levels of extracellular glutamate normally found in the brain, novel strategies are required, targeting eNMDARs to reduce or completely relieve the Mg^{2+} channel block at resting membrane potentials. The mechanisms underlying this phenomenon are yet to be fully understood.

Most NMDARs are tetrameric assemblies of two NR1 subunits and up to four different NR2 (NR2A, NR2B, NR2C, and NR2D) subunits (Paoletti et al., 2013; Wyllie et al., 2013; Hansen et al., 2018). The identity of the four different NR2 subunits determines the properties of the receptor. For example, NR2D-containing NMDARs are located exclusively extrasynaptically (Lozovaya et al., 2004). NR2D subunits confer a high agonist affinity (Erreger et al., 2007), a weak desensitization (Vicini et al., 1998; Wyllie et al., 1998) of NMDARs. Of note, NR2D receptors are less sensitive to Mg^{2+} block (Monyer et al., 1994; Kuner and Schoepfer, 1996; Momiyama et al., 1996; Huang and Gibb, 2014) and less Ca^{2+} permeable (Siegle Retchless et al., 2012) compared with NR2A and NR2B receptors. Therefore, NR2D-containing receptors may be ideal candidates for sensing low ambient glutamate concentrations in the extrasynaptic space and for generating tonic I_{NMDA} with less excitotoxicity in the brain. To date, the contribution of NR2D-containing NMDARs to tonic I_{NMDA} in nondepolarized neurons has not been investigated in the mature brain, partly because expression of the NR2D subunit is developmentally and regionally regulated. The NR2D subunit is widely expressed during late embryonic and early postnatal development; subsequently, it is generally preserved at low levels in a few regions of the adult brain (Watanabe et al., 1992; Monyer et al., 1994; Dunah et al., 1996; Wenzel et al., 1996; Liu and Wong-Riley, 2010). For example, among the four NR2 subunit mRNAs expressed (Al-Ghoul et al., 1997; Decavel and Curras, 1997), NR2B and NR2D are commonly expressed in the supraoptic nucleus (SON; Doherty and Sladek, 2011). However, the functional significance of NR2D expression in the mature brain remains unknown.

The SON consists of two different types of magnocellular neurosecretory cells (MNCs) that contribute to the hypothalamo-neurohypophyseal system (Swanson and Sawchenko, 1983). SON MNCs synthesize and release vasopressin (VP), a peptide hormone necessary for the regulation of fluid homeostasis, and oxytocin (OT), a closely related hormone that stimulates lactation and parturition. Salt intake increases both plasma sodium and osmolality. Therefore, VP neurons respond to increased osmolality by increasing neuronal activity and hormone release from the neurohypophysis during salt loading (SL). Here, we show that prolonged SL recruits NR2D-containing NMDARs to generate Mg^{2+} -resistant tonic I_{NMDA} in nondepolarized VP neurons. This leads to selective upregulation of the neuronal excitability of VP over OT neurons under SL conditions and contributes to fluid homeostasis via hormone secretion.

Materials and Methods

Animal experiments. All animal experimentation was approved by the Chungnam National University Institutional Animal Care and Use Committee (Registration #2025–1–21) and was conducted in accordance with the National Institutes of Health *Guide for the Care and Use of Laboratory Animals*. Male wild-type (WT) or NR2D knock-out (KO) C57BL/6N mice weighing 20–24 g, and Sprague Dawley (SD) rats and double-transgenic Wistar rats weighing 120–180 g, which express both the arginine VP (AVP)-enhanced green fluorescent protein (AVP-eGFP) fusion gene (Ueta et al., 2005) and the OT-monomeric red fluorescent protein 1 (OT-mRFP1) fusion gene (Katoh et al., 2011) were randomly assigned to either the euhydrated (EU) control group or SL group. NR2D KO C57BL/6N mice (Ikeda et al., 1995) were provided by the RIKEN BioResource Research Center (no. RBRC01840) through the National BioResource Project of the Ministry of Education, Culture, Sports, Science and Technology, Japan. All animals had access to food *ad libitum* throughout the experiments. The SL groups had their water replaced with 2% NaCl to drink for 3 or 7 d. At the end of the SL protocol, the animals were anesthetized with avertin (250 mg/kg, i.p.; Millipore Sigma) and decapitated to extract the brain. Blood was collected from each animal after decapitation and was centrifuged for serum collection. Serum osmolality was measured using the freezing point depression method and a micro-osmometer (model 210, Fiske Associates).

Seven days of SL resulted in increased serum osmolality in both WT SD (EU group: 307 ± 0.6 mOsm/kg, $n=6$; vs SL group: 367.8 ± 4.4 mOsm/kg, $n=7$) and transgenic Wistar rats (EU group: 308.1 ± 0.7 mOsm/kg, $n=31$; vs SL group: 368.8 ± 7.0 mOsm/kg, $n=43$). In C57BL/6 WT mice, serum osmolality increased to 349.3 ± 1.3 mOsm/kg ($n=14$) and 382.2 ± 5.5 mOsm/kg ($n=21$) in the 3 and 7 d SL groups, respectively, and to 327.3 ± 0.7 mOsm/kg ($n=17$) in the EU group. Seven days of SL also significantly increased serum osmolality in NR2D KO mice (EU group: 326.9 ± 1.8 mOsm/kg, $n=10$; vs SL group: 383.5 ± 7.6 mOsm/kg, $n=12$).

Electrophysiology and data analysis. Patch-clamp recordings from SON MNCs were obtained from hypothalamic slices (300 μ m) as previously described (Park et al., 2007). Briefly, the rats and mice were anesthetized with avertin and decapitated, and the brains were rapidly isolated. Slices were maintained in, and perfused with, artificial CSF (aCSF) containing 126 mM NaCl, 5 mM KCl, 1.2 mM $MgCl_2$, 26 mM $NaHCO_3$, 1.2 mM NaH_2PO_4 , 10 mM glucose, and 2.4 mM $CaCl_2$ at pH 7.3–7.4 and saturated with 95% O_2 and 5% CO_2 . In a subset of voltage-clamp recordings, low Mg^{2+} aCSF (20 μ M $MgCl_2$) was used to facilitate measurements of NMDA-mediated currents. The internal solution contained 130 mM K-gluconate, 10 mM KCl, 10 mM HEPES, 5 mM Mg^{2+} ATP, 0.9 mM $MgCl_2$, and 10 mM EGTA. Recordings were obtained at 35°C, using a Multiclamp 700B (Molecular Devices). Currents were filtered at 1 kHz and digitized at 10 kHz (Digidata 1440A and pClamp 9 software, Molecular Devices). Input–output (I–O) curves were plotted as previously described (Pandit et al., 2015). Briefly, the number of evoked spikes versus injected current in response to depolarizing step pulses (of 10 pA from 0 to 60 pA, 1.2 s duration) were plotted, and fitted by a Boltzmann function: $y = 1 / \{1 + \exp[(I_h - I_{50})/k]\}$, in which y is the number of spikes at a given injected current [hyperpolarization-activated cation current (I_h)], I_{50} the current at which half of the maximal spikes are evoked, and k is the slope factor. Data were excluded if the series resistance was not stable throughout the entire recording (20% change) or if neuronal input resistance (IR) was <550 M Ω at the beginning of the recording.

The NMDA receptor-mediated tonic current (tonic I_{NMDA}) was defined as changes in the holding current ($I_{holding}$) in the presence of ionotropic GABA receptor antagonist picrotoxin (200 μ M) and was calculated by the difference in $I_{holding}$ measured as the average of a 2 min steady-state baseline segment obtained before and after the application of NMDAR antagonists. I_{NMDA} was recorded and calculated at -70 mV unless otherwise stated. The effects of I_{NMDA} on firing discharge were studied using a current-clamp method. Firing rates (spontaneous or evoked using DC current injection) were recorded in continuous mode and calculated using Mini Analysis Program (Synaptosoft; RRID:SCR_002184) for a

period of ~3 min before and after application of the NMDAR antagonist (Park et al., 2006).

Drugs were added to the aCSF perfusing solution at differing concentrations. The final concentration of dimethylsulfoxide was <0.05%, when used as a vehicle. DL-2-amino-5-phospho-nopentanoic acid, (2S*,3R*)-1-(phenanthren-2-carbonyl) piperazine-2, 3-dicarboxylic acid (PPDA) and DL-threo- β -benzyloxyaspartic acid (TBOA) were all purchased from Tocris Bioscience. All other drugs were purchased from Sigma-Aldrich.

Single-cell RT-PCR. Single-cell RT-PCR was performed following patch-clamp recordings to categorize the VP neurons, as previously described with minor modifications (Lee et al., 2012). The cytoplasm of the neuron that showed Mg^{2+} -sensitive I_{NMDA} was pulled into a patch pipette with negative pressure and under visual control to ensure the nucleus was not contained. The cytoplasm in the pipette was dissipated into a prepared microtube containing 5 μ l of nuclease-free water (Qiagen) and then immediately stored at -70°C . Single-cell cDNA samples were synthesized using an Applied Biosystems High Capacity cDNA Reverse Transcription Kit (Thermo Fisher Scientific) according to the manufacturer instructions. PCR amplification was then performed using single-cell cDNA as a template. PCR mixtures contained 1 μ l of both forward and reverse 10 μ M primers, 10 μ l of 2 $\times$ Taq master mix buffer (GoTaq Green Master Mix, Promega), and 2–4 μ l of cDNA template. The annealing temperature in the thermal cycler was 60°C , and, in total, 50 cycles were performed. Primers targeting VP (5'-CCTCACCTCTGCCTGCTACTT-3' and 5'-GGGGGCGATGGCTCAGTAGAC-3') were synthesized by Bioneer. Final PCR products were visualized by electrophoresis using 1.8% agarose gels stained with ethidium bromide. All PCR products were purified using a PCR purification kit (Qiagen), and the purified products were then sequenced to confirm amplification of the correct sequences. Genomic DNA samples not treated with reverse transcriptase were used as negative controls to assess for contamination of samples.

Western blotting. SON sections were lysed with 1 $\times$ passive lysis buffer (Cell Signaling Technology) and quantified using a Coomassie Protein Assay Kit (Bio-Rad). Approximately 50 μ g of protein was loaded onto a 10% SDS polyacrylamide gel for electrophoresis and then transferred onto nitrocellulose membranes. The blots were then blocked with TBST (0.1% Tween 20 in 1 $\times$ Tris-buffered saline) containing 5% skimmed milk for 1 h at room temperature. The blots were then incubated at 4°C overnight with primary antibodies against the NMDAR NR2B subunit (1:1000; catalog #06–600, Millipore; RRID:AB_310193) and NR2D subunit (1:1000; catalog #AGC-020, Alomone Labs; RRID:AB_10658334) diluted in TBST containing 5% skim milk. Next the blots were incubated with a horseradish peroxidase-conjugated goat anti-rabbit (catalog #7074, Cell Signaling Technology; RRID:AB_2099233) and anti-mouse (catalog #31430, Thermo Fisher Scientific; RRID:AB_228307) secondary antibody (1:1000). A Pierce enhanced chemiluminescence detection kit (Thermo Fisher Scientific) was used to visualize antibody binding, and the intensity of the bands was measured using ImageJ software 1.42q (National Institutes of Health).

Statistical analysis and experimental design. Numerical data are presented as the mean $\pm$ SEM. To assess the differences in tonic I_{NMDA} under SL conditions, a hierarchical testing procedure was performed. In the first step, a Shapiro–Wilk test was used to test the null hypothesis that the data distribution was normal with a significance level of 5%. For data with a normal distribution, the statistical significance of comparisons was assessed using either a two-sample t test or a one-way ANOVA followed by a *post hoc* test (e.g., Bonferroni's test). If the null hypothesis was rejected, nonparametric tests were used with Microcal Origin software (RRID:SCR_002815). For all experiments male mice or rats were used. Tonic I_{NMDA} was categorized into two modalities, either the Mg^{2+} -sensitive tonic I_{NMDA} measured by I_{holding} shift was induced by increasing the extracellular Mg^{2+} concentration from 20 μ M to 1.2 mM at V_h , at -70 mV, or the Mg^{2+} -resistant tonic I_{NMDA} was measured by I_{holding} shift induced by NMDAR antagonists at V_h , at -70 mV using normal aCSF. Electrophysiological recordings were taken from three or more animals per group, and three to four slices were collected per animal. The Mann–Whitney test was used to compare two groups if the

data were not normally distributed. The pharmacological sensitivity of tonic I_{NMDA} to various NMDAR antagonists was investigated using both WT and NR2D KO animals. To compare I_{NMDA} amplitude and protein expression (NR2B and NR2D) in the EU group, the 3 d SL group and the 7 d SL group, a one-way ANOVA followed by a *post hoc* test was used. The effects of Mg^{2+} -resistant tonic I_{NMDA} on neuronal firing were investigated using NR2C/NR2D antagonists and compared using a Wilcoxon signed-rank test. Finally, the effects of Mg^{2+} -resistant tonic I_{NMDA} on the input–output function of VP neurons in the EU and SL groups were compared using a two-way repeated-measures (RM) ANOVA followed by a *post hoc* test.

Results

To assess the functional significance of tonic NMDAR activation in SON MNCs during an osmotic insult, we compared the tonic I_{NMDA} , defined as the shift in I_{holding} after the application of an NMDAR antagonist in the EU and SL animals. The total tonic I_{NMDA} , composed of Mg^{2+} -sensitive and Mg^{2+} -resistant tonic I_{NMDA} , was measured by increasing the extracellular Mg^{2+} concentration ($[Mg^{2+}]_o$) from 20 μ M to 1.2 mM followed by treatment with different NMDAR antagonists.

SL generated an Mg^{2+} -resistant tonic I_{NMDA} in nondepolarized VP neurons

First, we compared tonic I_{NMDA} in SD rats in the EU, 3 d SL, and 7 d SL groups in low $[Mg^{2+}]_o$ aCSF (Fig. 1A,B). An Mg^{2+} -sensitive tonic I_{NMDA} (indicated by an outward shift in I_{holding} after an increase in $[Mg^{2+}]_o$) was observed in all tested neurons, confirming previous findings that tonic persistent activation of NMDARs mediates tonic I_{NMDA} in SON MNCs (Fleming et al., 2011). Mg^{2+} -sensitive tonic I_{NMDA} did not differ among the EU (10.0 ± 0.6 pA; $n = 18$), 3 d SL (9.5 ± 1.6 pA; $n = 11$), and 7 d SL groups (8.5 ± 0.8 pA; $n = 21$; one-way ANOVA, $F_{(2,47)} = 1.75$, $p = 0.184$). Application of AP5 (100 μ M) in the presence of 1.2 mM $[Mg^{2+}]_o$ further shifted the I_{holding} outwardly (a phenomenon defined as an Mg^{2+} -resistant tonic I_{NMDA}) in 12 of 21 SON MNCs isolated from seven rats from the 7 d SL group (Fig. 1A,B). In contrast, AP5 caused minimal changes in I_{holding} in all SON slices obtained from rats in the EU and 3 d SL groups (0.3 ± 0.2 vs 0.2 ± 0.4 pA, respectively). Although 7 d SL generated Mg^{2+} -resistant tonic I_{NMDA} , total tonic I_{NMDA} values were not different among the three groups.

The SON consists of VP and OT neurons (Swanson and Sawchenko, 1983); therefore, our results showing that 7 d of SL generated an Mg^{2+} -resistant I_{NMDA} in a subgroup of SON MNCs may imply that an osmotic insult leads to the recruitment of Mg^{2+} -resistant tonic I_{NMDA} in a cell type-specific manner. To investigate this hypothesis, experiments were repeated in AVP-eGFP and OT-mRFP1 neurons (Fig. 1C–E). Here, we compared the effect of AP5 on I_{holding} at a holding potential (i.e., V_h) of -70 mV in normal aCSF. In the EU group, all tested VP neurons (11 neurons from four rats) and OT neurons (10 neurons from four rats) failed to express an Mg^{2+} -resistant tonic I_{NMDA} (Fig. 1D,E). Conversely, AP5 selectively produced an Mg^{2+} -resistant tonic I_{NMDA} in 11 of 12 VP neurons in the SL groups but this did not occur in OT neurons (four rats in each group; Fig. 1D,E). This result implies that SL recruits an Mg^{2+} -resistant tonic I_{NMDA} in a VP neuron-specific manner. In both the EU and SL groups, application of the ionotropic glutamate receptor antagonist, kynurenic acid (KynA; 5 mM) in addition to AP5 (AP5 + KynA) completely blocked PSCs but failed to cause a further significant change in I_{holding} (Fig. 1C). This finding supports the dominant role of NMDARs in the

generation of tonic glutamate currents under our recording conditions.

In a subset of experiments, we investigate the sensitivity of I_{NMDA} to intracellular MK-801, which has been used to selectively inhibit NMDARs in single cells (Berretta and Jones, 1996). Intracellular MK-801 (1 mM) contained in a recording pipette significantly reduced both total I_{NMDA} (and/or Mg^{2+} -sensitive I_{NMDA}) in the EU MNCs (from 12.2 ± 1.3 to 12.2 ± 1.3 pA) and 7 d SL MNCs (from 20.6 ± 1.7 to 10.0 ± 1.6 pA, $n = 6-7$ in each, $p < 0.01$ in both cases). Combined with the fact that intracellular MK-801 partially but selectively inhibits NMDAR current (Sun et al., 2018), these results suggested that I_{NMDA} reflects the blockade of NMDARs in SON MNCs being recorded in our recording conditions.

Next, we combined *post hoc* single-cell RT-PCR with electrophysiology in WT SD rats. The results are summarized in Figure 1F. An Mg^{2+} -resistant I_{NMDA} was prominent in 10 different identified VP neurons but was absent in 7 of the non-VP neurons, further supporting the conclusion that the SL-induced Mg^{2+} -resistant I_{NMDA} in VP neurons is not related to the genetic modification of AVP-eGFP neurons.

PPDA blocked Mg^{2+} -resistant tonic I_{NMDA} in nondepolarized SL-VP neurons

To investigate the molecular composition of NMDARs generating an Mg^{2+} -resistant I_{NMDA} , we assessed the pharmacological properties of the Mg^{2+} -resistant I_{NMDA} in VP neurons in the EU and SL groups (Fig. 2).

First, we directly compared the $I_{holding}$ shift in SL VP neurons produced by sequential application of an NR2C/NR2D-selective NMDAR antagonist, PPDA (Feng et al., 2004). PPDA (1 μ M) outwardly shifted the $I_{holding}$ of nondepolarized VP neurons from the SL groups in normal aCSF and subsequent addition of 100 μ M AP5 (PPDA + AP5) failed to induce a further shift in $I_{holding}$ (Fig. 2A,B). In contrast, ifenprodil (IFD; 10 μ M), a selective antagonist of the NR2B subunit (Chenard and Menniti, 1999) failed to alter $I_{holding}$ in VP neurons from the SL groups (Fig. 2B).

To investigate whether eNMDARs mediate Mg^{2+} -resistant tonic I_{NMDA} , we examined and compared the effect of memantine, an NMDAR antagonist that preferentially targets extrasynaptic receptors (Xia et al., 2010; Wu and Johnson, 2015) and MK-801, a use-dependent NMDAR open channel antagonist (Huettner and Bean, 1988), on $I_{holding}$ in nondepolarized VP neurons in the SL groups. The addition of memantine (1 μ M) resulted in Mg^{2+} -resistant I_{NMDA} comparable to that of PPDA, and subsequent addition of AP5 (memantine + AP5) did not

Figure 1. Cell type-specific changes in tonic I_{NMDA} in nondepolarized SON MNCs from rats in the euhydrated and salt-loading groups. **A**, Representative traces of neurons from the euhydrated and 7 d salt-loaded (7dSL) groups showing the outward shift in $I_{holding}$ induced by increased $[Mg^{2+}]_o$ (from 20 μ M to 1.2 mM) followed by the addition of AP5 (100 μ M). **B**, Total I_{NMDA} ($I_{holding}$ shift induced by $[Mg^{2+}]_o$ increase and the addition of AP5) are summarized for the EU, 3 d salt-loaded (3dSL) and 7dSL groups (left). Mg^{2+} -resistant I_{NMDA} ($I_{holding}$ shift induced by AP5 in the presence of 1.2 mM $[Mg^{2+}]_o$) values are summarized in the right panel. Note that Mg^{2+} -resistant I_{NMDA} was observed in neurons in the 7dSL group but not in the EU and 3dSL groups. **C**, Representative traces showing Mg^{2+} -resistant I_{NMDA} in AVP-eGFP neurons in one representative rat from the EU (EU-VP) and SL (SL-VP) groups. **D**, Summarized amplitudes of Mg^{2+} -resistant I_{NMDA} in the labeled neurons under the EU and SL conditions ($***p < 0.001$ compared with EU, two-sample t test). Representative images of AVP-eGFP (arrow) and non-VP (arrowhead) neurons from the recording shown in the inset. **E**, Summarized amplitudes of Mg^{2+} -resistant I_{NMDA} in labeled OT neurons under the EU and SL conditions. Representative images of OT-mRFP1 neurons (arrow) and non-OT neurons (arrowhead) from the recording shown in the inset. **F**, Mg^{2+} -resistant I_{NMDA} are summarized in non-VP [VP(-)] and VP neurons [VP(+)] assessed by *post hoc* single-cell RT-PCR. A representative PCR result in two groups is shown in the inset; $***p < 0.001$ compared with EU, two-sample t test.

further alter $I_{holding}$ (Fig. 2A,B). In contrast, MK-801 (40 μ M) failed to produce any effect on the $I_{holding}$ of VP neurons in the SL groups (Fig. 2B).

In summary, the amplitudes of Mg^{2+} -resistant I_{NMDA} produced by memantine, PPDA, and AP5 did not differ significantly in VP neurons in the SL groups (Fig. 2B), implying that Mg^{2+} -resistant I_{NMDA} is predominantly mediated by extrasynaptic NR2C/NR2D-containing receptors. All NMDAR antagonists (PPDA, memantine, MK-801, and ifenprodil) caused minimal changes to $I_{holding}$ in nondepolarized VP neurons in the EU group (data not shown), supporting the conclusion that SL

Figure 2. Pharmacological properties of tonic I_{NMDA} in nondepolarized VP neurons. **A**, Representative traces of VP neurons from rats in the SL group showing the outward shift in $I_{holding}$ induced by PPDA ($1 \mu M$, top) and memantine (Mem; $1 \mu M$, bottom) in normal aCSF with a nondepolarized membrane potential (V_h , -70 mV). Note that additional application of AP5 ($100 \mu M$) did not cause further changes in $I_{holding}$ in the same neurons. The dotted lines indicate the mean $I_{holding}$ under each condition. **B**, Mean changes in $I_{holding}$ induced by the addition of PPDA, ifenprodil (IFD, $10 \mu M$), Mem, MK801 ($40 \mu M$), and AP5 in VP neurons from the SL group, one-way ANOVA ($F_{(4,39)} = 16.4$, $p < 0.001$, with Bonferroni *post hoc* test, $***p < 0.001$, compared with PPDA).

Figure 3. Pharmacological properties of tonic I_{NMDA} in depolarized VP neurons. **A**, Representative traces showing the $I_{holding}$ changes induced by the sequential addition of $10 \mu M$ IFD and $1 \mu M$ PPDA (IFD + PPDA; top and bottom, respectively) in neurons from the EU and SL groups with a depolarized membrane potential (V_h , $+40$ mV). The dotted lines indicate the mean $I_{holding}$ under each condition. **B**, Summarized $I_{holding}$ changes induced by IFD and PPDA in VP neurons from both the EU and SL groups ($n = 9$ /group). $**p = 0.003$ compared with EU, Mann–Whitney test.

recruits additional NR2C/NR2D-containing receptors to mediate Mg^{2+} -resistant I_{NMDA} of VP neurons.

NR2B antagonist ifenprodil blocked tonic I_{NMDA} in depolarized VP neurons

To verify the contribution of NR2B-containing NMDARs to tonic I_{NMDA} in SON MNCs, we investigated the effects of ifenprodil on $I_{holding}$ in VP neurons in the EU and SL groups at a depolarized membrane potential.

In accordance with a previous report (Fleming et al., 2011), ifenprodil (IFD) efficiently produced tonic I_{NMDA} under depolarized conditions in normal aCSF (Fig. 3). At a V_h of $+40$ mV, $10 \mu M$ IFD induced a significant and similar inward $I_{holding}$ shift in VP neurons in both the EU and SL groups, and subsequent addition of $1 \mu M$ PPDA (IFD + PPDA) produced a further inward shift in $I_{holding}$ in VP neurons in the SL groups, but not in the EU group (Fig. 3A,B).

A similar result was observed with sequential application of PPDA and IFD (Fig. 3A), implying that an Mg^{2+} channel block is essential for both inward and outward tonic currents mediated by NR2B-containing NMDARs, whereas NR2C/NR2D-containing NMDARs contributed to the generation of tonic I_{NMDA} in nondepolarized VP neurons in the SL groups.

Increased EPSC frequency with hyperosmotic solution failed to recruit Mg^{2+} -resistant tonic I_{NMDA} in VP neurons

Hypertonic stimuli excite primary osmosensory neurons in the organum vasculosum lamina terminalis (OVLT; Sayer et al., 1984; Honda et al., 1990; Ciura and Bourque, 2006), sending target-selective excitatory inputs to VP neurons (Yang et al., 1994; Stachniak et al., 2014). Glutamate released from presynaptic terminals has the ability to diffuse and activate eNMDARs on neighboring cells (Asztely et al., 1997). Therefore, this enhanced glutamatergic input could recruit Mg^{2+} -resistant tonic I_{NMDA} in VP neurons after hyperosmotic stimulation. To assess this possibility, we examined whether acute hyperosmotic stimulation in brain slices (with the addition of 50 mM mannitol to aCSF) generated an Mg^{2+} -resistant tonic I_{NMDA} in VP neurons.

Direct application of the hyperosmotic solution (50 mM mannitol to aCSF) did not cause any significant change in the $I_{holding}$ of VP neurons in the EU group (Fig. 4A,B), although it significantly increased the frequency of EPSCs (Fig. 4C,D) but with no significant changes in amplitude or decay time kinetics, (Fig. 4E,F), in accordance with a previous report (Yokoyama et al., 2010). Similar results were observed in hypothalamic slices that were incubated in hyperosmotic solution for 20–30 min. Addition of AP5 ($100 \mu M$) did not cause any significant $I_{holding}$ changes in VP neurons in the EU group, even after incubation in hyperosmotic solution. These results disprove the notion that increased ambient glutamate with enhanced glutamatergic input from OVLT generates Mg^{2+} -resistant tonic I_{NMDA} in VP neurons.

Increased glutamate concentration was not sufficient to generate Mg^{2+} -resistant tonic I_{NMDA} in VP neurons

In addition to spillover of glutamate from the synaptic cleft, altered glutamate transporter (GluT) activity could lead to the accumulation of glutamate in the extracellular space, activating eNMDARs and therefore enhancing tonic I_{NMDA} strength (Fleming et al., 2011). To test this hypothesis, we investigated the effects of PPDA on $I_{holding}$ in the presence of $30 \mu M$ glutamate or TBOA in VP neurons in both the EU and SL groups (Fig. 5).

In the presence of TBOA ($100 \mu M$), a broad-spectrum GluT antagonist (Shimamoto et al., 1998), PPDA ($1 \mu M$) caused a significant change in $I_{holding}$ in VP neurons in the SL groups but not in the EU group (Fig. 5A,B); however, subsequent addition of 5 mM KynA (PPDA + KynA) caused a significant outward shift in $I_{holding}$ in all groups. Similar results were observed in the presence of $30 \mu M$ glutamate (Fig. 5C). PPDA failed to produce any significant change in $I_{holding}$ in VP neurons in the EU group while in the presence of glutamate. However, it did produce tonic I_{NMDA} in the SL groups, implying that an increase in ambient glutamate levels is not enough to generate Mg^{2+} -

resistant I_{NMDA} in SON MNCs. The PPDA-induced $I_{holding}$ shift of VP neurons in the SL groups did not differ in the absence (8.5 ± 1.1 pA; Fig. 2) or presence of glutamate or TBOA (one-way ANOVA, $F_{(2,25)} = 0.42$, $p = 0.66$), implying that Mg^{2+} -resistant tonic I_{NMDA} became saturated in VP neurons in the SL groups.

These results support the hypothesis that the Mg^{2+} -resistant I_{NMDA} of VP neurons in the SL groups cannot be attributed simply to increased ambient glutamate levels in the SON.

Increased coagonist concentration was not sufficient to generate Mg^{2+} -resistant tonic I_{NMDA} in VP neurons

NMDARs are unique among glutamate receptors in their requirement for the binding of coagonists, such as glycine or D-serine, in addition to glutamate. In SON MNCs, D-serine is the dominant coagonist at synapses, while glycine is more potent at extrasynaptic receptors (Papouin et al., 2012). We next explored whether the application of exogenous coagonists (glycine or D-serine) generated Mg^{2+} -resistant I_{NMDA} in VP neurons in the EU or SL groups. Glycine ($3 \mu M$) or D-serine ($100 \mu M$) failed to produce any changes in $I_{holding}$ in VP neurons in the EU or SL groups ($p = 0.991$ and 0.493 , $n = 4$ – 5 in all groups, two-sample t test). These results agree with those of previous studies and imply that the glycine binding sites of the eNMDARs mediating tonic I_{NMDA} become saturated in the brain (Le Meur et al., 2007; Papouin et al., 2012).

The NR2C/D subunit-selective potentiator CIQ failed to recruit Mg^{2+} -resistant tonic I_{NMDA} in VP neurons in the EU and SL groups

Next, we used an NR2C/NR2D subunit-selective potentiator, CIQ (Mullasseril et al., 2010), to investigate its effect on Mg^{2+} -resistant tonic I_{NMDA} in VP neurons. CIQ ($20 \mu M$) failed to produce any effect on $I_{holding}$ in VP neurons in the EU and SL groups ($p = 0.687$, $n = 6$ in both groups; two-sample t test), supporting the hypothesis that the SL-induced, NR2D-mediated Mg^{2+} -resistant tonic I_{NMDA} in VP neurons is maximally activated under these particular recording conditions.

NR2D KO prevented SL-induced Mg^{2+} -resistant tonic I_{NMDA} in SON MNCs

To confirm the role of NR2D-containing NMDARs in SL-induced Mg^{2+} -resistant tonic I_{NMDA} , we compared the Mg^{2+} -resistant tonic I_{NMDA} of C57BL/6 WT mice and of NR2D KO mice under both the EU and SL conditions.

First, we compared tonic I_{NMDA} in EU and SL WT mice (Fig. 6A–D) to examine whether SL-induced Mg^{2+} -resistant tonic I_{NMDA} is species specific. The Mg^{2+} -sensitive tonic I_{NMDA} was observed in all tested neurons, and no significant differences were observed in the EU or 3 and 7 d SL groups (Fig. 6A,B). The

Figure 4. Acute hyperosmotic stimulus failed to generate Mg^{2+} -resistant I_{NMDA} in VP neurons. **A**, A representative example showing the lack of change in $I_{holding}$ after a hyperosmotic stimulus containing 50 mM mannitol (360 mOsm/kg) and the addition of AP5 ($100 \mu M$) and KynA ($5 \mu M$). **B**, Mean $I_{holding}$ under the three different conditions, as in **A**, are summarized in the bar graph ($n = 6$). **C**, Expanded current traces showing the effects of a hyperosmotic solution on the degree of ongoing synaptic glutamatergic activity, as in **A**. **D–F**, Mean changes in the spontaneous EPSC frequency (**D**), amplitude (**E**), and decay time kinetics (**F**) induced by a hyperosmotic solution ($n = 6$ /condition). CTL, Control. * $p = 0.03$, compared with control, paired-sample t test.

addition of AP5 ($100 \mu M$) in the presence of normal $[Mg^{2+}]_o$ resulted in an Mg^{2+} -resistant tonic I_{NMDA} in 10 of 20 SON MNCs tested in the 7 d SL group, whereas minimal changes were observed in the SON MNCs of the EU and 3 d SL groups (Fig. 6A,C). Despite these findings, the total I_{NMDA} did not significantly differ among the three groups (Fig. 6D).

Next, we compared Mg^{2+} -resistant tonic I_{NMDA} between WT and NR2D KO mice under EU and 7 d SL conditions (Fig. 6E,F). PPDA ($1 \mu M$) produced an Mg^{2+} -resistant I_{NMDA} in 11 of 20 SON MNCs from seven different mice in the SL group, but no significant changes in $I_{holding}$ were observed in 10 neurons tested in the EU group from four different WT mice (Fig. 6E,F). In contrast, PPDA in combination with AP5 (PPDA + AP5) caused only minimal changes in $I_{holding}$ of all tested SON MNCs from NR2D KO mice in both the EU group (six neurons from three mice) and SL groups (14 neurons from five mice; Fig. 6E,F), implying that NR2D-containing NMDARs play an essential role in mediating SL-induced Mg^{2+} -resistant tonic I_{NMDA} in SON MNCs.

Chronic SL selectively increased NR2D expression over NR2B subunit expression in the SON

To investigate whether SL induces Mg^{2+} -resistant tonic I_{NMDA} via altered expression of extrasynaptic NMDARs in the SON, we

Figure 5. Increased extracellular glutamate failed to generate Mg^{2+} -resistant I_{NMDA} of VP neurons. **A**, Representative examples of the effects of PPDA (1 μM) and KynA (5 mM) on $I_{holding}$ in the presence of 100 μM TBOA, a glutamate transporter antagonist. PPDA produced tonic I_{NMDA} in the SL groups but not in the EU group. **B**, PPDA-induced $I_{holding}$ changes (I_{PPDA}) in the presence of TBOA, as shown in the bar graph ($n = 8$; *** $p < 0.001$, compared with EU, two-sample t test). **C**, I_{PPDA} in the presence of 30 μM glutamate as shown in the bar graph ($n = 8$; *** $p < 0.001$, compared with EU, two-sample t test).

Figure 6. NR2D KO prevented the generation of SL-induced Mg^{2+} -resistant I_{NMDA} . **A**, Representative traces showing that increased $[Mg^{2+}]_o$ (from 20 μM to 1.2 mM) and the addition of AP5 (100 μM) caused an outward shift in $I_{holding}$ in SON MNCs in C57BL/6 mice. AP5 induced an $I_{holding}$ shift in the 7 d SL (7dSL) group but not in the euhydrated or 3 d SL (3dSL) groups. The dotted lines indicate mean $I_{holding}$ level under each condition. **B**, **C**, $I_{holding}$ changes induced by sequential application of Mg^{2+} (Mg^{2+} -sensitive I_{NMDA} ; **B**) and AP5 (Mg^{2+} -resistant I_{NMDA} ; **C**) in each neuron type are plotted. **D**, Summarized total I_{NMDA} in the three groups. **E**, Representative current traces showing the effect of PPDA (1 μM) on $I_{holding}$ in SON MNCs in normal aCSF in the four different groups. Note that SL induced Mg^{2+} -resistant I_{NMDA} in the wild-type mice but not in the NR2D KO mice. **F**, Amplitudes of Mg^{2+} -resistant I_{NMDA} measured by PPDA are shown for the four different groups.

compared the expression of NMDAR subunits NR2B and NR2D in the SON of the EU group with that of mice in the 3 and 7 d SL groups. In all three groups, Western blot analysis showed the presence of NR2B and NR2D in the SON (Fig. 7A). Importantly, while NR2D subunit polypeptide expression was significantly higher in the 7 d SL mice than in the EU and 3 d SL groups (Fig. 7C), NR2B expression was similar in all three groups (Fig. 7B).

Blockage of Mg^{2+} -resistant tonic I_{NMDA} downregulated the neuronal firing of SL-VP neurons

To determine the contribution of Mg^{2+} -resistant tonic I_{NMDA} to resting V_m and IR in SON MNCs, recordings were obtained using the current-clamp mode. We investigated the association between I_{NMDA} attenuation and increased IR in VP neurons in the SL groups using PPDA, which selectively blocks Mg^{2+} -resistant tonic I_{NMDA} (Table 1).

A two-way RM ANOVA revealed that IR was dependent on the addition of 1 μM PPDA ($F_{(3,41)} = 5.89$, $p = 0.04$), but not on SL ($F_{(3,41)} = 2.56$, $p = 0.13$). A *post hoc* test revealed that PPDA significantly increased IR in VP neurons in the SL groups ($p < 0.05$), whereas it caused minimal changes to VP neurons in the EU group ($p > 0.05$; Table 1). Similarly, V_m was dependent on the addition of PPDA ($F_{(3,41)} = 6.45$, $p = 0.02$), but not on SL ($F_{(3,41)} = 0.02$, $p = 0.82$). In accordance with its effect on $I_{holding}$ (Fig. 2), PPDA caused minimal effects on V_m (from -58.2 ± 1.1 to -57.7 ± 1.0 mV, $n = 11$, $p > 0.05$) in VP neurons in the EU group, whereas it significantly hyperpolarized V_m (from -56.5 ± 1.1 to -59.0 ± 1.0 mV, $n = 11$, $p < 0.05$) in VP neurons in the SL groups.

To assess further the functional significance of Mg^{2+} -resistant tonic I_{NMDA} in SL SON MNCs, we examined the effect of PPDA on the spontaneous firing activity of VP and OT neurons (Fig. 8). The spontaneous firing frequency was not dependent on the cell type (two-way ANOVA, $F_{(3,25)} = 0.004$, $p = 0.94$). Furthermore, SL had no significant effect on the spontaneous firing frequency of VP (EU group: 3.2 ± 1.3 Hz, $n = 6$; vs SL group: 5.7 ± 1.5 Hz, $n = 11$) or OT neurons (EU group: 3.4 ± 2.2 Hz, $n = 4$; vs SL group: 5.2 ± 2.1 Hz, $n = 7$; two-way ANOVA, $F_{(3,25)} = 0.93$, $p = 0.34$) or on the proportion of spontaneously active neurons of the VP (6 of 11 neurons in the EU group vs 11 of 13 neurons in the SL group, $p = 0.18$, Fisher's exact test) and OT neurons (4 of 7 neurons in the EU group vs 7 of 9 neurons in the SL groups, $p = 0.60$, Fisher's exact test). All tested neurons that were silent were then subjected to a current injection to produce a firing frequency of ~ 2 Hz to investigate the effect of PPDA on ongoing firing activity (Fig. 8).

In accordance with its effect on Mg^{2+} -resistant tonic I_{NMDA} , PPDA (1 μM) significantly decreased the neuronal firing rate in VP neurons in the SL groups ($n = 13$, $p = 0.0002$, Wilcoxon signed-rank test), while it caused a minimal decrease in VP neurons in the EU group ($n = 11$, $p = 0.311$, Wilcoxon signed-rank test; Fig. 8A,B). PPDA did not cause any significant changes in the neuronal firing of OT neurons in the EU or SL groups (Fig.

Figure 7. SL increased NR2D expression in the SON. **A**, Representative Western blot analysis showing NMDAR NR2B and NR2D subunit expression under EU and SL conditions. **B**, NR2B and **C**, NR2D subunit expression in the SON in the EU and SL groups is summarized and compared, respectively. The protein expression was normalized to the level detected in the EU group and compared with the expression in SL animals. Summarized data shown are as the mean $\pm$ SE ($n = 5$ and $n = 11$ for NR2B and NR2D subunits, respectively). One-way ANOVA, $F_{(2,30)} = 8.48$, $p < 0.05$; with Bonferroni *post hoc* test, $**p < 0.01$.

8A,B). Combined with the results that PPDA increased the input resistance in VP neurons in the SL groups (Table 1), these results further suggested that altered input resistance by the blockade of Mg^{2+} -resistant tonic I_{NMDA} could be sufficient to increase the neuronal firing in the neurons.

To determine whether the decrease in SON neuronal firing after blockade of Mg^{2+} -resistant tonic I_{NMDA} required an intact ionotropic receptor-mediated GABAergic synaptic input, experiments were repeated in the presence of the ionotropic GABA receptor antagonist picrotoxin (200 μM). All recorded neurons were spontaneously active in the presence of picrotoxin. The results are summarized in Fig. 8, C and D. In the presence of picrotoxin, PPDA significantly decreased the neuronal firing rate in

Table 1. Basic membrane properties of VP SON MNCs

		<i>n</i>	IR (m Ω)	<i>V_m</i> (mV)
EU	CTL	11	746.1 $\pm$ 40.2	−58.2 $\pm$ 1.1
	PPDA	11	749.2 $\pm$ 45.2	−57.7 $\pm$ 1.0
SL	CTL	11	659.4 $\pm$ 29.8	−56.5 $\pm$ 1.1
	PPDA	11	705.6 $\pm$ 28.4*	−59.0 $\pm$ 1.0*

CTL, Control.

* $p < 0.05$ compared with respective CTL, two-way RM ANOVA with Bonferroni *post hoc* test.

VP neurons in the SL groups ($n = 8$, $p = 0.02$, paired samples *t* test) but not in the VP neurons in the EU group. In VP neurons in the SL groups, the effect of PPDA on ongoing firing in the presence of picrotoxin ($64.2 \pm 9.2\%$ of control, $n = 8$) was no different from that produced in the absence of picrotoxin ($66 \pm 8.1\%$ of control, $n = 13$, $p = 0.82$, two-sample *t* test).

Together, these results imply that blockade of Mg^{2+} -resistant tonic I_{NMDA} is sufficient to decrease neuronal excitability and firing discharge of VP neurons under SL conditions, even in the absence of an active GABAergic input.

Blockade of Mg^{2+} -resistant tonic I_{NMDA} shifted the input–output function in VP neurons in the SL groups

To assess the functional significance of Mg^{2+} -resistant tonic I_{NMDA} on repetitive firing activity, the effect of PPDA on the I–O function in VP neurons was compared between the EU and SL groups. Depolarizing current steps of increasing amplitude were applied and the numbers of resulting spikes were plotted as a function of the depolarizing steps (Fig. 9). At all levels of current injection, the numbers of action potentials elicited in VP neurons did not differ significantly between the EU and SL groups (Fig. 9A). The I–O function was expressed using the Boltzmann equation with a half-maximal current (I_{50}) of 31.9 ± 1.8 pA and a slope factor (k) of 8.7 ± 0.6 pA ($n = 11$) in VP neurons in the EU group (Fig. 9B), which were no different from the values for VP neurons in the SL groups ($I_{50} = 31.5 \pm 1.6$ pA, $k = 8.4 \pm 0.3$ pA, $n = 11$). A two-way RM ANOVA revealed that the firing responses were dependent on current injection ($F_{(1,140)} = 132.80$, $p < 0.001$) but not on SL ($F_{(1,140)} = 0.05$, $p = 0.94$). PPDA (1 μM) significantly decreased the firing response in VP neurons in the SL groups but not in the EU group (Fig. 9A). As a result, PPDA significantly increased the I_{50} in VP neurons in the SL groups to 38.5 ± 1.6 pA ($F_{(3,41)} = 18.91$, with Bonferroni *post hoc* test, $p < 0.001$), while the changes in the EU group did not show statistical significance (32.8 ± 2.1 pA, $n = 11$, $p > 0.05$).

Discussion

The main findings of this study are that (1) an osmotic challenge mediated by 7 d (but not 3 d) of SL generated an Mg^{2+} -resistant tonic I_{NMDA} in nondepolarized VP neurons; (2) the Mg^{2+} -resistant tonic I_{NMDA} in VP neurons in the SL groups was not a result of increased ambient glutamate concentration; (3) pharmacological blockade and genetic deletion of NR2D subunits inhibited the Mg^{2+} -resistant tonic I_{NMDA} in VP neurons in the SL groups; (4) the Mg^{2+} -resistant tonic I_{NMDA} selectively regulated the firing activity of VP (but not of OT) neurons in the SL groups; and (5) SL increased NR2D expression in the SON. These findings indicate that increased expression and activation of NR2D-containing NMDARs is a novel mechanism generating an Mg^{2+} -resistant tonic I_{NMDA} in VP neurons, which may contribute to elevated neuronal activity and, in turn, VP hormone release to aid fluid homeostasis following the chronic osmotic insult. To the best of our knowledge, this is the first evidence that

Figure 8. Mg^{2+} -resistant tonic I_{NMDA} upregulated neuronal firing in SON MNCs. **A**, Representative traces of neuronal firing in VP and OT neurons from rats in the EU and SL groups. Blockade of Mg^{2+} -resistant tonic I_{NMDA} (PPDA, 1 μM) decreased the firing activity in SL-VP but not in EU-VP, EU-OT, and SL-OT neurons, while subsequent application of 100 μM AP5 (PPDA + AP5) efficiently decreased firing activity in all groups. **B**, Firing frequency under each condition [*** $p < 0.001$, compared with control (CTL) Wilcoxon signed-rank test; $n = 11$, EU-VP; $n = 13$, SL-VP; $n = 7$, EU-OT; $n = 9$, SL-OT]. **C**, Representative traces showing the effect of PPDA on the neuronal firing of EU-VP and SL-VP neurons in the presence of an ionotropic $GABA_A$ receptor antagonist, picrotoxin (PTX; 200 μM). **D**, Firing frequency of VP neurons before and during the application of PPDA in the presence of PTX under the EU ($n = 5$) and SL ($n = 8$) conditions; * $p = 0.02$, compared with control, paired-sample t test.

recruitment of NR2D-containing NMDARs in the adult brain contributes to generation of a unique, functionally relevant extrasynaptic excitatory signaling mechanism. We propose that this novel mechanism plays an important adaptive role in these neuroendocrine cells in response to sustained physiological challenges.

Cellular mechanisms generating Mg^{2+} -resistant tonic I_{NMDA} in VP neurons under SL conditions

SON MNCs represent a conspicuous example of activity-dependent neuronal–glial plasticity (Tweedle and Hatton, 1977, 1984; Theodosis and Poulain, 1993) that involves reversible glial

retraction from and between neighboring neurons leading to increased formation on synaptic inputs on the targeted neurons. Glial retraction with reduced GLT-1 activity increases extracellular glutamate and activates eNMDARs in SON MNCs in response to osmotic challenges (Fleming et al., 2011; Joe et al., 2014). However, in the present study, the increased extracellular glutamate level evoked by exogenous glutamate or GluT blocker was not sufficient to generate or potentiate PPDA-sensitive tonic I_{NMDA} in the EU and SL SON MNCs. These results imply that increased ambient glutamate levels, in the context of an altered topography of the tripartite synapse and/or GluT activity itself, are not enough to generate an Mg^{2+} -resistant I_{NMDA} in VP neurons under SL conditions.

Given that ambient glutamate is sufficient to activate eNMDARs (Sah et al., 1989; Herman and Jahr, 2007; Papouin et al., 2012), altered expression and/or composition of eNMDARs could generate a tonic I_{NMDA} in SON MNCs regardless of ambient glutamate concentrations. Indeed, the lack of change in the abundance of NMDAR subunits previously reported after 3 d of SL (Doherty and Sladek, 2011) and passive dehydration (Carreño et al., 2011) seems to be consistent with our results showing that 3 d of SL was not enough to induce Mg^{2+} -resistant tonic I_{NMDA} in SON MNCs (Fig. 1). Similarly, our results demonstrating that 7 d of SL increased NR2D expression, together with previous reports of NR1 upregulation in chronic SL (7–10 d; Decavel and Curras, 1997), imply that an increased abundance of NMDARs generates Mg^{2+} -resistant tonic I_{NMDA} after 7 d SL in VP neurons. Our results showing that pharmacological and genetic intervention of NR2D prevented Mg^{2+} -resistant tonic I_{NMDA} provide further evidence that NR2D-containing receptors play an essential role in the generation of Mg^{2+} -resistant tonic I_{NMDA} in VP neurons under SL conditions.

Together, our results imply that, in addition to the changes in the neuronal–glial microenvironment, SL recruits NR2D-containing NMDARs, which in turn mediate this unique extrasynaptic signaling, shown in VP neurons.

Diversity of eNMDARs mediating tonic I_{NMDA}

Given that SON MNCs express NR2B and NR2D subunit proteins (Doherty and Sladek, 2011) and that NR1/NR2B/NR2D triheteromers form functional NMDARs in both synaptic (Brothwell et al., 2008) and extrasynaptic membranes (Brickley et al., 2003), it is plausible that, in addition to NR2D

diheteromers, triheteromeric eNMDARs composed of NR1/NR2B/NR2D subunits mediate Mg^{2+} -resistant tonic I_{NMDA} in VP neurons under SL conditions. However, NR1/NR2B/NR2D triheteromers are also sensitive to ifenprodil, although less so than diheteromeric NR1/NR2B receptors (Brothwell et al., 2008), so our results showing that ifenprodil failed to block Mg^{2+} -resistant tonic I_{NMDA} of VP neurons under SL conditions (Fig. 2B) challenge the hypothesis that triheteromeric NR1/NR2B/NR2D receptors or diheteromeric NR2B receptors contribute to the Mg^{2+} -resistant tonic I_{NMDA} of VP neurons under SL conditions. It is also noteworthy that extracellular MK-801 failed to block Mg^{2+} -resistant tonic I_{NMDA} of VP neurons in SL conditions (Fig. 2B). Combined with the fact that extracellular MK-801 preferentially interacts and blocks the NMDAR currents mediated by NR2A- and NR2B-containing NMDARs compared with NR2C- or NR2D-containing receptors in nondepolarized cells (Yamakura et al., 1993; Laurie and Seeburg, 1994; Bresink et al., 1996), insensitivity of Mg^{2+} -resistant I_{NMDA} to the open channel blocker, MK-801 is also in line with the fact that NR2C/NR2D-containing receptors mediate Mg^{2+} -resistant I_{NMDA} of VP neurons.

Although NR2B does not contribute to the Mg^{2+} -resistant tonic I_{NMDA} of VP neurons under SL conditions, it is broadly accepted that NR2B receptors preferentially located in extrasynaptic regions (Tovar and Westbrook, 1999) contribute to tonic I_{NMDA} in the brain. Like previous studies (Fleming et al., 2011), we found that ifenprodil efficiently blocked tonic I_{NMDA} in depolarized SON MNCs (Fig. 3). While still debated, it is generally accepted that eNMDARs are composed of NR2B-containing NMDARs. Since NR2D receptors are less sensitive than NR2B receptors to Mg^{2+} ion channel blockade (Kuner and Schoepfer, 1996), our results imply that both NR2B- and NR2D-containing NMDARs mediate tonic I_{NMDA} , with their relative contributions being dependent on the depolarizing state of neurons (e.g., NR2D is predominant when neurons are hyperpolarized and NR2B and NR2D are predominant when neurons are depolarized).

Cell type-specific generation of NR2D-mediated Mg^{2+} -resistant tonic I_{NMDA} in SON MNCs under SL conditions

SON MNCs release VP and OT hormones into the blood in response to specific physiological demands, which are tightly linked to the electrical activity of neurons (Poulain and Wakerley, 1982; Renaud and Bourque, 1991). The excitatory inputs from the OVLT to SON MNCs are target selective for VP neurons (Yang et al., 1994; Stachniak et al., 2014), implying that they may selectively modulate the excitability of VP neurons in response to a hyperosmotic challenge. However, our results showing that enhanced glutamate input, putatively including those from OVLT, did not generate or potentiate I_{NMDA} in SON MNCs (Figs. 4, 5) argue against this interpretation.

It has been proposed that differential NMDA receptor composition may shape the neurophysiological properties of VP and OT neurons (Al-Ghoul et al., 1997). Indeed, VP neurons

Figure 9. Blockade of Mg^{2+} -resistant tonic I_{NMDA} shifted the I–O function. **A**, Representative examples of action potentials evoked by depolarizing pulses of increasing amplitude before and during the application of PPDA ($1 \mu M$) to VP neurons in the EU and SL groups. Note the decreased numbers of spikes produced in the presence of PPDA by neurons in the SL group at each current amplitude step. V_m was held at -58 mV under all conditions. **B**, Mean numbers of spikes produced as a function of depolarizing current amplitude before (open) and during the application of PPDA (filled) in VP neurons from the EU (circle) and SL (square) groups. Blockade of Mg^{2+} -resistant tonic I_{NMDA} significantly shifted the I–O relationship to the right under SL conditions (see Results) compared with control (CTL; $n = 11$ in each), two-way RM ANOVA.

expressed similar levels of mRNA of the NR2D (31%), NR2B (39%), and NR2C (32%) subunits, whereas the NR2B subunit is predominantly expressed (65%) in OT neurons (Al-Ghoul et al., 1997). These findings, in addition to the results showing that NR2C/NR2D receptors exhibit weaker magnesium block and higher affinity for glycine than NR2B receptors, imply that the high NR2C/NR2D subunit could cause VP neurons to be more susceptible to glutamatergic activation than OT neurons. In this study, pharmacological and genetic intervention of NR2D receptors blocked tonic I_{NMDA} and suppressed the neuronal firing of VP neurons under SL conditions, which is in accordance with previous studies reporting that, at the protein level, NR2C expression is absent in the SON (Doherty and Sladek, 2011) despite the fivefold higher NR2C mRNA in VP compared with OT neurons (Al-Ghoul et al., 1997). Overall, these results imply that during a sustained osmotic challenge, NR2D-mediated tonic I_{NMDA} contributes to the regulation of SON firing activity in a cell type-dependent manner. However, we cannot completely exclude the possibility that SL selectively alters the microenvironment of VP neurons rather than OT neurons, contributing to the cell type-specific nature of this effect.

Functional significance of NR2D-mediated Mg^{2+} -resistant tonic I_{NMDA} in VP neurons under SL conditions

It has long been thought that nonfunctional extrasynaptic receptors may be recruited to active synapses or functionally combined with synaptic signaling, on demand, in the brain. For example, enhanced synaptic activity functionally recruits NR2D-containing eNMDARs to modify excitatory postsynaptic currents in the hippocampus and cerebellum (Clark and Cull-Candy, 2002; Harney et al., 2008). Our results add to the known functional repertoire of eNMDARs, showing that NR2D-containing NMDARs can also be recruited in an “on-demand” basis to extrasynaptic sites to generate a functionally relevant Mg^{2+} -resistant tonic I_{NMDA} in VP neurons.

Although tonic NMDAR activation has been proposed to modulate the input–output function (Sah et al., 1989; Wu et al., 2012), such activation may also lead to cell death, possibly by an excessive Ca^{2+} influx (Hardingham et al., 2002). The permeability of NMDARs to Ca^{2+} ions may confer to NMDARs a central

role in both synaptic plasticity under physiological conditions and neuronal death under excitotoxic pathologic conditions. In addition to other unique biophysical properties of NR2D subunit-containing NMDARs, including weak desensitization, slow deactivation, low Mg^{2+} sensitivity, and high glutamate affinity (Yamakura and Shimoji, 1999; Bai et al., 2013; Paoletti et al., 2013; Wyllie et al., 2013), NR2D receptors show reduced Ca^{2+} permeability (Sieglar Retchless et al., 2012). Therefore, NR2D-containing eNMDARs may generate tonic I_{NMDA} in nondepolarized VP neurons under physiological levels of Mg^{2+} concentrations in the brain and contribute to upregulation of neuronal activity on an “on-demand” basis, while minimizing potential excitotoxic effects because of their lower Ca^{2+} permeability. However, it is noteworthy that 7 d SL increased serum osmolality to an equivalent level in WT and NR2D KO mice, suggesting that there is central and/or peripheral adaptation maintaining fluid homeostasis in NR2D KO mice in the SL condition. Given that increases in plasma VP induced by salt intake promotes water retention and vasoconstriction (Prager-Khoutorsky et al., 2017), future studies are warranted to further investigate the functional significance of the Mg^{2+} -resistant tonic I_{NMDA} to facilitate VP release regulating sodium homeostasis, blood pressure, and kidney function in WT and NR2D KO mice.

This study shows that the recruitment of functional NR2D-containing NMDARs constitutes a novel mechanism by which VP neurons generate an Mg^{2+} -resistant tonic I_{NMDA} and contribute to adaptive firing responses to a sustained osmotic challenge. Moreover, our results expand the general mechanisms by which neurons in the mature brain are capable of regulating a membrane potential-sensitive excitatory signaling modality in an “on-demand” manner.

References

- Al-Ghoul WM, Meeker RB, Greenwood RS (1997) Differential expression of five N-methyl-D-aspartate receptor subunit mRNAs in vasopressin and oxytocin neuroendocrine cells. *Brain Res Mol Brain Res* 44:262–272.
- Asztely F, Erdemli G, Kullmann DM (1997) Extrasynaptic glutamate spillover in the hippocampus: dependence on temperature and the role of active glutamate uptake. *Neuron* 18:281–293.
- Bai N, Hayashi H, Aida T, Namekata K, Harada T, Mishina M, Tanaka K (2013) Dock3 interaction with a glutamate-receptor NR2D subunit protects neurons from excitotoxicity. *Mol Brain* 6:22.
- Berretta N, Jones RS (1996) Tonic facilitation of glutamate release by presynaptic N-methyl-D-aspartate autoreceptors in the entorhinal cortex. *Neuroscience* 75:339–344.
- Bresink I, Benke TA, Collett VJ, Seal AJ, Parsons CG, Henley JM, Collingridge GL (1996) Effects of memantine on recombinant rat NMDA receptors expressed in HEK 293 cells. *Br J Pharmacol* 119:195–204.
- Brickley SG, Misra C, Mok MHS, Mishina M, Cull-Candy SG (2003) NR2B and NR2D subunits coassemble in cerebellar Golgi cells to form a distinct NMDA receptor subtype restricted to extrasynaptic sites. *J Neurosci* 23:4958–4966.
- Brothwell SL, Barber JL, Monaghan DT, Jane DE, Gibb AJ, Jones S (2008) NR2B- and NR2D-containing synaptic NMDA receptors in developing rat substantia nigra pars compacta dopaminergic neurones. *J Physiol* 586:739–750.
- Carreño FR, Walch JD, Dutta M, Nedungadi TP, Cunningham JT (2011) Brain-derived neurotrophic factor-tyrosine kinase B pathway mediates NMDA receptor NR2B subunit phosphorylation in the supraoptic nuclei following progressive dehydration. *J Neuroendocrinol* 23:894–905.
- Chenard BL, Menniti FS (1999) Antagonists selective for NMDA receptors containing the NR2B subunit. *Curr Pharm Des* 5:381–404.
- Ciura S, Bourque CW (2006) Transient receptor potential vanilloid 1 is required for intrinsic osmoreception in organum vasculosum lamina terminalis neurons and for normal thirst responses to systemic hyperosmolality. *J Neurosci* 26:9069–9075.
- Clark BA, Cull-Candy SG (2002) Activity-dependent recruitment of extrasynaptic NMDA receptor activation at an AMPA receptor-only synapse. *J Neurosci* 22:4428–4436.
- Decavel C, Curras MC (1997) Increased expression of the N-methyl-D-aspartate receptor subunit, NR1, in immunohistochemically identified magnocellular hypothalamic neurons during dehydration. *Neuroscience* 78:191–202.
- Doherty FC, Sladek CD (2011) NMDA receptor subunit expression in the supraoptic nucleus of adult rats: dominance of NR2B and NR2D. *Brain Res* 1388:89–99.
- Dunah AW, Yasuda RP, Wang YH, Luo J, Dávila-García M, Gbadegesin M, Vicini S, Wolfe BB (1996) Regional and ontogenic expression of the NMDA receptor subunit NR2D protein in rat brain using a subunit-specific antibody. *J Neurochem* 67:2335–2345.
- Erreger K, Geballe MT, Kristensen A, Chen PE, Hansen KB, Lee CJ, Yuan H, Le P, Lyuboslavsky PN, Micale N, Jørgensen L, Clausen RP, Wyllie DJA, Snyder JP, Traynelis SF (2007) Subunit-specific agonist activity at NR2A-, NR2B-, NR2C-, and NR2D-containing N-methyl-D-aspartate glutamate receptors. *Mol Pharmacol* 72:907–920.
- Feng B, Tse HW, Skifter DA, Morley R, Jane DE, Monaghan DT (2004) Structure-activity analysis of a novel NR2C/NR2D-preferring NMDA receptor antagonist: 1-(phenanthrene-2-carbonyl) piperazine-2,3-dicarboxylic acid. *Br J Pharmacol* 141:508–516.
- Fleming TM, Scott V, Naskar K, Joe N, Brown CH, Stern JE (2011) State-dependent changes in astrocyte regulation of extrasynaptic NMDA receptor signalling in neurosecretory neurons. *J Physiol* 589:3929–3941.
- Hansen KB, Yi F, Perszyk RE, Furukawa H, Wollmuth LP, Gibb AJ, Traynelis SF (2018) Structure, function, and allosteric modulation of NMDA receptors. *J Gen Physiol* 150:1081–1105.
- Hanson E, Armbruster M, Lau LA, Sommer ME, Klatf ZJ, Swanger SA, Traynelis SF, Moss SJ, Noubary F, Chadchankar J, Dulla CG (2019) Tonic activation of GluN2C/GluN2D-containing NMDA receptors by ambient glutamate facilitates cortical interneuron maturation. *J Neurosci* 39:3611–3626.
- Hardingham GE, Fukunaga Y, Bading H (2002) Extrasynaptic NMDARs oppose synaptic NMDARs by triggering CREB shut-off and cell death pathways. *Nat Neurosci* 5:405–414.
- Harney SC, Jane DE, Anwyl R (2008) Extrasynaptic NR2D-containing NMDARs are recruited to the synapse during LTP of NMDAR-EPSCs. *J Neurosci* 28:11685–11694.
- Herman MA, Jahr CE (2007) Extracellular glutamate concentration in hippocampal slice. *J Neurosci* 27:9736–9741.
- Honda K, Negoro H, Dyball RE, Higuchi T, Takano S (1990) The osmoreceptor complex in the rat: evidence for interactions between the supraoptic and other diencephalic nuclei. *J Physiol* 431:225–241.
- Huang Z, Gibb AJ (2014) Mg^{2+} block properties of triheteromeric GluN1-GluN2B-GluN2D NMDA receptors on neonatal rat substantia nigra pars compacta dopaminergic neurones. *J Physiol* 592:2059–2078.
- Huettnet JE, Bean BP (1988) Block of N-methyl-D-aspartate-activated current by the anticonvulsant MK-801: selective binding to open channels. *Proc Natl Acad Sci U S A* 85:1307–1311.
- Ikeda K, Araki K, Takayama C, Inoue Y, Yagi T, Aizawa S, Mishina M (1995) Reduced spontaneous activity of mice defective in the epsilon 4 subunit of the NMDA receptor channel. *Brain Res Mol Brain Res* 33:61–71.
- Joe N, Scott V, Brown CH (2014) Glial regulation of extrasynaptic NMDA receptor-mediated excitation of supraoptic nucleus neurones during dehydration. *J Neuroendocrinol* 26:35–42.
- Katoh A, Fujiwara H, Ohbuchi T, Onaka T, Hashimoto T, Kawata M, Suzuki H, Ueta Y (2011) Highly visible expression of an oxytocin-monomeric red fluorescent protein 1 fusion gene in the hypothalamus and posterior pituitary of transgenic rats. *Endocrinology* 152:2768–2774.
- Krupp JJ, Vissel B, Heinemann SF, Westbrook GL (1998) N-terminal domains in the NR2 subunit control desensitization of NMDA receptors. *Neuron* 20:317–327.
- Kuner T, Schoepfer R (1996) Multiple structural elements determine subunit specificity of Mg^{2+} block in NMDA receptor channels. *J Neurosci* 16:3549–3558.
- Laurie DJ, Seeburg PH (1994) Ligand affinities at recombinant N-methyl-D-aspartate receptors depend on subunit composition. *Eur J Pharmacol* 268:335–345.

- Le Meur K, Galante M, Angulo MC, Audinat E (2007) Tonic activation of NMDA receptors by ambient glutamate of non-synaptic origin in the rat hippocampus. *J Physiol* 580:373–383.
- Lee SK, Lee S, Shin SY, Ryu PD, Lee SY (2012) Single cell analysis of voltage-gated potassium channels that determines neuronal types of rat hypothalamic paraventricular nucleus neurons. *Neuroscience* 205:49–62.
- Liu Q, Wong-Riley MT (2010) Postnatal development of N-methyl-D-aspartate receptor subunits 2A, 2B, 2C, 2D, and 3B immunoreactivity in brain stem respiratory nuclei of the rat. *Neuroscience* 171:637–654.
- Lozovaya NA, Grebenyuk SE, Tsintsadze TS, Feng B, Monaghan DT, Krishtal OA (2004) Extrasynaptic NR2B and NR2D subunits of NMDA receptors shape “superslow” afterburst EPSC in rat hippocampus. *J Physiol* 558:451–463.
- Momiyama A, Feldmeyer D, Cull-Candy SG (1996) Identification of a native low-conductance NMDA channel with reduced sensitivity to Mg^{2+} in rat central neurones. *J Physiol* 494:479–492.
- Monyer H, Burnashev N, Laurie DJ, Sakmann B, Seeburg PH (1994) Developmental and regional expression in the rat brain and functional properties of four NMDA receptors. *Neuron* 12:529–540.
- Mullasseril P, Hansen KB, Vance KM, Ogden KK, Yuan H, Kurtkaya NL, Santangelo R, Orr AG, Le P, Vellano KM, Liotta DC, Traynelis SF (2010) A subunit-selective potentiator of NR2C- and NR2D-containing NMDA receptors. *Nat Commun* 1:90.
- Pandit S, Jo JY, Lee SU, Lee YJ, Lee SY, Ryu PD, Lee JU, Kim HW, Jeon BH, Park JB (2015) Enhanced astroglial GABA uptake attenuates tonic GABA inhibition of the presympathetic hypothalamic paraventricular nucleus neurons in heart failure. *J Neurophysiol* 114:914–926.
- Paoletti P, Bellone C, Zhou Q (2013) NMDA receptor subunit diversity: impact on receptor properties, synaptic plasticity and disease. *Nat Rev Neurosci* 14:383–400.
- Papouin T, Oliet SH (2014) Organization, control and function of extrasynaptic NMDA receptors. *Philos Trans R Soc Lond B Biol Sci* 369:20130601.
- Papouin T, Ladépêche L, Ruel J, Sacchi S, Labasque M, Hanini M, Groc L, Pollegioni L, Mothet JP, Oliet SH (2012) Synaptic and extrasynaptic NMDA receptors are gated by different endogenous coagonists. *Cell* 150:633–646.
- Park JB, Skalska S, Stern JE (2006) Characterization of a novel tonic gamma-aminobutyric acid receptor-mediated inhibition in magnocellular neurosecretory neurons and its modulation by glia. *Endocrinology* 147:3746–3760.
- Park JB, Skalska S, Son S, Stern JE (2007) Dual GABA_A receptor-mediated inhibition in rat presympathetic paraventricular nucleus neurons. *J Physiol* 582:539–551.
- Poulain DA, Wakerley JB (1982) Electrophysiology of hypothalamic magnocellular neurones secreting oxytocin and vasopressin. *Neuroscience* 7:773–808.
- Prager-Khoutorsky M, Choe KY, Levi DI, Bourque CW (2017) Role of vasopressin in rat models of salt-dependent hypertension. *Curr Hypertens Rep* 19:42.
- Renaud LP, Bourque CW (1991) Neurophysiology and neuropharmacology of hypothalamic magnocellular neurons secreting vasopressin and oxytocin. *Prog Neurobiol* 36:131–169.
- Sah P, Hestrin S, Nicoll RA (1989) Tonic activation of NMDA receptors by ambient glutamate enhances excitability of neurons. *Science* 246:815–818.
- Sayer RJ, Hubbard JL, Sirett NE (1984) Rat organum vasculosum laminae terminalis in vitro: responses to transmitters. *Am J Physiol* 247:R374–R379.
- Shimamoto K, Lebrun B, Yasuda-Kamatani Y, Sakaitani M, Shigeri Y, Yumoto N, Nakajima T (1998) DL-threo-beta-benzoyloxyaspartate, a potent blocker of excitatory amino acid transporters. *Mol Pharmacol* 53:195–201.
- Siegler Retchless B, Gao W, Johnson JW (2012) A single GluN2 subunit residue controls NMDA receptor channel properties via intersubunit interaction. *Nat Neurosci* 15:406–413, S1–S2.
- Stachniak TJ, Trudel E, Bourque CW (2014) Cell-specific retrograde signals mediate antiparallel effects of angiotensin II on osmoreceptor afferents to vasopressin and oxytocin neurons. *Cell Rep* 8:355–362.
- Sun W, Wong JM, Gray JA, Carter BC (2018) Incomplete block of NMDA receptors by intracellular MK-801. *Neuropharmacology* 143:122–129.
- Swanson LW, Sawchenko PE (1983) Hypothalamic integration: organization of the paraventricular and supraoptic nuclei. *Annu Rev Neurosci* 6:269–324.
- Theodosis DT, Poulain DA (1993) Activity-dependent neuronal-glia and synaptic plasticity in the adult mammalian hypothalamus. *Neuroscience* 57:501–535.
- Tovar KR, Westbrook GL (1999) The incorporation of NMDA receptors with a distinct subunit composition at nascent hippocampal synapses *in vitro*. *J Neurosci* 19:4180–4188.
- Traynelis SF, Wollmuth LP, McBain CJ, Menniti FS, Vance KM, Ogden KK, Hansen KB, Yuan H, Myers SJ, Dingledine R (2010) Glutamate receptor ion channels: structure, regulation, and function. *Pharmacol Rev* 62:405–496.
- Tweedle CD, Hatton GI (1977) Ultrastructural changes in rat hypothalamic neurosecretory cells and their associated glia during minimal dehydration and rehydration. *Cell Tissue Res* 181:59–72.
- Tweedle CD, Hatton GI (1984) Synapse formation and disappearance in adult rat supraoptic nucleus during different hydration states. *Brain Res* 309:373–376.
- Ueta Y, Fujihara H, Serino R, Dayanithi G, Ozawa H, Matsuda K, Kawata M, Yamada J, Ueno S, Fukuda A, Murphy D (2005) Transgenic expression of enhanced green fluorescent protein enables direct visualization for physiological studies of vasopressin neurons and isolated nerve terminals of the rat. *Endocrinology* 146:406–413.
- Vicini S, Wang JF, Li JH, Zhu WJ, Wang YH, Luo JAH, Wolfe BB, Grayson DR (1998) Functional and pharmacological differences between recombinant N-methyl-D-aspartate receptors. *J Neurophysiol* 79:555–566.
- Villarroel A, Regalado MP, Lerma J (1998) Glycine-independent NMDA receptor desensitization: localization of structural determinants. *Neuron* 20:329–339.
- Watanabe M, Inoue Y, Sakimura K, Mishina M (1992) Developmental changes in distribution of NMDA receptor channel subunit mRNAs. *Neuroreport* 3:1138–1140.
- Wenzel A, Villa M, Mohler H, Benke D (1996) Developmental and regional expression of NMDA receptor subtypes containing the NR2D subunit in rat brain. *J Neurochem* 66:1240–1248.
- Wu YN, Johnson SW (2015) Memantine selectively blocks extrasynaptic NMDA receptors in rat substantia nigra dopamine neurons. *Brain Res* 1603:1–7.
- Wu YW, Grebenyuk S, McHugh TJ, Rusakov DA, Semyanov A (2012) Backpropagating action potentials enable detection of extrasynaptic glutamate by NMDA receptors. *Cell Rep* 1:495–505.
- Wyllie DJA, Behe P, Colquhoun D (1998) Single-channel activations and concentration jumps: comparison of recombinant NR1a/NR2A and NR1a/NR2D NMDA receptors (vol 510, pg 1, 1998). *J Physiol* 510:1–18.
- Wyllie DJA, Livesey MR, Hardingham GE (2013) Influence of GluN2 subunit identity on NMDA receptor function. *Neuropharmacology* 74:4–17.
- Xia P, Chen HS, Zhang D, Lipton SA (2010) Memantine preferentially blocks extrasynaptic over synaptic NMDA receptor currents in hippocampal autapses. *J Neurosci* 30:11246–11250.
- Yamakura T, Shimoji K (1999) Subunit- and site-specific pharmacology of the NMDA receptor channel. *Prog Neurobiol* 59:279–298.
- Yamakura T, Mori H, Masaki H, Shimoji K, Mishina M (1993) Different sensitivities of NMDA receptor channel subtypes to non-competitive antagonists. *Neuroreport* 4:687–690.
- Yang CR, Senatorov VV, Renaud LP (1994) Organum vasculosum lamina terminalis-evoked postsynaptic responses in rat supraoptic neurones in vitro. *J Physiol* 477:59–74.
- Yokoyama T, Saito T, Ohbuchi T, Hashimoto H, Suzuki H, Otsubo H, Fujihara H, Nagatomo T, Ueta Y (2010) TRPV1 gene deficiency attenuates miniature EPSC potentiation induced by mannitol and angiotensin II in supraoptic magnocellular neurons. *J Neurosci* 30:876–884.