

Research Articles | Systems/Circuits

## Comparing the limbic-frontal connectome across the primate order: conservation of connections and implications for translational neuroscience

<https://doi.org/10.1523/JNEUROSCI.0377-25.2026>

Received: 14 February 2025

Revised: 29 November 2025

Accepted: 19 January 2026

Copyright © 2026 Folloni et al.

This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](#), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

---

*This Early Release article has been peer reviewed and accepted, but has not been through the composition and copyediting processes. The final version may differ slightly in style or formatting and will contain links to any extended data.*

**Alerts:** Sign up at [www.jneurosci.org/alerts](http://www.jneurosci.org/alerts) to receive customized email alerts when the fully formatted version of this article is published.

**Comparing the limbic-frontal connectome across the primate order: conservation of connections and implications for translational neuroscience**

Davide Folloni<sup>1,2,3\*</sup>, Lea Roumazeilles<sup>4</sup>, Katherine L Bryant<sup>4,5</sup>, Paul R Manger<sup>6</sup>, Mads F Bertelsen<sup>7</sup>, Alexandre A Khrapitchev<sup>8</sup>, Peter H Rudebeck<sup>1,2</sup>, Rogier B Mars<sup>4,9</sup>

**Affiliations:**

<sup>1</sup> Nash Family Department of Neuroscience and Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place, New York, NY, 10029, USA

<sup>2</sup> Lipschultz Center for Cognitive Neuroscience, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA

<sup>3</sup> Vita-Salute San Raffaele University, Milan, Italy

<sup>4</sup> Wellcome Centre for Integrative Neuroimaging Nuffield Department of Clinical Neurosciences, John Radcliffe Hospital, University of Oxford, Headington, Oxford OX3 9DU, United Kingdom

<sup>5</sup> Laboratoire de Psychologie Cognitive, Aix-Marseille Université, Marseille, France

<sup>6</sup> School of Anatomical Sciences, Faculty of Health Sciences, University of Witwatersrand, Johannesburg, South Africa

<sup>7</sup> Centre for Zoo and Wild Animal Health, Copenhagen Zoo, Frederiksberg, Denmark

<sup>8</sup> Department of Oncology, Cancer Research UK and Medical Research Council Oxford Institute for Radiation Oncology, University of Oxford, Oxford, United Kingdom;

<sup>9</sup> Donders Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen, the Netherlands

\*To whom correspondence should be addressed:

Dr. Davide Folloni,

Icahn School of Medicine at Mount Sinai

One Gustave L. Levy Place

New York, NY, 10029, USA

Tel: +1 212 824-9307

E-mail: [folloni.davide@hsr.it](mailto:folloni.davide@hsr.it)

## ABSTRACT

The interaction of the limbic system and frontal cortex of the primate brain is important in many affective behaviors. For this reason, it is heavily implicated in a number of psychiatric conditions. This system is often studied in the macaque monkey, the most largely-used non-human primate model species. However, how evolutionary conserved this system is and how well results obtained in any model species translate to the human brain can only be understood by studying its organization across the primate order. Here, we present an investigation of the topology of limbic-frontal connections across seven species, representing all major branches of the primate family tree: humans (13 females, 11 males), chimpanzee (1 female), gorilla (1 male), gibbon (1 male), macaque (1 female, 2 males), squirrel monkey (1 female, 2 males), lemur (3 males). We show that dichotomous organization of amygdalofugal and uncinate connections with frontal cortex is conserved across all species. Subgenual connectivity of the cingulum bundle, however, seems less prominent in prosimian and New World monkey brains. These results inform both translational neuroscience and primate brain evolution.

## SIGNIFICANCE STATEMENT:

The interaction between the limbic system and the frontal cortex is critical for affective behaviors and is implicated in psychiatric conditions. While often studied in macaques, understanding how conserved these circuits are across primates is essential for translational relevance. Here, we investigate limbic-frontal connections across seven primate species, spanning all major evolutionary branches. We demonstrate that the dichotomous organization of amygdalofugal and uncinate pathways is conserved, while subcallosal cingulate connectivity of the cingulum bundle is less prominent in prosimian and New World monkeys. These findings provide key insights into the evolution of primate brains and enhance our understanding of the translational potential of non-human primate models for studying human brain function and disorders.

## INTRODUCTION

The limbic system and frontal cortex form an interconnected system of brain areas that are affected in disorders of mood, motivation, and anxiety (Kenwood et al., 2022; Klein-Flügge et al., 2022; H. Mayberg, 2001). Reciprocal connectivity between the parts that form this system are essential to support their roles in associative learning, decision-making and behavioral control (Munuera et al., 2018; Murray & Wise, 2010; Saez et al., 2015). As such, these connections are often the target of clinical interventions such as deep brain stimulation (Johansen-Berg et al., 2008; H. S. Mayberg et al., 2005; Riva-Posse et al., 2014). Advances in non-invasive stimulation of deep cortical structures (Folloni, Verhagen, et al., 2019; Folloni, 2022; Fouragnan et al., 2019; Verhagen et al., 2019) mean that functional interactions between connectional systems can potentially be targeted without surgical interventions. However, to advantageously and judiciously use these advances, a better understanding of the limbic-frontal connectome is required.

Much of our knowledge of these systems is based on invasive work in non-human animals, using techniques that cannot be ethically and legally applied to humans. The development of neuroimaging techniques that can be applied to both humans and non-human animals with increasing levels of resolution has enabled formal tests of how well results obtained in model species translate to humans (Mars et al., 2014). For example, comparative diffusion MRI-based tractography work has concentrated on assessing the principles of frontal cortex organization and connectivity in humans and macaque monkeys (Barrett et al., 2020; Croxson et al., 2005; Sallet et al., 2013). Recent investigations demonstrate how such methods can also be used to identify even very specific limbic-frontal connections. For instance, the amygdalofugal pathway, a small pathway running between the amygdala and frontal cortex (PFC) (Aggleton et al., 1980; Lehman et al., 2011; Nauta, 1961, 1962), could be detected in humans using tractography, as long as the approach is guided by anatomical knowledge obtained in macaques using both invasive tracers and tractography (Folloni, Sallet, et al., 2019). Comparative neuroimaging can thus be a valuable tool for investigating how limbic-frontal circuits vary across species.

Neuroimaging has the additional advantage that it can be applied to a larger number of species than can often be used in comparative neuroanatomical studies. Comparisons do not have to be limited to studies of humans and macaques and can be extended to a broader range of primate species, allowing the characterization of patterns of evolutionary consistency or change. Understanding such patterns will lead to a better understanding of how well results obtained in any model species are likely to generalize to the species of interest, which in most cases is the human. The increased availability of MRI datasets from non-human primates

(Milham et al., 2018; Tandler et al., 2022), combined with new analytical frameworks (Mars et al., 2021), make such studies feasible.

Frontal cortex is thought to be quite variable across primate species, both in term of its relative size and connectivity with other brain regions (Donahue et al., 2018; Mars et al., 2018; Passingham & Smaers, 2014). Therefore, we set out to compare the major pathways that connect to frontal cortex from the limbic system. It is possible that organizational variances of these frontal cortical areas among primates may be associated with differently organized patterns of connectivity. Alternatively, the course and organization of the amygdala-cingulate-prefrontal circuits may be conserved across primate species. To test these possibilities, we reconstructed the cingulum bundle (CB), ventral amygdalofugal pathway (AmF), and the uncinate fascicle (UF) across an unprecedented range of primate species using probabilistic tractography (**Figure 1**). We observed that, at the level of diffusion-tractography estimates of anatomical connectivity, the amygdala-cingulate-prefrontal circuits are similar in their course and overall pattern of organization not only in anthropoid brains, but also in prosimians.

## METHODS

### *Data acquisition and preprocessing*

**Human data.** Human in vivo diffusion MRI data were provided by the Human Connectome Project (HCP), WU-Minn Consortium (Principal Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience Research; and by the McDonnell Center for Systems Neuroscience at Washington University (Van Essen et al., 2013). The minimally preprocessed datasets of the first twenty-four subjects from the Q2 public data release were used (age range 22-35 years; thirteen females). Data acquisition and preprocessing methods are detailed in Ugurbil et al. (Ugurbil et al., 2013), Sotiropoulos et al. (Sotiropoulos et al., 2013), and Glasser et al. (Glasser et al., 2013). The diffusion data were collected at a 1.25 mm isotropic resolution across the entire brain on a customized 3T Siemens Skyra scanner using a monopolar Stejskal-Tanner diffusion scheme with a slice-accelerated EPI readout. Sampling in q-space included 3 shells at  $b = 1000, 2000, \text{ and } 3000 \text{ s/mm}^2$ . For each shell 90 diffusion gradient directions and 6 non-diffusion weighted images ( $b = 0 \text{ s/mm}^2$ ) were acquired with reversed phase-encoding direction for TOPUP distortion correction (Andersson et al., 2003). Voxel-wise model fitting of diffusion orientations was performed using FSL's bedpostX to fit a crossing fiber model to the data (Behrens et al., 2007). A multi-shell extension was used to

reduce overfitting of crossing fibers due to non-monoexponential diffusion decay (Jbabdi et al., 2012). Up to three fiber orientations per voxel were allowed.

Chimpanzee and gorilla. Whole brain samples from one chimpanzee (*Pan troglodytes*, 1 female, 54 years of age at time of death) and one gorilla (*Gorilla gorilla*, male, 12 years of age at time of death) were obtained from the London Zoological Society and the Primate Brain Bank. The samples were acquired from animals euthanized for management reasons unrelated to this research project. These brains were collected within minutes following death by initial perfusion with cold saline followed by cold paraformaldehyde. Details of sample preparation, scanning procedure, and data preprocessing have been communicated before (K. L. Bryant et al., 2021; Roumazeilles et al., 2020). In brief, samples were stored in formalin, rehydrated using a phosphate-buffered saline solution one week prior to scanning and placed in fluorinert for the scanning procedure. Samples were imaged using a 7T whole body scanner with a 28-channel knee coil (QED). Diffusion MRI data were acquired using a 3D diffusion weighted steady state free precession (DW-SSFP) pulse sequence (McNab et al., 2009). DW-SSFP data comprising of 240 diffusion-weighted ( $q = 300 \text{ cm}^{-1}$ , resolution =  $0.6 \times 0.6 \times 0.6 \text{ mm}^3$ ) and six non-diffusion weighted ( $q = 20 \text{ cm}^{-1}$ ) imaging volumes were acquired over the whole post mortem brains. To account for the T1, T2 and flip-angle dependencies of the DW-SSFP signal (Buxton, 1993; McNab & Miller, 2010), T1, T2 and B1 datasets were acquired via a turbo inversion-recovery (TIR), turbo spin-echo (TSE), and actual flip angle imaging (AFI) acquisition (Yarnykh, 2007). Prior to processing, a Gibbs ringing correction (Kellner et al., 2016) was applied to the DW-SSFP, TIR and TSE datasets. Quantitative T1 and T2 maps were generated from the TIR and TSE datasets assuming mono-exponential signal evolution. A B1 map was generated from the AFI data following the methodology described in (Yarnykh, 2007). All co-registrations within and between imaging modalities were performed with FLIRT (Jenkinson et al., 2002; Jenkinson & Smith, 2001) via a six degrees of freedom (translation and rotation) transformation. The DW-SSFP data, along with the T1, T2 and B1 maps were fitted with the full DW-SSFP signal equation (Buxton, 1993) to both a diffusion tensor model and a ball-and-two-stick model using cuDIMOT (Hernandez-Fernandez et al., 2019).

Gibbon, macaque monkey, squirrel monkey, and lemur data. Whole-brain samples were obtained from one gibbon (*H. lar*, male, 5.5 years old), three rhesus macaques (*M. mulatta*, between 11 and 15 years old, 1 female, 2 males), three squirrel monkeys (*S. boliviensis*, between 2 and 19 years old, 1 female, 2 males) and three ring-tailed lemurs (*L. catta*, between 3 and 11 years, 3 males). The samples were obtained from Copenhagen Zoo (gibbon, squirrel monkeys, and lemurs), the London Zoological Society (squirrel monkey), and the University of Oxford's Biomedical Sciences (macaques). The samples were acquired from deceased

animals that died of causes unrelated to this research project. Details of sample preparation, scanning procedure, and data preprocessing have been described previously (Bryant et al., 2023; Roumazeilles et al., 2022). In brief, all brains were extracted and fixed within 24 h after the death of the animal. All brains were rehydrated in a PBS solution 1 week prior to scanning and placed in fomblin or fluorinert for the scanning procedure. The diffusion-weighted MRI data were acquired from the whole brain using a 7 T preclinical MRI scanner (Varian, Oxford UK). The scanner bore diameter is 210 mm, the gradient coil references are the following: 205\_120\_HD (Varian, Oxford UK) with a Gmax of 50 G/cm. The radiofrequency coil was made by Rapid Biomedical GmbH (Rimpar, Germany) and is a birdcage transmit receive coil with 72 mm ID. We used a 2D diffusion-weighted spin-echo multi-slice protocol with single line readout (DW-SEMS; TR = 10 s; TE = 26 ms; Matrix size = 128 × 128 with a sufficient number of slices to cover each brain; resolution for ring-tailed lemurs: 0.5 mm<sup>3</sup>; resolution for gibbon and macaques: 0.6 mm<sup>3</sup>; resolution for squirrel monkey 0.5 mm<sup>3</sup>. A total of 16 non-diffusion-weighted (b = 0 s/mm<sup>2</sup>) and 128 diffusion-weighted (b = 4000 s/mm<sup>2</sup>) volumes were acquired with diffusion encoding directions evenly distributed over the whole sphere (single shell protocol). Diffusion tensors were fitted to each voxel using DTIFIT (Behrens et al., 2003). Subsequent preprocessing and analytical steps were specific for each species and are described in the following sections. Voxel-wise model fitting of diffusion orientations was performed using FSL's BedpostX to fit a crossing fiber model to the data (Behrens et al., 2007).

### *Probabilistic tractography*

Tractography was run using FSL's probtrackx2 using the following parameters. Probabilistic tractography does not suffer from some of the problems inherent in deterministic methods, in that it allows us to visualize the full distributions of reconstructed pathways. We adjust some parameters to the size of the brain (Jbabdi et al., 2015), but most importantly we normalize each tract to its maximum within the sample, such that we can obtain a comparable distribution across species. The parameters, similar to our approach in previous studies, were kept as similar as possible across species for consistency and translatability. Human: maximum of 3200 steps per sample; 10000 samples; step size of 0.25 mm; curvature threshold of 0.2. Macaque: maximum 3200 steps per sample; 10000 samples; step size of 0.1 mm; curvature threshold of 0.2. Chimpanzee and Gorilla: maximum 3200 steps per sample; 10000 samples; step size of 0.1 mm; curvature threshold of 0.2. Gibbon: maximum 3200 steps per sample; 10000 samples; step size of 0.1 mm; curvature threshold of 0.2. Squirrel Monkey: maximum 3200 steps per sample; 10000 samples; step size of 0.1 mm; curvature threshold of 0.2. Ring-tailed lemur: maximum 3200 steps per sample; 10000 samples; step size of 0.1 mm; curvature

threshold of 0.2. It should be noted that diffusion MRI cannot distinguish between direct and indirect connections, so the identified fibers could represent either type of projection and travel in either direction.

#### *Comparative probabilistic tractography protocol*

Probabilistic tractography of the anterior limb of the amygdalofugal pathway (AmF), uncinate fasciculus (UF), and the frontal segment of the cingulum bundle (CB) was performed in diffusion space. Descriptions of anatomical locations were made by reference to previously published atlases (Nieuwenhuys et al., 2008; Paxinos et al., 2000), tract-tracing studies (Fudge et al., 2002; Lehman et al., 2011; Schmahmann & Pandya, 2006), and diffusion tractography studies (Croxson et al., 2005; Jbabdi et al., 2013).

In all species seed (**Suppl. Figure 1**), exclusion and waypoint masks were reconstructed in the same way with regard to established white and grey matter regions. The seed mask of the AmF was drawn in the sub-commissural white matter perforating the substantia innominata, a region between the dorsal amygdala and the bed nuclei of the stria terminalis (BNST). The mask was located medially to the ventral pallidum (VP) and sublenticular extended amygdala (SLEA) and dorsally to the nucleus basalis of Meynert (NBM) (Mai et al., 2015; Paxinos et al., 2000). Its position carefully aimed to reproduce macaque tract tracing studies (deCampo & Fudge, 2013; Fudge & Haber, 2001; Schmahmann & Pandya, 2006). In accordance with the local trajectory of the AmF (as estimated by tract tracing), the seed mask was constrained to contiguous voxels showing high fractional anisotropy in an anterior-posterior direction. UF was seeded axially in the anterior temporal lobe, in the white matter rostro-lateral to the amygdala. An axial seed mask was chosen to account for the strong curve of the fibers from dorsal-ventral to anterior-posterior orientation to the orientation as they enter into the frontal lobe (Folloni, Sallet, et al., 2019). The CB seed mask was drawn in a coronal plane capturing the WM dorsal and medial to the corpus callosum, within the cingulate gyrus. To increase the anatomical accuracy of our tracts, two additional tracts, the extreme/external capsule complex and the anterior limb of the internal capsule, were created and their seed masks used as exclusion regions for AmF, UF and CB. Excluding fibers related to these latter two tracts is important because they are expected to run or terminate in close proximity to AmF, UF and CB.

To constrain the tractography algorithm, exclusion masks were drawn in all species to exclude false-positive results in areas of high crossing fibers as follows: (1) within the basal ganglia to avoid picking up spurious subcortical tracts; (2) posterior to the seeds to prevent

the projections from running backwards as the prefrontal cortical streamlines were the focus of this study; (3) an axial slice at the level of the superior temporal gyrus to prevent tracts from running in a ventral direction in an unconstrained manner (except for fibers reaching the amygdala); (4) on the axial-coronal slices cutting across the thalamus, basal ganglia, and corpus callosum to exclude subcortical and callosal projections; (5) in the dorsal cingulate cortex to avoid leakage of tracts to nearby bundles as a result of high curvature of the tracts (except for CB tracking); (6) an L-shaped coronal mask from the paracingulate cortex and the inferior frontal sulcus to the vertex to exclude tracking in the superior longitudinal fascicle; and (7) a mask excluding the opposite hemisphere to only track ipsilateral tracts. In humans, an exclusion mask of the CSF was used in each subject to prevent fibers from jumping across sulci during tracking. Streamlines encountering any of the exclusion masks were excluded from the tractography results. A coronal waypoint section was drawn in the frontal lobe at the level of the caudal genu of the corpus callosum to ensure that the fibers emanating from the seeds were projecting to the prefrontal cortex.

Streamlines were seeded from each voxel within each subject's seed mask within the body of the tract of interest. We aimed to keep all seeding and tracking parameters constant across tracts and species, only adjusting step size based on brain size and white matter density. Each streamline followed local orientations sampled from the posterior distribution given by BedpostX (Behrens et al., 2007). A visitation map or tractogram was constructed for each individual in order to allow comparison of these maps between tracts, subjects, and species. To counteract underweighting of distant voxels, the tractograms were log-transformed. Subsequently, to allow direct comparison across tracts, subjects, and species, independent of brain size and scan resolution, we further standardized the data by dividing a tract's voxel values by the 75<sup>th</sup> percentile value across the tract, thereby removing potential bias of differences in numbers of streamlines. In both human and macaques, the focus of the investigation was on the tracts' connections with PFC in both hemispheres. If multiple subjects per species were available, the normalized tractogram was then warped to each species' group template and averaged into a species-specific tractogram. For the purposes of visualization only, the normalized tractograms were subsequently thresholded. Figures were created with BioRender.com and FSL (Smith et al., 2004).

#### *Data sharing statement*

Data were analyzed using tools from FSL (Smith et al., 2004), which is available online at [fsl.fmrib.ox.ac.uk](http://fsl.fmrib.ox.ac.uk). Human MRI data were obtained from the Human Connectome Project (Van Essen et al., 2013) and available online from [humanconnectome.org](http://humanconnectome.org). The non-human primate

MRI data are available from the Digital Brain Bank (Tendler et al., 2022) online at [open.win.ox.ac.uk/DigitalBrainBank](https://open.win.ox.ac.uk/DigitalBrainBank).

## RESULTS

We reconstructed the principal connections between the limbic system and the prefrontal cortex (PFC) in the human brain and compared their organization with that in great (chimpanzee and gorilla) and lesser (gibbon) apes, an Old World (macaque monkey) and a New World monkey (squirrel monkey), and a prosimian (ring-tailed lemur). The evolutionary relationship between these species is shown in Figure 1. The goal was to assess how these connectional systems may have varied or remained consistent among primates. Our anatomical reconstruction focused on the temporal-prefrontal rostral branch of the amygdalofugal pathway (AmF), the uncinate fascicle (UF), and the anterior section of the cingulum bundle (CB). We first describe reconstructions of all three tracts in the human and macaque monkey in detail and then use them as baselines for subsequent descriptions of similarities and differences among the other species investigated. Importantly, here we defined anatomical and comparative approaches to reconstruct the anterior connections of the CB, AmF, and UF as they innervate frontal and temporal cortex. The approach used in each species were designed to be as similar as possible, allowing us to directly compare the organization of these fibre tracts within the methodological limitations. This follows prior studies that describe different brains in terms of directly comparable, homologous, features (K. Bryant et al., 2023; Elston et al., 2011; Mars et al., 2018; Preuss & Wise, 2022).

### *Reconstruction of limbic-frontal connections in macaque and human*

The UF is a C-shaped fiber bundle that reciprocally connects the temporopolar cortex and amygdala with the orbital and ventrolateral PFC. In our reconstruction, in a ventral section, the macaque UF extended along the outer surface of the temporal lobe and curved through the superior temporal gyrus (Figure 2). While the main direction of fibers in this region is from dorsal to ventral, some fibers branched along a medial-to-lateral axis towards the amygdala, thus connecting adjacent fibers near the anterior temporal region and the lateral prefrontal cortex (Figure 2). At the level of the limen insulae, UF fibers traveled in the white matter ventral to the claustrum, ventrolateral to the putamen and globus pallidus, and medial to the insular cortex (Figure 2). The human UF maintained a similar C-shaped structure projecting between the rostromedial temporal regions and the ventrolateral prefrontal cortex (Figure 2). Like the macaque temporal UF, it exhibited an anatomical organization primarily along a dorsal-ventral

gradient, but with lateral fibers extending towards amygdaloid nuclei, parahippocampal areas, and rhinal areas (Figure 2).

Within the macaque prefrontal cortex, the UF exhibited strong connectivity with the white matter bordering the caudolateral orbitofrontal cortex (OFC), as well as opercular and inferior frontal areas. Additionally, a subset of fibers extended alongside the subgenual cingulate area and the medial bank of the posterior OFC (Figure 3-5). In this region, the more medial UF fibers joined with the fibers of the AmF and the CB to innervate the white matter of the anterior cingulate cortex (ACC) (Figure 3-5). Overall, the primary portion of the UF in the ventral prefrontal cortex spanned across the lateral, central, and medial orbital gyri, forming a complex network within the OFC that extended into the frontopolar cortex (Figure 2,4-5). Similarly, in the human, at the junction of the prefrontal and temporal cortices, the UF closely followed the insular cortex and areas within the caudal region of the inferior frontal gyrus (Figure 2). Along the basal ganglia, the human UF encompasses the lateral ventral striatum and extends to a more lateral section of the ventral prefrontal cortex (Figure 2). In this region, the main body of the macaque and human UF continues to traverse laterally, providing extensive innervation to the white matter of the ventrolateral prefrontal cortex. Further, a subset of fibers also courses more medially, adjacent to subgenual anterior cingulate cortex (sACC) and, more prominently, towards the posterior OFC alongside the AmF and the CB (Figure 3-5). Within the anterior region of the ventral prefrontal cortex, both macaque and human UF extends across medial and lateral orbital areas. A prominent bundle of fibers occupies a substantial portion of the white matter in the lateral OFC and adjacent cortical areas on the ventral surface of the brain in both species. Another branch of the UF extends further rostrally in the white matter adjacent to the frontopolar cortex (FPC) in both macaque and human brains (Figure 2,4-5).

In the macaque brain, the frontal portion of the AmF was represented by a thin group of fibers that ran in the white matter adjacent the longitudinal fissure. At the level of the anterior commissure (AC), the fibers in the medially projecting limb of the AmF pathway traveled within the white matter dorsal to the amygdala, between amygdala nuclei, sublentiform extended amygdala area and medial longitudinal fissure (Figure 2,4). In this species, one set of AmF fibers ran in the white matter region between the nucleus basalis of Meynert, the lentiform nucleus, the ventral pallidum, and the piriform cortex. These medial AmF fibers divided into different sets of projections. Some extended downward to innervate the anterior hypothalamic areas near the paraventricular-supraoptic nuclei, while others coursed dorsally to connect the amygdala with the thalamus, the BNST, and the septum of the macaque monkey. Another set of macaque AmF fibers ran in the white matter surrounding the substantia innominata, ventral pallidum, and the lower bank of the AC, entering, further rostrally, into the white matter beneath

the ventral striatum/nucleus accumbens (Figure 2,4). These AmF pathways were found in both macaque monkeys and humans. Despite the strong medial-lateral primary diffusion direction characterizing the AC, some weaker AmF fibers that overlapped with the AC were still detectable in our tractography data within both macaque monkey and human brains. It is important to note that as the AmF passes through the basal forebrain and beneath the striatum, it is likely to interact with other connectional systems.

Eventually, the anterior fibers of the AmF of macaques and humans extended into the orbital cortex and into the rostral PFC by running in the white matter adjacent to the subgenual anterior cingulate cortex (sACC or Brodmann area 25) and the adjacent areas in subcallosal cortex (Figure 3-5). Near the sACC, the AmF of macaques and humans split into two subsections. The first subsection ran upward alongside the CB in the anterior cingulate sulcus and the surrounding white matter (Figure 5). The second branch, on the other hand, stretched between the sACC and the frontopolar cortex, passing underneath the medial OFC between the gyrus rectus and the medial orbital gyrus (Figure 2,4-5).

Frontal connections of the macaque and human CB project along the full extent of the anterior cingulate cortex. This bundle is composed by a mixture of fibers entering the large cingulum bundle briefly to reach nearby areas or longer-distance connections reaching distal brain regions. Core CB fibers surround the corpus callosum throughout its course in PFC in both species. A subsection of fibers in rostral CB exits the main bundle and contains axons towards and arriving from both cortical and subcortical areas (Figure 5). At the level of subcallosal and orbitofrontal cortex, in both macaques and humans, CB fibers interact with axons travelling in the AmF and UF (Figure 3,5). Here, CB fibers either terminate by innervating adjacent areas or continue coursing in a rostral direction towards anterior PFC and frontopolar cortex. In both species, another set of connections continue running along the genu of the corpus callosum, next to area 23, 24 and 32, and more dorsally, in the CB branches extending into dorsal ACC regions, premotor, lateral PFC (partly through intermixing with UF fibers) and medial-dorsal PFC areas 8 and 9 (Figure 5).

Overall, these results, obtained by probabilistic tractography accounting for multiple possible fiber orientations (Behrens et al., 2007), show that AmF, UF and prefrontal CB fibers exhibit a high degree of similarity between macaque monkeys and humans (Figure 3-4). In both species, fibers from the three bundles merge and intertwine at the level of the subgenual cingulate cortex giving before partially separating and running independently in anterior prefrontal and temporal cortices (Figure 3-5). The protocols employed here to reconstruct UF, AmF, and CB in macaque and human were based on Folloni et al. (Folloni, Sallet, et al., 2019), tailoring them to obtain results best matching those previously observed using macaque tracer

data (Heilbronner & Haber, 2014; Oler et al., 2017). This enabled us to avoid false positives and false negatives that may be created when employing tractography algorithms and ensured the truthfulness of the reconstructed bundles. Figure 3 shows the macaque and human tractography results side by side with previously published tracer data to corroborate these original findings. Having established a protocol that allowed both matching of macaque tractography data with established macaque tracer data and matching of macaque and human data, we felt confident using these protocols in chimpanzee, gorilla, gibbon, squirrel monkey, and ring-tailed lemur. We describe the course of the tracts in these species at the levels of anterior temporal cortex, subcallosal frontal cortex, and more anterior prefrontal cortex.

#### *Course of the AmF and UF between limbic/temporal and frontal cortex across all species*

The most prominent shared feature observed across lemur, squirrel and macaque monkey, gibbon, gorilla, chimpanzee and human brains is their widespread connectivity between prefrontal and anterior temporal areas, both subcortical medial and lateral cortical structures (Figure 4). Consistently across all the species examined, fibers connect amygdala, piriform and inferior temporal cortices on the temporal side with medial and lateral regions in anterior PFC on the opposite end. As the anterior limb of the AmF and the UF travel between temporal and frontal poles, these bundles innervate striatal, insular, basal forebrain, ventromedial and ventrolateral PFC areas (Figure 2). Consistently across all the species examined, the AmF exits the amygdala via the medial wall of the temporal lobe and then runs dorsally towards limbic, striatal and prefrontal structures along other forebrain and thalamic fibers (Figure 2). The UF, instead, unfolds as a curve between the anterior temporal lobe and PFC carrying prefrontal, forebrain, amygdalar, rhinal, piriform, parahippocampal and temporal associative projections.

A clear distinction between AmF and UF is observable across the six primate species at the level of the amygdala, temporal cortex, and peri-commissural white matter, showing the dichotomous medial-lateral organization in striatal and basal forebrain we see in macaque and human (Figure 4). The AmF occupies the white matter perforating the ventral pallidum and expanding underneath the ventral striatum, primarily branching towards the anterior ventromedial striatum, rostrally, and the hypothalamus, bed nuclei, stria terminalis and septum, medially. UF fibers course through the sub-capsular white matter along the ventral insular cortex and putamen. A few fibers leave the main body of the UF and merge with the AmF, approaching from a central position. This ventral-dorsal intertwining of the two paths is observable across all primate brains examined and becomes complete when AmF and UF approach the subgenual ACC (Figure 2). In the white matter innervating the striatal and basal

forebrain territory, however, AmF and UF in the gibbon overlapped to a higher degree than in the other species, where they were still clearly distinguishable (Figure 2). Interestingly, the medial-lateral organization of UF and AmF is also preserved in the strepsirrhine lemur (Figure 2) despite the anatomical expansion of the temporal lobe and of the PFC in anthropoids (Eldridge et al., 2021; Gurche, 1982; Ni et al., 2019; Passingham & Wise, 2012).

#### *Level of the subcallosal cortex*

Fibers coursing alongside the subgenual ACC, underneath the ventral floor of the genu of the corpus callosum, display a common organization across Old World monkey, lesser and great ape, and human brains. After coursing through the temporal lobe and near the ventral striatum, AmF and UF approach the subgenual cortex from a medial and lateral position respectively. This is also where the main body of the cingulum bundle (CB) joins the territory of AmF and UF, but its main body runs primarily medial to AmF and UF (Figure 3,5,6). In both the macaque and human brains, the CB fibers merge with fibers from AF and UF in this region, confirming results from previous tract-tracing studies (Haber & Behrens, 2014).

In caudal subgenual ACC, AmF and UF not only begin sharing important portions of white matter with CB, but they also start merging into a single limbic bundle innervating ventral PFC and anterior temporal cortex, in addition to medial cortex at their extremes. Although AmF and UF here begin merging into a single connectional system as they interacted with CB they maintained a medio-lateral organisation until they approach posterior OFC. The most dorso-medial WM territory surrounding the subgenual ACC is primarily occupied by fibers carried by an additional dorsal limbic tract, the CB. Within the PFC, this tract carries fibers running in the middle, anterior and subgenual cingulate cortex (Figure 5-6).

Interestingly, CB fibers did not extend into the caudal sACC as clearly as in the Prosimian (ring-tailed lemur) and New World monkey (squirrel monkey) but we could still observe a similar organization of AmF and UF as described above for the macaque monkey, gibbon, gorilla, chimpanzee and human brains (Figure 6-7). This pattern was consistent even when adjusting for different values of the curvature parameter used in our analyses aiming to account for the anatomy of the genu of the corpus callosum in these two species. (Figure 2).

#### *Level of anterior prefrontal cortex*

Thus far, the organization of UF, AmF, and CB is similar across all of the primates studied here. However, prefrontal cortex has differentially expanded in different primate lineages, showing a relative increase in size in anthropoid primates (Passingham and Wise, 2012) and

perhaps again in the ape and especially the human lineage (Donahue et al., 2018; Passingham & Smaers, 2014). It is therefore of particular interest to examine how these tracts project to anterior prefrontal cortex.

A merging of amygdala, temporal cingulate and prefrontal connections in posterior OFC was clearly observable along all of the representatives of the primate order examined (Figure 4). As previously reported for macaque and human brains (Folloni, Sallet, et al., 2019), AmF and UF transition from a dichotomous organization in temporal and subcortical regions to an intertwined bundle in more anterior PFC white matter. Although in medial portions of anterior PFC, AmF, and UF occupy a similar portion of white matter, we observed that AmF occupies the territory adjacent to the medial frontal wall, whereas UF extends to lateral ventral areas.

In anterior PFC, AmF, and UF fibers coursed together along the medial orbital and frontopolar cortices consistently across lemurs, squirrel and macaque monkeys, gibbons, gorilla, chimpanzees and humans regardless of their frontal lobe expansion (Figure 4). In addition, UF fibers branched out to caudolateral OFC, ventro-lateral PFC as well as opercular and inferior frontal areas (Figure 4). Axons from both AmF and UF reached granular PFC in the primate species that possess this cell type. Anatomical interactions among AmF, UF and CB fibers in caudal-ventral PFC were shared across all species, except in the gibbon, where AmF and UF converged more posteriorly, beginning in the transition territory between subgenual and orbital cortex. Although we cannot entirely rule out the influence of false positives given the small sample size for this species, it is noticeable that the gibbon is the only species among those examined here that primarily lives in very small groups, and spends the least amount of time in socially and emotionally bonding behaviors (Cunningham & Mootnick, 2009; Guan et al., 2018). This peculiarity in social behavior, either it being environmentally induced or inherent in the gibbon nature, may be associated with our finding that there is a difference in AmF and UF organization in posterior OFC for this species.

## DISCUSSION

In this study, we reconstructed the limbic-prefrontal connections in species from all major branches of the primate radiation. By using the same methodology, we showed that the overall organization of AmF, UF, and CB is evolutionarily conserved, but with some species-specific variances (Figure 8). This is important given the need to improve translation from animal models to humans for developing new brain treatments. Notably, our diffusion tractography results reveal that the anatomy of limbic-frontal fibers resembled previous tracer investigations

focused on these specific tracts in macaque monkeys (Fudge et al., 2002; Fudge & Haber, 2001; Heilbronner & Haber, 2014; Kunishio & Haber, 1994; Lehman et al., 2011; Oler et al., 2017) and diffusion imaging studies that compared only a subset of the species investigated here (Folloni, Sallet, et al., 2019; Roumazeilles et al., 2020; Thiebaut de Schotten et al., 2012).

Primate brains, including humans, share a general organizational plan but exhibit species-specific adaptations (Preuss & Kaas, 2003). Temporal and frontal cortices expanded in hominoids (Braunsdorf et al., 2021; Chaplin et al., 2013; Donahue et al., 2018), accompanied by increased organizational complexity and differentiation in both grey and white matter (Barrett et al., 2020; Eichert et al., 2019; Roumazeilles et al., 2020). To our knowledge, this is the first study to compare the organization of specific white matter tracts across all major branches of the primate family tree. Previous work has either examined more global connectivity measures (Ardesch et al., 2022; Suarez et al., 2022) or focused on a limited number of species (Barrett et al., 2020; Roumazeilles et al., 2020).

Despite anatomical heterogeneities in temporal and frontal cortex across primates, the overall organization of the AmF, UF and CB appears conserved. In all species, amygdala-temporal-frontal connections travel in three distinct tracts with similar courses and relative topology. This suggests that their organization represents a shared anatomical feature, possibly originating in early primates. Inter-species similarities in connectivity within the anterior temporal and frontal lobes may be indicative of how the environment shaped white matter organization in early primates. Further, these shared features may also relate to common foraging strategies across primates, which rely heavily on visual foraging and associative learning (Bongioanni et al., 2021; Chau et al., 2015; Eacott & Gaffan, 1992; Folloni et al., 2021.; Fouragnan et al., 2019; Gutnikov et al., 1997; Joyce & Barbas, 2018; Murray & Wise, 2010; Passingham & Wise, 2012; Vogt, 2005; Wittmann et al., 2020).

Reconstructing fiber bundles in inhomogeneous tissue such as cingulate, striatal, and ventral prefrontal regions is challenging because fractional anisotropy does not fully capture tract integrity. This can produce false negatives or positives in deterministic tractography. By contrast, the probabilistic tractography approach used here, informed by prior anatomical knowledge, reveals a distribution and reconstruction of streamlines consistent with previous tracer studies.

It is important to properly contextualize this result. We show that the organization of two dissociable limbic-prefrontal pathways, the AmF and UF, in the basal forebrain and subgenual cortex, is conserved across primates. This is a particularly intriguing finding in lemurs and squirrel monkeys, which have markedly lower white matter density in this region. Our methodology focused on reconstructing the main bodies of fibre bundles throughout the

brain, a reliable approach for cross-species comparisons because it identifies homologous white-matter structures (Mars et al., 2018) and avoids the false positives common when inferring connectivity directly from grey matter (Maier-Hein et al., 2017). This approach underpins tools used to compare white matter architecture across species, development, and individuals (Warrington et al., 2020, 2022). However, this does mean we do make claims on the precise grey matter areas innervated by these bundles. Indeed, the conserved fiber systems identified in this work exist within neural systems that do show phylogenetic diversity, as mentioned above. A particularly prominent difference is the addition of granular areas in the prefrontal cortex (Preuss & Wise, 2022). Future work will determine how these converged pathways relate to diverse grey matter architecture (Eldridge et al., 2021) and the implications of this for function (Bramson et al., 2020).

Despite overall conservation, our results suggest a potential difference in innervation of subgenual cortex across species. Across primate evolution, cingulate fibres increasingly expand within the subcallosal cortex, transitioning from focal inputs in prosimians and New World monkeys to more expanded projections in macaques, gibbons, great apes, and humans. This expansion is accompanied by a widening separation between limbic and frontotemporal pathways ventrally, reflecting increased differentiation of amygdala-cingulate-prefrontal circuits.

Although still poorly understood, subcallosal cortex is a key hub for mood and affect, as dysregulation of this region is a core feature of patients with mood disorders and anhedonia (Drevets et al., 1997, 2008). Lesions of this area cause deficits in anticipatory arousal (Rudebeck et al., 2014), whereas hyperactivation is associated with autonomic changes and deficits in reward processing (Alexander et al., 2019) and aversive behavior (Alexander et al., 2019, 2020). These findings implicate subgenual cortex as critical to the orchestration of neural circuits regulating affective behavior and survival (Wood et al., 2023). In most species, the CB interacts in subgenual cortex with AmF and UF before the three bundles separate into their own specific trajectories. These results replicate tracer studies in Old-World monkeys (Heilbronner & Haber, 2014; Joyce & Barbas, 2018; Vogt, 2005). We did not observe the same organization in lemurs and squirrel monkeys: at comparable thresholds, none of the reconstructed cingulum fibres reached the subcallosal white matter, regardless of the curvature parameters used.

It is important to interpret this result carefully. It is unlikely that the cingulum entirely fails to reach caudal subgenual and ventromedial PFC in these species, as tracer work in squirrel monkeys shows fibres in this white matter region (Leichnetz & Astruc, 1976). However, the density of fibers reaching this territory, as indicated indirectly by the likelihood of the probabilistic tractography reaching there, is much lower in these species. This is

important, as the density of connections is a very important predictor of information exchange between areas (Honey et al., 2010). Our results suggest that the growing use of New World monkeys such as the squirrel monkey in translational research (Royo et al., 2021) should be paired with extensive validation studies to confirm that results from other model species replicate.

The overall preservation of the CB, AmF, and UF across prosimians, New- and Old-World monkeys, gibbons, and hominoids underscores a powerful translational opportunity not only in evolutionary terms but also for developing and improving either novel or existing technologies modulating activity in subcallosal region. This also cautions against directly translating findings between New- and Old-World monkey models without accounting for species differences. Nevertheless, even in prosimians and New-World monkeys, where subcallosal white-matter fibres are reduced relative to macaques and anthropoids, the organization and separation of limbic–prefrontal tracts are still evident. This is notable given the broader divergences in cortical cytoarchitecture, association-cortex expansion, and prefrontal lamination across these lineages. The presence of homologous pathway scaffolds in these evolutionarily distant species indicates that the core architecture supporting affective and cognitive integration is deeply conserved.

The white-matter similarities and differences identified here have direct relevance for therapeutic interventions targeting the subcallosal cortex, including deep-brain stimulation, transcranial ultrasound neuromodulation, and pharmacological approaches. These findings can inform species-specific parameter selection, interpretation of outcomes, and behavioural phenotypes across diverse models. Notably, the reduced extent of cingulate fibres in prosimians and New-World monkeys highlights that translational studies in these species must consider not only pathway presence but also quantitative features such as stimulation trajectory, strength, and depth—that may cause similar manipulations to engage amygdala–cingulate–prefrontal circuits differently across species.

This manuscript includes, to our knowledge, the largest set of species in which a specific group of tracts has been reconstructed in detail. Such reconstructions enable quantitative, whole-brain comparisons, now feasible with datasets like the one presented here (e.g., Assimopoulos et al., 2025). Although we outline organizational principles across several primate families, future work could broaden species coverage. For example, we included the prosimian *Lemur catta* but not smaller-brained species such as *Galago* or the Platyrrhine tamarin. Tarsiers, the only non-anthropoid haplorhines, are also interesting, though MRI data are lacking. Many limbic targets appear conserved across mammals (Murray et al., 2011), raising the question of how widely these principles extend. While tract reconstructions in non-primates remain limited (but see Inglis et al., 2024 and Wright et al., 2018), the recent release

of carnivore scans (Boch et al., 2025) provides an opportunity to examine these evolutionary patterns more broadly.

In conclusion, we show that despite differences in phylogeny, life history, behaviour, and ecology, the topology of limbic–frontal connections is shared not only across humans, apes, and monkeys, but also with simians and prosimians. Although these limbic-frontal connections innervate systems that vary in size and morphology, their overall architecture and course is conserved. An exception are the cingulum connections to subgenual cortex, a key target for brain therapies. Together, this work shows how a large-scale comparative approach can help guide translational neuroscience, as well as provide insights into brain evolution.

## Acknowledgements

The work of DF was supported by grants from the National Institute of Mental Health (1K99MH137315) and the European Research Council Starting grant (project number 101220843). The work of RBM was funded by the Biotechnology and Biological Sciences Research Council (BBSRC) UK [BB/N019814/1] and the Medical Research Council (MRC) UK [MR/Y010698/1]. The Wellcome Centre for Integrative Neuroimaging is supported by core funding from the Wellcome Trust [203139/Z/16/Z]. The work of KLB was funded by Institut Convergence ILCB (ANR-16-CONV-0002). For the purpose of Open Access, the author has applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission. Data will be uploaded to the Wellcome Centre for Integrative Neuroimaging Gitlab and released upon publication.

## Author contributions

D.F., R.B.M. planned and designed the study. D.F., R.B.M., P.M., M.F.B., A.A.K., acquired the data. D.F., R.B.M., L.R., K.L.B. analyzed the data. D.F., R.B.M., K.L.B., P.H.R., interpreted the results. All authors contributed to manuscript preparation.

## Competing interests

None to declare.

- Aggleton, J. P., Burton, M. J., & Passingham, R. E. (1980). Cortical and subcortical afferents to the amygdala of the rhesus monkey (*Macaca mulatta*). *Brain Research*, 190(2), 347–368. [https://doi.org/10.1016/0006-8993\(80\)90279-6](https://doi.org/10.1016/0006-8993(80)90279-6)
- Alexander, L., Gaskin, P. L. R., Sawiak, S. J., Fryer, T. D., Hong, Y. T., Cockcroft, G. J., Clarke, H. F., & Roberts, A. C. (2019). Fractionating Blunted Reward Processing Characteristic of Anhedonia by Over-Activating Primate Subgenual Anterior Cingulate Cortex. *Neuron*, 101(2), 307–320.e6. <https://doi.org/10.1016/j.neuron.2018.11.021>
- Alexander, L., Wood, C. M., Gaskin, P. L. R., Sawiak, S. J., Fryer, T. D., Hong, Y. T., McIver, L., Clarke, H. F., & Roberts, A. C. (2020). Over-activation of primate subgenual cingulate cortex enhances the cardiovascular, behavioral and neural responses to threat. *Nature Communications*, 11(1), 5386. <https://doi.org/10.1038/s41467-020-19167-0>
- Andersson, J. L. R., Skare, S., & Ashburner, J. (2003). How to correct susceptibility distortions in spin-echo echo-planar images: Application to diffusion tensor imaging. *NeuroImage*, 20(2), 870–888. [https://doi.org/10.1016/S1053-8119\(03\)00336-7](https://doi.org/10.1016/S1053-8119(03)00336-7)
- Ardesch, D. J., Scholtens, L. H., de Lange, S. C., Roumazeilles, L., Khrapitchev, A. A., Preuss, T. M., Rilling, J. K., Mars, R. B., & van den Heuvel, M. P. (2022). Scaling Principles of White Matter Connectivity in the Human and Nonhuman Primate Brain. *Cerebral Cortex*, 32(13), 2831–2842. <https://doi.org/10.1093/cercor/bhab384>
- Assimopoulos, S., Warrington, S., Folloni, D., Bryant, K., Tang, W., Jbabdi, S., Heilbronner, S., Mars, R. B., & Sotiropoulos, S. N. (2025). Cross-species Standardised Cortico-Subcortical Tractography. *eLife*, 14. <https://doi.org/10.7554/eLife.107012.1>
- Barrett, R. L. C., Dawson, M., Dyrby, T. B., Krug, K., Ptito, M., D'Arceuil, H., Croxson, P. L., Johnson, P. J., Howells, H., Forkel, S. J., Dell'Acqua, F., & Catani, M. (2020). Differences in Frontal Network Anatomy Across Primate Species. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 40(10), 2094–2107. <https://doi.org/10.1523/JNEUROSCI.1650-18.2019>
- Behrens, T. E. J., Berg, H. J., Jbabdi, S., Rushworth, M. F. S., & Woolrich, M. W. (2007). Probabilistic diffusion tractography with multiple fibre orientations: What can we gain? *NeuroImage*, 34(1), 144–155. <https://doi.org/10.1016/j.neuroimage.2006.09.018>
- Behrens, T. E. J., Woolrich, M. W., Jenkinson, M., Johansen-Berg, H., Nunes, R. G., Clare, S., Matthews, P. M., Brady, J. M., & Smith, S. M. (2003). Characterization and propagation of uncertainty in diffusion-weighted MR imaging. *Magnetic Resonance in Medicine*, 50(5), 1077–1088. <https://doi.org/10.1002/mrm.10609>
- Boch, M., Karadachka, K., Loh, K. K., Benn, R. A., Roumazeilles, L., Bertelsen, M. F., Manger, P. R., Wriggelsworth, E., Spiro, S., Spocter, M. A., Johnson, P. J., Avelino-de-Souza, K., Patzke, N., Lamm, C., Miller, K. L., Sallet, J., Khrapitchev, A. A., Tandler, B. C., & Mars, R. B. (2025). Comparative neuroimaging of the carnivoran brain: Neocortical sulcal anatomy. *eLife*, 13. <https://doi.org/10.7554/eLife.100851.2>
- Bongioanni, A., Folloni, D., Verhagen, L., Sallet, J., Klein-Flügge, M. C., & Rushworth, M. F. S. (2021). Activation and disruption of a neural mechanism for novel choice in monkeys. *Nature*, 591(7849), 270–274. <https://doi.org/10.1038/s41586-020-03115-5>
- Bramson, B., Folloni, D., Verhagen, L., Hartogsveld, B., Mars, R. B., Toni, I., & Roelofs, K. (2020). Human Lateral Frontal Pole Contributes to Control over Emotional Approach-Avoidance Actions. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 40(14), 2925–2934. <https://doi.org/10.1523/JNEUROSCI.2048-19.2020>

- Braunsdorf, M., Blazquez Freches, G., Roumazeilles, L., Eichert, N., Schurz, M., Uithol, S., Bryant, K. L., & Mars, R. B. (2021). Does the temporal cortex make us human? A review of structural and functional diversity of the primate temporal lobe. *Neuroscience & Biobehavioral Reviews*, 131, 400–410. <https://doi.org/10.1016/j.neubiorev.2021.08.032>
- Bryant, K. L., Ardesch, D. J., Roumazeilles, L., Scholtens, L. H., Khrapitchev, A. A., Tendler, B. C., Wu, W., Miller, K. L., Sallet, J., van den Heuvel, M. P., & Mars, R. B. (2021). Diffusion MRI data, sulcal anatomy, and tractography for eight species from the Primate Brain Bank. *Brain Structure and Function*, 226(8), 2497–2509. <https://doi.org/10.1007/s00429-021-02268-x>
- Bryant, K., Manger, P. R., Bertelsen, M. F., Khrapitchev, A. A., Sallet, J., Benn, R. A., & Mars, R. (2023). A comprehensive atlas of white matter tracts in a lesser ape, the lar gibbon. *Manuscript Submitted for Publication*.
- Buxton, R. B. (1993). The diffusion sensitivity of fast steady-state free precession imaging. *Magnetic Resonance in Medicine*, 29(2), 235–243. <https://doi.org/10.1002/mrm.1910290212>
- Chaplin, T. A., Yu, H.-H., Soares, J. G. M., Gattass, R., & Rosa, M. G. P. (2013). A Conserved Pattern of Differential Expansion of Cortical Areas in Simian Primates. *The Journal of Neuroscience*, 33(38), 15120–15125. <https://doi.org/10.1523/JNEUROSCI.2909-13.2013>
- Chau, B. K. H., Sallet, J., Papageorgiou, G. K., Noonan, M. P., Bell, A. H., Walton, M. E., & Rushworth, M. F. S. (2015). Contrasting Roles for Orbitofrontal Cortex and Amygdala in Credit Assignment and Learning in Macaques. *Neuron*, 87(5), 1106–1118. <https://doi.org/10.1016/j.neuron.2015.08.018>
- Croxson, P. L., Johansen-Berg, H., Behrens, T. E. J., Robson, M. D., Pinsk, M. A., Gross, C. G., Richter, W., Richter, M. C., Kastner, S., & Rushworth, M. F. S. (2005). Quantitative investigation of connections of the prefrontal cortex in the human and macaque using probabilistic diffusion tractography. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 25(39), 8854–8866. <https://doi.org/10.1523/JNEUROSCI.1311-05.2005>
- Cunningham, C., & Mootnick, A. (2009). Gibbons. *Current Biology: CB*, 19(14), R543-544. <https://doi.org/10.1016/j.cub.2009.05.013>
- deCampo, D. M., & Fudge, J. L. (2013). Amygdala projections to the lateral bed nucleus of the stria terminalis in the macaque: Comparison with ventral striatal afferents. *The Journal of Comparative Neurology*, 521(14), 3191–3216. <https://doi.org/10.1002/cne.23340>
- Donahue, C. J., Glasser, M. F., Preuss, T. M., Rilling, J. K., & Van Essen, D. C. (2018). Quantitative assessment of prefrontal cortex in humans relative to nonhuman primates. *Proceedings of the National Academy of Sciences of the United States of America*, 115(22), E5183–E5192. <https://doi.org/10.1073/pnas.1721653115>
- Drevets, W. C., Price, J. L., & Furey, M. L. (2008). Brain structural and functional abnormalities in mood disorders: Implications for neurocircuitry models of depression. *Brain Structure & Function*, 213(1–2), 93–118. <https://doi.org/10.1007/s00429-008-0189-x>
- Drevets, W. C., Price, J. L., Simpson, J. R., Todd, R. D., Reich, T., Vannier, M., & Raichle, M. E. (1997). Subgenual prefrontal cortex abnormalities in mood disorders. *Nature*, 386(6627), 824–827. <https://doi.org/10.1038/386824a0>
- Eacott, M. J., & Gaffan, D. (1992). Inferotemporal-frontal Disconnection: The Uncinate Fascicle and Visual Associative Learning in Monkeys. *The European Journal of Neuroscience*, 4(12), 1320–1332.

- Eichert, N., Verhagen, L., Folloni, D., Jbabdi, S., Khrapitchev, A. A., Sibson, N. R., Mantini, D., Sallet, J., & Mars, R. B. (2019). What is special about the human arcuate fasciculus? Lateralization, projections, and expansion. *Cortex; a Journal Devoted to the Study of the Nervous System and Behavior*, 118, 107–115. <https://doi.org/10.1016/j.cortex.2018.05.005>
- Eldridge, M. A. G., Hines, B. E., & Murray, E. A. (2021). The visual prefrontal cortex of anthropoids: Interaction with temporal cortex in decision making and its role in the making of “visual animals”. *Current Opinion in Behavioral Sciences*, 41, 22–29. <https://doi.org/10.1016/j.cobeha.2021.02.012>
- Elston, G. N., Benavides-Piccione, R., Elston, A., Manger, P. R., & DeFelipe, J. (2011). Pyramidal Cells in Prefrontal Cortex of Primates: Marked Differences in Neuronal Structure Among Species. *Frontiers in Neuroanatomy*, 5, 2. <https://doi.org/10.3389/fnana.2011.00002>
- Folloni, D. (2022). Ultrasound neuromodulation of the deep brain. *Science (New York, N.Y.)*, 377(6606), 589. <https://doi.org/10.1126/science.add4836>
- Folloni, D., Fouragnan, E., Wittmann, M. K., Roumazeilles, L., Tankelevitch, L., Verhagen, L., Attali, D., Aubry, J.-F., Sallet, J., & Rushworth, M. F. S. (2021). Ultrasound modulation of macaque prefrontal cortex selectively alters credit assignment–related activity and behavior. *Science Advances*, 7(51), eabg7700. <https://doi.org/10.1126/sciadv.abg7700>
- Folloni, D., Sallet, J., Khrapitchev, A. A., Sibson, N., Verhagen, L., & Mars, R. B. (2019). Dichotomous organization of amygdala/temporal-prefrontal bundles in both humans and monkeys. *eLife*, 8. <https://doi.org/10.7554/eLife.47175>
- Folloni, D., Verhagen, L., Mars, R. B., Fouragnan, E., Constans, C., Aubry, J.-F., Rushworth, M. F. S., & Sallet, J. (2019). Manipulation of subcortical and deep cortical activity in the primate brain using transcranial focused ultrasound stimulation. *Neuron*, 101, 1009–1116. <https://doi.org/10.1101/342303>
- Fouragnan, E. F., Chau, B. K. H., Folloni, D., Kolling, N., Verhagen, L., Klein-Flügge, M., Tankelevitch, L., Papageorgiou, G. K., Aubry, J.-F., Sallet, J., & Rushworth, M. F. S. (2019). The macaque anterior cingulate cortex translates counterfactual choice value into actual behavioral change. *Nature Neuroscience*, 22(5), 797. <https://doi.org/10.1038/s41593-019-0375-6>
- Fudge, J. L., & Haber, S. N. (2001). Bed nucleus of the stria terminalis and extended amygdala inputs to dopamine subpopulations in primates. *Neuroscience*, 104(3), 807–827.
- Fudge, J. L., Kunishio, K., Walsh, P., Richard, C., & Haber, S. N. (2002). Amygdaloid projections to ventromedial striatal subterritories in the primate. *Neuroscience*, 110(2), 257–275.
- Glasser, M. F., Sotiropoulos, S. N., Wilson, J. A., Coalson, T. S., Fischl, B., Andersson, J. L., Xu, J., Jbabdi, S., Webster, M., Polimeni, J. R., Van Essen, D. C., Jenkinson, M., & WU-Minn HCP Consortium. (2013). The minimal preprocessing pipelines for the Human Connectome Project. *NeuroImage*, 80, 105–124. <https://doi.org/10.1016/j.neuroimage.2013.04.127>
- Guan, Z.-H., Ma, C.-Y., Fei, H.-L., Huang, B., Ning, W.-H., Ni, Q.-Y., Jiang, X.-L., & Fan, P.-F. (2018). Ecology and social system of northern gibbons living in cold seasonal forests. *Zoological Research*, 39(4), 255–265. <https://doi.org/10.24272/j.issn.2095-8137.2018.045>
- Gurche, J. A. (1982). Early Primate Brain Evolution. In E. Armstrong & D. Falk (Eds), *Primate Brain Evolution: Methods and Concepts* (pp. 227–246). Springer US. [https://doi.org/10.1007/978-1-4684-4148-2\\_15](https://doi.org/10.1007/978-1-4684-4148-2_15)

- Gutnikov, S. A., Ma, Y. Y., & Gaffan, D. (1997). Temporo-frontal disconnection impairs visual-visual paired association learning but not configural learning in Macaca monkeys. *The European Journal of Neuroscience*, 9(7), 1524–1529. <https://doi.org/10.1111/j.1460-9568.1997.tb01507.x>
- Haber, S. N., & Behrens, T. E. J. (2014). The neural network underlying incentive-based learning: Implications for interpreting circuit disruptions in psychiatric disorders. *Neuron*, 83(5), 1019–1039. <https://doi.org/10.1016/j.neuron.2014.08.031>
- Heilbronner, S. R., & Haber, S. N. (2014). Frontal cortical and subcortical projections provide a basis for segmenting the cingulum bundle: Implications for neuroimaging and psychiatric disorders. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 34(30), 10041–10054. <https://doi.org/10.1523/JNEUROSCI.5459-13.2014>
- Hernandez-Fernandez, M., Regulý, I., Jbabdi, S., Giles, M., Smith, S., & Sotiropoulos, S. N. (2019). Using GPUs to accelerate computational diffusion MRI: From microstructure estimation to tractography and connectomes. *NeuroImage*, 188(December 2018), 598–615. <https://doi.org/10.1016/j.neuroimage.2018.12.015>
- Honey, C. J., Thivierge, J.-P., & Sporns, O. (2010). Can structure predict function in the human brain? *NeuroImage*, 52(3), 766–776. <https://doi.org/10.1016/j.neuroimage.2010.01.071>
- Inglis, F. M., Taylor, P. A., Andrews, E. F., Pascalau, R., Voss, H. U., Glen, D. R., & Johnson, P. J. (2024). A diffusion tensor imaging white matter atlas of the domestic canine brain. *Imaging Neuroscience (Cambridge, Mass.)*, 2, 1–21. [https://doi.org/10.1162/imag\\_a\\_00276](https://doi.org/10.1162/imag_a_00276)
- Jbabdi, S., Lehman, J. F., Haber, S. N., & Behrens, T. E. (2013). Human and monkey ventral prefrontal fibers use the same organizational principles to reach their targets: Tracing versus tractography. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 33(7), 3190–3201. <https://doi.org/10.1523/JNEUROSCI.2457-12.2013>
- Jbabdi, S., Sotiropoulos, S. N., Haber, S. N., Van Essen, D. C., & Behrens, T. E. (2015). Measuring macroscopic brain connections in vivo. *Nature Neuroscience*, 18(11), 1546–1555. <https://doi.org/10.1038/nn.4134>
- Jbabdi, S., Sotiropoulos, S. N., Savio, A. M., Graña, M., & Behrens, T. E. (2012). Model-based analysis of multi-shell diffusion MR data for tractography: How to get over fitting problems. *Magnetic Resonance in Medicine : Official Journal of the Society of Magnetic Resonance in Medicine / Society of Magnetic Resonance in Medicine*, 68(6), 1846–1855. <https://doi.org/10.1002/mrm.24204>
- Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved Optimization for the Robust and Accurate Linear Registration and Motion Correction of Brain Images. *NeuroImage*, 17(2), 825–841. <https://doi.org/10.1006/nimg.2002.1132>
- Jenkinson, M., & Smith, S. (2001). A global optimisation method for robust affine registration of brain images. *Medical Image Analysis*, 5(2), 143–156. [https://doi.org/10.1016/S1361-8415\(01\)00036-6](https://doi.org/10.1016/S1361-8415(01)00036-6)
- Johansen-Berg, H., Gutman, D. A., Behrens, T. E. J., Matthews, P. M., Rushworth, M. F. S., Katz, E., Lozano, A. M., & Mayberg, H. S. (2008). Anatomical connectivity of the subgenual cingulate region targeted with deep brain stimulation for treatment-resistant depression. *Cerebral Cortex (New York, N.Y.: 1991)*, 18(6), 1374–1383. <https://doi.org/10.1093/cercor/bhm167>
- Joyce, M. K. P., & Barbas, H. (2018). Cortical Connections Position Primate Area 25 as a Keystone for Interoception, Emotion, and Memory. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 38(7), 1677–1698. <https://doi.org/10.1523/JNEUROSCI.2363-17.2017>

- Kellner, E., Dhital, B., Kiselev, V. G., & Reisert, M. (2016). Gibbs-ringing artifact removal based on local subvoxel-shifts. *Magnetic Resonance in Medicine*, 76(5), 1574–1581. <https://doi.org/10.1002/mrm.26054>
- Kenwood, M. M., Kalin, N. H., & Barbas, H. (2022). The prefrontal cortex, pathological anxiety, and anxiety disorders. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*, 47(1), 260–275. <https://doi.org/10.1038/s41386-021-01109-z>
- Klein-Flügge, M. C., Jensen, D. E. A., Takagi, Y., Priestley, L., Verhagen, L., Smith, S. M., & Rushworth, M. F. S. (2022). Relationship between nuclei-specific amygdala connectivity and mental health dimensions in humans. *Nature Human Behaviour*, 6(12), 1705–1722. <https://doi.org/10.1038/s41562-022-01434-3>
- Kunishio, K., & Haber, S. N. (1994). Primate cingulostriatal projection: Limbic striatal versus sensorimotor striatal input. *The Journal of Comparative Neurology*, 350(3), 337–356. <https://doi.org/10.1002/cne.903500302>
- Lehman, J. F., Greenberg, B. D., McIntyre, C. C., Rasmussen, S. A., & Haber, S. N. (2011). Rules ventral prefrontal cortical axons use to reach their targets: Implications for diffusion tensor imaging tractography and deep brain stimulation for psychiatric illness. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 31(28), 10392–10402. <https://doi.org/10.1523/JNEUROSCI.0595-11.2011>
- Leichnetz, G. R., & Astruc, J. (1976). The efferent projections of the medial prefrontal cortex in the squirrel monkey (*Saimiri sciureus*). *Brain Research*, 109(3), 455–472. [https://doi.org/10.1016/0006-8993\(76\)90027-5](https://doi.org/10.1016/0006-8993(76)90027-5)
- MacLachy, L., Orlikoff, E., & Cosman, M. N. (2020). Hominoidea Morphology. In J. Vonk & T. Shackelford (Eds), *Encyclopedia of Animal Cognition and Behavior* (pp. 1–14). Springer International Publishing. [https://doi.org/10.1007/978-3-319-47829-6\\_2001-1](https://doi.org/10.1007/978-3-319-47829-6_2001-1)
- Mai, J. K., Majtanik, M., & NHMRC, G. P. A. (2015). *Atlas of the Human Brain* (4 edition). Academic Press.
- Maier-Hein, K. H., Neher, P. F., Houde, J.-C., Côté, M.-A., Garyfallidis, E., Zhong, J., Chamberland, M., Yeh, F.-C., Lin, Y.-C., Ji, Q., Reddick, W. E., Glass, J. O., Chen, D. Q., Feng, Y., Gao, C., Wu, Y., Ma, J., Renjie, H., Li, Q., ... Descoteaux, M. (2017). The challenge of mapping the human connectome based on diffusion tractography. *Nature Communications*, 8(1), 1349. <https://doi.org/10.1038/s41467-017-01285-x>
- Mars, R. B., Jbabdi, S., & Rushworth, M. F. S. (2021). A Common Space Approach to Comparative Neuroscience. *Annual Review of Neuroscience*, 44, 69–86. <https://doi.org/10.1146/annurev-neuro-100220-025942>
- Mars, R. B., Neubert, F.-X., Verhagen, L., Sallet, J., Miller, K. L., Dunbar, R. I. M., & Barton, R. A. (2014). Primate comparative neuroscience using magnetic resonance imaging: Promises and challenges. *Frontiers in Neuroscience*, 8, 298. <https://doi.org/10.3389/fnins.2014.00298>
- Mars, R. B., Sotiropoulos, S. N., Passingham, R. E., Sallet, J., Verhagen, L., Khrapitchev, A. A., Sibson, N., & Jbabdi, S. (2018). Whole brain comparative anatomy using connectivity blueprints. *eLife*, 7. <https://doi.org/10.7554/eLife.35237>
- Mayberg, H. (2001). Depression and frontal-subcortical circuits: Focus on prefrontal-limbic interactions. In *Frontal-subcortical circuits in psychiatric and neurological disorders* (pp. 177–206). The Guilford Press.
- Mayberg, H. S., Lozano, A. M., Voon, V., McNeely, H. E., Seminowicz, D., Hamani, C., Schwab, J. M., & Kennedy, S. H. (2005). Deep brain stimulation for treatment-resistant depression. *Neuron*, 45(5), 651–660. <https://doi.org/10.1016/j.neuron.2005.02.014>

- McNab, J. A., Jbabdi, S., Deoni, S. C. L., Douaud, G., Behrens, T. E. J., & Miller, K. L. (2009). High resolution diffusion-weighted imaging in fixed human brain using diffusion-weighted steady state free precession. *NeuroImage*, 46(3), 775–785. <https://doi.org/10.1016/j.neuroimage.2009.01.008>
- McNab, J. A., & Miller, K. L. (2010). Steady-state diffusion-weighted imaging: Theory, acquisition and analysis. *NMR in Biomedicine*, 23(7), 781–793. <https://doi.org/10.1002/nbm.1509>
- Milham, M. P., Ai, L., Koo, B., Xu, T., Amiez, C., Balezeau, F., Baxter, M. G., Blezer, E. L. A., Brochier, T., Chen, A., Croxson, P. L., Damatac, C. G., Dehaene, S., Everling, S., Fair, D. A., Fleysher, L., Freiwald, W., Froudast-Walsh, S., Griffiths, T. D., ... Schroeder, C. E. (2018). An Open Resource for Non-human Primate Imaging. *Neuron*, 100(1), 61-74.e2. <https://doi.org/10.1016/j.neuron.2018.08.039>
- Munuera, J., Rigotti, M., & Salzman, C. D. (2018). Shared neural coding for social hierarchy and reward value in primate amygdala. *Nature Neuroscience*, 21(3), 415–423. <https://doi.org/10.1038/s41593-018-0082-8>
- Murray, E. A., & Wise, S. P. (2010). Interactions between orbital prefrontal cortex and amygdala: Advanced cognition, learned responses and instinctive behaviors. *Current Opinion in Neurobiology*, 20(2), 212–220. <https://doi.org/10.1016/j.conb.2010.02.001>
- Murray, E. A., Wise, S. P., & Rhodes, S. E. V. (2011). What Can Different Brains Do with Reward? In J. A. Gottfried (Ed.), *Neurobiology of Sensation and Reward*. CRC Press/Taylor & Francis. <http://www.ncbi.nlm.nih.gov/books/NBK92798/>
- Nauta, W. J. (1961). Fibre degeneration following lesions of the amygdaloid complex in the monkey. *Journal of Anatomy*, 95, 515–531.
- Nauta, W. J. (1962). Neural associations of the amygdaloid complex in the monkey. *Brain: A Journal of Neurology*, 85, 505–520. <https://doi.org/10.1093/brain/85.3.505>
- Ni, X., Flynn, J. J., Wyss, A. R., & Zhang, C. (2019). Cranial endocast of a stem platyrrhine primate and ancestral brain conditions in anthropoids. *Science Advances*, 5(8), eaav7913. <https://doi.org/10.1126/sciadv.aav7913>
- Nieuwenhuys, R., Voogd, J., & Huijzen, C. van. (2008). *The Human Central Nervous System: A Synopsis and Atlas* (4th edn). Steinkopff-Verlag Heidelberg. <https://www.springer.com/gb/book/9783540346845>
- Oler, J. A., Tromp, D. P. M., Fox, A. S., Kovner, R., Davidson, R. J., Alexander, A. L., McFarlin, D. R., Birn, R. M., Berg, B., deCampo, D. M., Kalin, N. H., & Fudge, J. L. (2017). Connectivity between the central nucleus of the amygdala and the bed nucleus of the stria terminalis in the non-human primate: Neuronal tract tracing and developmental neuroimaging studies. *Brain Structure & Function*, 222(1), 21–39. <https://doi.org/10.1007/s00429-016-1198-9>
- Passingham, R. E., & Smaers, J. B. (2014). Is the prefrontal cortex especially enlarged in the human brain allometric relations and remapping factors. *Brain, Behavior and Evolution*, 84(2), 156–166. <https://doi.org/10.1159/000365183>
- Passingham, R. E., & Wise, S. P. (2012). *The neurobiology of the prefrontal cortex: Anatomy, evolution, and the origin of insight* (1st ed). Oxford University Press.
- Paxinos, G., Huang, X. F., & Toga, A. W. (2000). *The Rhesus Monkey Brain in Stereotaxic Coordinates*. Academic Press.
- Preuss, T. M., & Kaas, J. H. (2003). The discovery of cerebral diversity: An unwelcome scientific revolution. In *Evolutionary anatomy of the primate cerebral cortex* (pp. 1147–1166). Cambridge University Press.
- Preuss, T. M., & Wise, S. P. (2022). Evolution of prefrontal cortex. *Neuropsychopharmacology*, 47(1), Article 1. <https://doi.org/10.1038/s41386-021-01076-5>

- Riva-Posse, P., Choi, K. S., Holtzheimer, P. E., McIntyre, C. C., Gross, R. E., Chaturvedi, A., Crowell, A. L., Garlow, S. J., Rajendra, J. K., & Mayberg, H. S. (2014). Defining Critical White Matter Pathways Mediating Successful Subcallosal Cingulate Deep Brain Stimulation for Treatment-Resistant Depression. *Biological Psychiatry*, 76(12), 963–969. <https://doi.org/10.1016/j.biopsych.2014.03.029>
- Roumazeilles, L., Eichert, N., Bryant, K. L., Folloni, D., Sallet, J., Vijayakumar, S., Foxley, S., Tendler, B. C., Jbabdi, S., Reveley, C., Verhagen, L., Dershowitz, L. B., Guthrie, M., Flach, E., Miller, K. L., & Mars, R. B. (2020). Longitudinal connections and the organization of the temporal cortex in macaques, great apes, and humans. *PLoS Biology*, 18(7), e3000810. <https://doi.org/10.1371/journal.pbio.3000810>
- Roumazeilles, L., Lange, F. J., Benn, R. A., Andersson, J. L. R., Bertelsen, M. F., Manger, P. R., Flach, E., Khrapitchev, A. A., Bryant, K. L., Sallet, J., & Mars, R. B. (2022). Cortical Morphology and White Matter Tractography of Three Phylogenetically Distant Primates: Evidence for a Simian Elaboration. *Cerebral Cortex (New York, N.Y.: 1991)*, 32(8), 1608–1624. <https://doi.org/10.1093/cercor/bhab285>
- Royo, J., Forkel, S. J., Pouget, P., & Thiebaut de Schotten, M. (2021). The squirrel monkey model in clinical neuroscience. *Neuroscience and Biobehavioral Reviews*, 128, 152–164. <https://doi.org/10.1016/j.neubiorev.2021.06.006>
- Rudebeck, P. H., Putnam, P. T., Daniels, T. E., Yang, T., Mitz, A. R., Rhodes, S. E. V., & Murray, E. A. (2014). A role for primate subgenual cingulate cortex in sustaining autonomic arousal. *Proceedings of the National Academy of Sciences of the United States of America*, 111(14), 5391–5396. <https://doi.org/10.1073/pnas.1317695111>
- Saez, A., Rigotti, M., Ostojic, S., Fusi, S., & Salzman, C. D. (2015). Abstract Context Representations in Primate Amygdala and Prefrontal Cortex. *Neuron*, 87(4), 869–881. <https://doi.org/10.1016/j.neuron.2015.07.024>
- Sallet, J., Mars, R. B., Noonan, M. P., Neubert, F.-X., Jbabdi, S., O'Reilly, J. X., Filippini, N., Thomas, A. G., & Rushworth, M. F. (2013). The Organization of Dorsal Frontal Cortex in Humans and Macaques. *Journal of Neuroscience*, 33(30), 12255–12274. <https://doi.org/10.1523/JNEUROSCI.5108-12.2013>
- Schmahmann, J. D., & Pandya, D. N. (2006). *Fiber Pathways of the Brain*. Oxford University Press.
- Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E. J., Johansen-Berg, H., Bannister, P. R., De Luca, M., Drobnjak, I., Flitney, D. E., Niazy, R. K., Saunders, J., Vickers, J., Zhang, Y., De Stefano, N., Brady, J. M., & Matthews, P. M. (2004). Advances in functional and structural MR image analysis and implementation as FSL. *NeuroImage*, 23 Suppl 1, S208–219. <https://doi.org/10.1016/j.neuroimage.2004.07.051>
- Sotiropoulos, S. N., Jbabdi, S., Xu, J., Andersson, J. L., Moeller, S., Auerbach, E. J., Glasser, M. F., Hernandez, M., Sapiro, G., Jenkinson, M., Feinberg, D. A., Yacoub, E., Lenglet, C., Van Essen, D. C., Ugurbil, K., Behrens, T. E. J., & WU-Minn HCP Consortium. (2013). Advances in diffusion MRI acquisition and processing in the Human Connectome Project. *NeuroImage*, 80, 125–143. <https://doi.org/10.1016/j.neuroimage.2013.05.057>
- Suarez, L. E., Yovel, Y., van den Heuvel, M. P., Sporns, O., Assaf, Y., Lajoie, G., & Misic, B. (2022). A connectomics-based taxonomy of mammals. *eLife*, 11, e78635. <https://doi.org/10.7554/eLife.78635>
- Tendler, B. C., Hanayik, T., Ansorge, O., Bangerter-Christensen, S., Berns, G. S., Bertelsen, M. F., Bryant, K. L., Foxley, S., Martijn van den Heuvel, Prof., Howard, A. F., Huszar, I. N., Khrapitchev, A. A., Leonte, A., Manger, P. R., Menke, R. A., Mollink, J., Mortimer, D., Pallebage-Gamarallage, M., Roumazeilles, L., ... Miller, K. L. (2022).

- The Digital Brain Bank, an open access platform for post-mortem datasets. *eLife*, 11, e73153. <https://doi.org/10.7554/eLife.73153>
- Thiebaut de Schotten, M., Dell'Acqua, F., Valabregue, R., & Catani, M. (2012). Monkey to human comparative anatomy of the frontal lobe association tracts. *Cortex; a Journal Devoted to the Study of the Nervous System and Behavior*, 48(1), 82–96. <https://doi.org/10.1016/j.cortex.2011.10.001>
- Uğurbil, K., Xu, J., Auerbach, E. J., Moeller, S., Vu, A. T., Duarte-Carvajalino, J. M., Lenglet, C., Wu, X., Schmitter, S., Van de Moortele, P. F., Strupp, J., Sapiro, G., De Martino, F., Wang, D., Harel, N., Garwood, M., Chen, L., Feinberg, D. A., Smith, S. M., ... WU-Minn HCP Consortium. (2013). Pushing spatial and temporal resolution for functional and diffusion MRI in the Human Connectome Project. *NeuroImage*, 80, 80–104. <https://doi.org/10.1016/j.neuroimage.2013.05.012>
- Van Essen, D. C., Smith, S. M., Barch, D. M., Behrens, T. E. J., Yacoub, E., Ugurbil, K., & WU-Minn HCP Consortium. (2013). The WU-Minn Human Connectome Project: An overview. *NeuroImage*, 80, 62–79. <https://doi.org/10.1016/j.neuroimage.2013.05.041>
- Verhagen, L., Gallea, C., Folloni, D., Constans, C., Jensen, D. E., Ahnine, H., Roumazeilles, L., Santin, M., Ahmed, B., Lehericy, S., Klein-Flügge, M. C., Krug, K., Mars, R. B., Rushworth, M. F., Pouget, P., Aubry, J.-F., & Sallet, J. (2019). Offline impact of transcranial focused ultrasound on cortical activation in primates. *eLife*, 8. <https://doi.org/10.7554/eLife.40541>
- Vogt, B. A. (2005). Pain and emotion interactions in subregions of the cingulate gyrus. *Nature Reviews. Neuroscience*, 6(7), 533–544. <https://doi.org/10.1038/nrn1704>
- Warrington, S., Bryant, K. L., Khrapitchev, A. A., Sallet, J., Charquero-Ballester, M., Douaud, G., Jbabdi, S., Mars, R. B., & Sotiropoulos, S. N. (2020). XTRACT - Standardised protocols for automated tractography in the human and macaque brain. *NeuroImage*, 217, 116923. <https://doi.org/10.1016/j.neuroimage.2020.116923>
- Warrington, S., Thompson, E., Bastiani, M., Dubois, J., Baxter, L., Slater, R., Jbabdi, S., Mars, R. B., & Sotiropoulos, S. N. (2022). Concurrent mapping of brain ontogeny and phylogeny within a common space: Standardized tractography and applications. *Science Advances*, 8(42), eabq2022. <https://doi.org/10.1126/sciadv.abq2022>
- Wittmann, M. K., Fouragnan, E., Folloni, D., Klein-Flügge, M. C., Chau, B. K. H., Khamassi, M., & Rushworth, M. F. S. (2020). Global reward state affects learning and activity in raphe nucleus and anterior insula in monkeys. *Nature Communications*, 11. <https://doi.org/10.1038/s41467-020-17343-w>
- Wood, C. M., Alexander, L., Alsö, J., Santangelo, A. M., McIver, L., Cockcroft, G. J., & Roberts, A. C. (2023). Chemogenetics identifies separate area 25 brain circuits involved in anhedonia and anxiety in marmosets. *Science Translational Medicine*, 15(690), eade1779. <https://doi.org/10.1126/scitranslmed.ade1779>
- Wright, A. K., Theilmann, R. J., Ridgway, S. H., & Scadeng, M. (2018). Diffusion tractography reveals pervasive asymmetry of cerebral white matter tracts in the bottlenose dolphin (*Tursiops truncatus*). *Brain Structure and Function*, 223(4), 1697–1711. <https://doi.org/10.1007/s00429-017-1525-9>
- Yarnykh, V. L. (2007). Actual flip-angle imaging in the pulsed steady state: A method for rapid three-dimensional mapping of the transmitted radiofrequency field. *Magnetic Resonance in Medicine*, 57(1), 192–200. <https://doi.org/10.1002/mrm.21120>

**Figure 1. Primate family tree.** Phylogram showing evolutionary relationships of prosimians, Old and New World monkeys, lesser and great apes, and humans (Eldridge et al., 2021; MacLatchy et al., 2020; Ni et al., 2019; Preuss & Wise, 2022). Brains are not to scale. Mya = million years ago.

**Figure 2.** Limbic-frontal connections in macaque monkeys and humans. The anterior limb of the AmF (red-yellow), UF (blue-light blue) and CB (green) are displayed across the two species on five coronal brain slices (A-B) and three axial brain (C-D) slices showing their respective interactions and organization in limbic and prefrontal cortex. Location of slices from A-D are reported in E for the human brain (top) and macaque monkey brain (bottom).

**Figure 3. Histology-diffusion MRI comparison of connections.** Tract-tracing-driven (left panel; reconstructed from Figure 15 in Heilbronner and Haber(Heilbronner & Haber, 2014) reconstruction of amygdala-cingulate-prefrontal connections using non-invasive diffusion MRI (left and right) in macaque monkeys (right hemisphere) and humans (left hemisphere). In both macaque and human brains AmF (red-yellow), UF (blue-light blue) and CB (green) are shown.

**Figure 4. Coronal views of temporal-frontal connections in all species.** The anterior limb of the AmF (red-yellow), UF (blue-light blue) are displayed across each species on three coronal brain slices showing their respective interactions and organization along a posterior (leftmost column) – anterior (rightmost column) gradient. The brains are displayed from the species most closely related to humans (top row) to the least related (Ring-tailed lemur, bottom row).


**Figure 5. Organization of temporal-cingulate-frontal connections across all species.** The anterior limb of the AmF (red-yellow), UF (blue-light blue) and CB (green) are displayed across each species on a sagittal brain slice showing their organization within cingulate and prefrontal cortices, striatum and basal forebrain. The brains are displayed in the following order from left to right: ring-tailed lemur (prosimians), squirrel monkey (New World monkey), macaque monkey (Old World monkey), gibbon (lesser ape), gorilla (great ape), chimpanzee (great ape) and human (*Homo sapiens sapiens*).


**Figure 6. Evolutionary expansion of cingulate projections into the subcallosal cortex across primates.** Extension of cingulum bundle (green), amygdalofugal pathway (red-yellow) and uncinate fasciculus (blue) into the subcallosal part of the anterior cingulate cortex from prosimians to humans. Shown from left to right are representative coronal slices from the lemur, squirrel monkey, macaque monkey, gibbon, gorilla, chimpanzee, and human, aligned to emphasise homologous subcallosal landmarks. Cingulum fibres can be observed in all primates from the macaque monkey to the Human brains but not in lemurs and new-world squirrel monkeys. Organization and extension of Amygdalofugal and uncinate fascicle are preserved across all species.


**Figure 7. Principles of organization and prefrontal targets across strepsirrhini and haplorhini.** Organization and prefrontal targets of cingulum bundle (green), amygdalofugal pathway (red-yellow) and uncinate fascicle (blue) across strepsirrhini (lemurs. left) and haplorrhine (macaque monkeys and humans, right). Amygdalofugal pathway and uncinate fasciculus organization is similar across the two suborders or primates, with separation in


more caudal regions and convergence in the anterior frontal cortex. Extension of the Cingulum bundle fibres in the subcallosal cortex, however, differs between the two suborders.


**Figure 8. Schematic diagram of similarities and differences across strepsirrhini and haplorhini in the limbic-prefrontal connectome.** Schematic template representing the organization of AmF (red), UF (blue) and CB (green) in the ventral prefrontal cortex in the lemur (left), monkey (middle) and human brains displayed on an axial (top row) and sagittal slice (bottom row). The dashed line (red) shows the subcortical AmF fibers running underneath and through the ventral striatum.











## Extension of cingulate fibers in the subcallosal cortex across primates

